Extensions of the multicentric functional calculus

Diana Andrei

Extensions of the multicentric functional calculus

Diana Andrei

A doctoral dissertation completed for the degree of Doctor of Science (Technology) to be defended, with the permission of the Aalto University School of Science, at a public examination held at the lecture hall M1 of the school on 10 December at 12 noon.

Aalto University
School of Science
Department of Mathematics and Systems Analysis

Supervising professor

Professor Juha Kinnunen, Aalto University, Finland

Thesis advisor

Emeriti Professor Olavi Nevanlinna, Aalto University, Finland

Preliminary examiners

Professor Mikael Lindström, Åbo Akademi University, Finland

University Lecturer Hans-Olav Tylli, University of Helsinki, Finland

Opponent

University Lecturer Dr. Jari Taskinen, University of Helsinki, Finland

Aalto University publication series

DOCTORAL DISSERTATIONS 174/2021

© 2021 Diana Andrei

ISBN 978-952-64-0624-4 (printed)

ISBN 978-952-64-0625-1 (pdf)

ISSN 1799-4934 (printed)

ISSN 1799-4942 (pdf)

<http://urn.fi/URN:ISBN:978-952-64-0625-1>

Unigrafia Oy

Helsinki 2021

Finland

Author

Diana Andrei

Name of the doctoral dissertation

Extensions of the multicentric functional calculus

Publisher School of Science**Unit** Department of Mathematics and Systems Analysis**Series** Aalto University publication series DOCTORAL DISSERTATIONS 174/2021**Field of research** Mathematics**Manuscript submitted** 17 November 2021**Date of the defence** 10 December 2021**Permission for public defence granted (date)** 16 November 2021**Language** English☐ **Monograph**☒ **Article dissertation**☐ **Essay dissertation****Abstract**

In operator theory, one of the central concepts is the spectrum of an operator and if one knows how to separate the spectrum into components, then the multicentric calculus is a useful tool, introduced by Olavi Nevanlinna in 2011.

This thesis presents extensions of the multicentric calculus from single operators to n -tuples of commuting operators, for both holomorphic and non-holomorphic functions. It also covers the same calculus when replacing the polynomials with rational functions.

The multicentric representation of holomorphic functions gives a simple way to generalize the von Neumann result, i.e., the unit disc is a spectral set for contractions in Hilbert spaces. In other words, this calculus provides a way of representing the spectrum of a bounded operator T , by searching for a polynomial p that maps the spectrum to a small disc around origin. Since the von Neumann inequality works for contractions with spectrum in the unit disc, the multicentric representation applies a suitable polynomial p to the operator T , so that $p(T)$ becomes a contraction with spectrum in the unit disc and thus the usual holomorphic functional calculus holds. When extending the calculus to n -tuples of commuting operators, a constant and some extra conditions are needed for the von Neumann inequality to hold true.

In order to extend the calculus to non-holomorphic functions, the Banach algebra is the tool to use in finding those functions for which it is possible to have a simple functional calculus by using suitable polynomial p . For a given bounded operator T on a Hilbert space, the polynomial p is such that $p(T)$ is diagonalizable or similar to normal. The operators here are considered to be matrices. In particular, the calculus provides a natural approach to deal with non-trivial Jordan blocks. The extension of this calculus is done for a pair of commuting matrices, by constructing a suitable tensor product Banach algebra that can be identified with the space of continuous functions of two variables.

When replacing the polynomial by a rational function, one can apply the calculus to functions that are not polynomially convex, thus extending it to a larger class of functions.

Keywords multicentric calculus, lemniscates, spectral projection, commuting operators, Banach algebra, tensor norm, rational functions, series expansions.

ISBN (printed) 978-952-64-0624-4**ISBN (pdf)** 978-952-64-0625-1**ISSN (printed)** 1799-4934**ISSN (pdf)** 1799-4942**Location of publisher** Helsinki**Location of printing** Helsinki **Year** 2021**Pages** 124**urn** <http://urn.fi/URN:ISBN:978-952-64-0625-1>

Preface

I want to express my gratitude to several people who made this dissertation possible.

First, I want to thank my thesis advisor, Professor Olavi Nevanlinna, who took me into his research group and acted as my supervisor for my first years into the doctoral program. I am deeply grateful for his guidance of superior quality, his advice and ideas that he gave me over the years, as well as for his patience and unconditional support. I also want to thank Professor Juha Kinnunen for supervising my last years into the program and for the opportunity to do this work and my co-author, Tiina Vesanen, for her valuable work and contributions. I thank my preliminary examiners, for their valuable comments.

Then my gratitude goes to everyone at Jenny and Wihuri Foundation who partly supported my research and thesis completion.

Finally, I want to thank my family, my parents and sister for their trust and support. My last words go to my husband Cristian. Thank you for your loving pressure to finish.

Espoo, November 17, 2021,

Diana Andrei

Contents

Preface	1
Contents	3
List of Publications	5
Author's Contribution	7
1. Introduction	9
1.1 History and background	9
1.2 Organization of the thesis	10
2. Multicentric Calculus	11
2.1 Basis of multicentric calculus	11
2.2 Application to Riesz projection	14
2.3 Separation of the spectrum	16
3. Extension of multicentric holomorphic calculus to multiple commuting operators	19
3.1 On a pair of commuting operators	19
3.2 On the n -tuples of commuting operators	21
4. A Banach algebra with a multicentric calculus	25
4.1 Construction of the algebra	25
4.2 Application to the Gelfand theory	29
5. Change of variables using rational functions	31
5.1 Representation using rational functions as variables . . .	31
5.2 Applications and remarks	33
References	35
Publications	37

List of Publications

This thesis consists of an overview and of the following publications which are referred to in the text by their Roman numerals.

- I** Diana Apetrei, Olavi Nevanlinna. Multicentric calculus and the Riesz projection. *Journal of Numerical Analysis and Approximation Theory*, Volume 44, no.2, pp. 127-145, 2015.
- II** Diana Andrei. Multicentric holomorphic calculus for n -tuples of commuting operators. *Advances in Operator Theory*, Volume 4, no.2, pp. 447-461, 2019.
- III** Diana Andrei. A tensor product algebra with functional calculus for commuting pair of matrices. Submitted to *Advances in Operator Theory*, September 22th 2021.
- IV** Diana Andrei, Olavi Nevanlinna, Tiina Vesanen. Rational functions as new variables. Submitted to *Banach Journal of Mathematical Analysis*, May 24th 2021.

Author's Contribution

Publication I: “Multicentric calculus and the Riesz projection”

Diana Apetrei proved the model problem, did the numerical estimates and wrote that up. Olavi Nevanlinna proposed the problem and wrote the introduction and concluding remarks.

Publication II: “Multicentric holomorphic calculus for n -tuples of commuting operators”

This represents independent work by the author.

Publication III: “A tensor product algebra with functional calculus for commuting pair of matrices”

This represents independent work by the author.

Publication IV: “Rational functions as new variables”

Diana Andrei and Olavi Nevanlinna came up with the proposed problem after long collaboration time and worked out the key results equally. Olavi Nevanlinna did the writing. Tiina Vesanen did the numerical estimates and visualizations.

1. Introduction

1.1 History and background

The base for the multicentric holomorphic representation is the Jacobi series. This series was introduced by C.G.J. Jacobi [10] in the 19th century, without the use of Cauchy integral. Jacobi was interested in extending functions using polynomials as powers and using one degree lower polynomials as coefficients. Cauchy introduced the Cauchy integral [3] and function theory earlier, but for some reason Jacobi did not use it at all. Jacobi's topic had not been much discussed before Alfred Kienast [11] in 1906 when, in his dissertation, he looked at the Jacobi series using the function theory, how it fits in using the Cauchy integral but did not take polynomial as a new variable. One reason why it looked complicated and not attractive was that things had to be separately worked out in each mathematical domain.

Again, going back to the 19th century, orthogonal polynomials were created earlier than the Jacobi series and Jacobi worked with them. In particular, orthogonal polynomial numerically were attractive when looking at interval in real line and whenever one could move the discussion into that environment. On the other hand, because of using orthogonal polynomials and restricting attention to spectra in real line, that is, working with self-adjoint and normal operators, the tools that worked in some way for such operators were further developed, while for non-normal operators all the work assumed a particular type.

Later, many studied and developed analytic functions in the disc, what behaviour they had when one approached the boundary. It is known that, if a function is analytic inside the disc and continuous up to boundary, then polynomials are dense, otherwise things are tricky. M. Fekete [8], in 1932, touched this by looking at the Jacobi series inside a lemniscate, but without mapping the discussion back to unit circle. Most likely the interest from pure mathematics was not there.

In 1941, at the suggestion of professor J.L. Walsh, J.H Curtiss [5] studied the behaviour of the Jacobi series on the boundary of its regions of convergence.

Although conformal maps became available and were well understood, the multicentric holomorphic calculus was introduced by Olavi Nevanlinna [19] in the 21st century. His interest started when he studied polynomial numerical hulls, then operator valued meromorphic functions and in particular low rank perturbation theory. It then became natural for him to ask whether one could compute the spectrum and if the resolvent could be represented by a single expression using holomorphic functional calculus. The power of this multicentric calculus given functional analysis calculus is that it doesn't depend on the normality of the operators.

This thesis will cover the basis of multicentric functional calculus for both holomorphic and non-holomorphic functions and then presents my work in extending it to multiple variables. One further development for the single variable case is presented at the end of this paper, this time using rational functions instead of polynomials as new variables.

1.2 Organization of the thesis

Each section describes a publication in chronological order. The first section, Section 2, includes the basic notions and background for the multicentric calculus as presented in [17] - [19], that are used throughout the thesis.

Section 2 also covers Publication I where the authors look at powers of a polynomial instead of looking at its norm, in order to get spectral radius and how they were able to apply the multicentric holomorphic calculus to represent the Riesz spectral projection. Then spectrum separation issues by the lemniscate are presented through examples.

Section 3 describes Publication II where the author extends the single variable multicentric holomorphic calculus to n -tuples of commuting variables starting with the case when $n = 2$. Then the variables are replaced by operators and formulas are worked out accordingly.

Section 4 covers Publication III which extends even further the multicentric calculus in the multivariable setup. Here the author follows the non-holomorphic case presented in [20] for single variable. To that end an algebra is defined and basic notions and background are presented at the beginning of the section.

Section 5 presents the last publication, Publication IV, in which the authors look at a different change of variables, replacing the monic polynomial by a rational function.

Each section explains the main results and also points out challenges and possible constraints used to meet the desired goals.

2. Multicentric Calculus

2.1 Basis of multicentric calculus

This section gathers all basic notions and results from [17], [18], and [19]. In [18], O. Nevanlinna shows how the multicentric representation of functions gives a simple way to generalize the von Neumann result, i.e., the unit disc is a spectral set for contractions in Hilbert spaces. The discussion in his paper regards the multicentric representation of holomorphic functions and briefly it goes as follows.

Let $\mathcal{A}$ be a unital Banach algebra. Recall that a set $\Sigma \subset \mathbb{C}$ is a K -spectral set for $a \in \mathcal{A}$, if for all rational functions r with poles off Σ one has that $\|r(a)\| \leq K \sup_{z \in \Sigma} |r(z)|$, with a fixed K .

If A denotes a bounded operator in a Hilbert space H , denote by $V_p(A)$ the set

$$V_p(A) = \{z \in \mathbb{C} : |p(z)| \leq \|p(A)\|\} \quad (2.1.1)$$

where p is a monic polynomial with distinct roots, $\lambda_1, \dots, \lambda_d$. Denote by Λ the set $\{\lambda_1, \dots, \lambda_d\}$. Then these sets are K -spectral, whenever the lemniscate, that is, the boundary of $V_p(A)$, $\{z \in \mathbb{C} : |p(z)| = \|p(A)\|\}$, does not pass through any critical point of p . Since any compact set can be the spectrum of an operator, it is important that lemniscates have good approximation properties. Furthermore, in [17] is given an algorithm which, for any given A , provides a sequence of monic polynomials p with distinct roots so that the sets $V_p(A)$ squeeze around the polynomially convex hull of the spectrum of A .

To be able to estimate a holomorphic function φ effectively at an operator A , it is used the approach introduced in [19], that is, each polynomial

$$p(z) = \prod_{k=1}^d (z - \lambda_k)$$

with simple roots λ_k induces a unique multicentric representation of φ ,

$$\varphi(z) = \sum_{j=1}^d \delta_j(z) f_j(w) \quad \text{with } w = p(z), \quad (2.1.2)$$

where δ_j denote the polynomials of degree $d - 1$ taking the value 1 at λ_j and vanishes at the other roots. In [19] the practical computation of the Taylor series of f_j is discussed. In fact, the coefficients may be computed in a recursive way if the derivatives of the original function φ are available at the local centers λ_j .

The representation (2.1.2) allows an obvious avenue for analysis, estimation and computation in complicated sets. One just treats the functions f_j in discs $|w| \leq \rho$ and combines the estimates for φ in the sets satisfying $|p(z)| \leq \rho$.

This approach is demonstrated in [18] by generalizing a well-known result of von Neumann on contractions in Hilbert spaces. In order to do this one needs an estimate of the following form

$$\sup_{|w| \leq \rho} |f_j(w)| \leq C(\rho) \sup_{|p(z)| \leq \rho} |\varphi(z)|. \quad (2.1.3)$$

This would then imply that the sets $V_p(A)$ are K -spectral sets with some K . Now, let γ_ρ denote the lemniscate

$$\gamma_\rho = \{z \in \mathbb{C} : |p(z)| = \rho\}.$$

For small ρ the lemniscate consists of d separate circular curves, for large ρ it reduces to just one circular curve. In general the lemniscate is smooth except if it contains a critical point, where the derivative of p vanishes. Thus there are at most $d - 1$ such exceptional values ρ . Let $s(\rho)$ denote the distance from γ_ρ to the set of critical points.

The key result here is the following theorem.

Theorem 2.1.1. *If p is a monic polynomial of degree d with distinct roots, then there exists a constant C such that if φ is holomorphic for $|p(z)| \leq \rho$, then the functions f_k in (2.1.2) are holomorphic for $|w| \leq \rho$ and if γ_ρ does not contain any critical points of p the estimate (2.1.3) holds with some $C(\rho)$ satisfying*

$$C(\rho) \leq 1 + \frac{C}{s(\rho)^{d-1}}. \quad (2.1.4)$$

Remark 2.1.2. If $C(\rho)$ denotes the smallest constant such that (2.1.3) holds for all φ then $C(\rho) \rightarrow 1$ as $\rho \rightarrow 0$ or $\rho \rightarrow \infty$. Generically the critical points are simple and then the constant is proportional to $1/s(\rho)$.

Below we state a fundamental result by von Neumann for contractions in Hilbert spaces.

Theorem 2.1.3. *If $A \in \mathcal{B}(H)$, and $\|A\| \leq 1$, then the closed unit disc is a spectral set for A .*

This can be reformulated as follows:

$$\|\varphi(A)\| \leq \sup_{|z| \leq \|A\|} |\varphi(z)| \quad (2.1.5)$$

provided φ is holomorphic in $|z| \leq \|A\|$.

The results are formulated for holomorphic functions rather than for polynomials or rational functions as there are considered sets which may consist of several simply connected components. In particular, the results apply as such for Riesz spectral projections. The main result in [18] is the following:

Theorem 2.1.4. *Suppose we are given a monic polynomial $p \in \mathbb{P}_d$ with distinct roots and a bounded operator $A \in \mathcal{B}(H)$ in a Hilbert space H . Let $\rho \geq 0$ satisfying $\|p(A)\| \leq \rho$ and be such that the lemniscate γ_ρ contains no critical points of p . Then for all φ which are holomorphic for $|p(z)| \leq \rho$ there holds*

$$\|\varphi(A)\| \leq K \sup_{|p(z)| \leq \rho} |\varphi(z)|, \quad (2.1.6)$$

where the constant K satisfies

$$K \leq C(\rho) \sum_{j=1}^d \|\delta_j(A)\|, \quad (2.1.7)$$

with $C(\rho)$ as in Theorem 2.1.1.

A simple but useful application of this theorem is the Riesz spectral projection. Suppose γ_ρ consists of several components and is free from critical points. Then one can define φ to be identically 1 in some open neighbourhood of some of the components and to vanish in a neighbourhood of all the others. If $A \in \mathcal{B}(H)$ is such that $\|p(A)\| \leq \rho$, then the resulting operator is simply the Riesz spectral projection to the invariant subspace with respect to the part of the spectrum where φ equals 1.

Another application is related to the power boundedness. One can apply Theorem 2.1.4 with $\rho = 0$ but then A has to be an algebraic operator and all eigenvalues are non-defective. Requiring $p(A) = 0$ for p with simple zeroes says exactly that. Now let $\mathbb{D}$ denote the open unit disc. Suppose that $V_p(A) \subset \overline{\mathbb{D}}$.

Corollary 2.1.5. *Let p be monic with simple zeros and suppose $\rho \geq 0$ is such that γ_ρ contains no critical points and $\gamma_\rho \subset \mathbb{D}$. If $A \in \mathcal{B}(H)$ is such that $\|p(A)\| \leq \rho$, then A is power bounded and with the constant $C(\rho)$ provided by Theorem 2.1.1 we have for all $n \geq 1$,*

$$\|A^n\| \leq C(\rho) \sum_{j=1}^d \|\delta_j(A)\|. \quad (2.1.8)$$

Remark 2.1.6. If A is algebraic then one can take p to be the minimal polynomial. Recall that a polynomial is called minimal if it is monic, $p(A) = 0$ and the polynomial is of smallest possible degree. Then Corollary 2.1.5 and Remark 2.1.2 yield

$$\|A^n\| \leq \sum_{j=1}^d \|\delta_j(A)\|. \quad (2.1.9)$$

Remark 2.1.7. In his papers Olavi Nevanlinna made the convention that a function is holomorphic in a compact set if it is holomorphic in an open neighbourhood containing that compact set.

2.2 Application to Riesz projection

In Publication I one can find a follow up on the papers of O. Nevanlinna [18], [17] and [19]. There we discuss about multicentric holomorphic calculus in which one represents the function φ using a new variable $w = p(z)$ in such a way that when it is evaluated at the operator A , then $p(A)$ is small in norm. As before, p is assumed to have distinct roots.

The separation of a compact set, such as the spectrum, into different components by a polynomial lemniscate is presented with few examples, as well as applications of the Calculus to the computation and estimation of the Riesz spectral projection. It may then be desirable the use of $p(z)^n$ as a new variable and a small dimensional problem is also presented to see how the size of coupling can affect on the need of taking a high power of $p(A)$.

To apply the multicentric calculus to the Riesz projection, when getting the spectral radius it is natural to look at the power of p instead of looking at the norm. For this we need the basic formula for expressing a given function $\varphi(z)$ as a linear combination of functions $f_{j,k}(w^n)$ when $w = p(z)$, for n a given positive integer. The function f_j in the multicentric representation (2.1.2) are obtained from φ with the formula from [18]

$$f_j(w) = \sum_{l=1}^d \delta_l(\lambda_j, w) \varphi(\zeta_l(w)), \quad (2.2.1)$$

where $\zeta_l(w)$ denote the roots of $p(\lambda) - w = 0$ and

$$\delta_l(\lambda, w) = \frac{p(\lambda) - w}{p'(\zeta_l(w))(\lambda - \zeta_l(w))}.$$

When φ is holomorphic, f_j can also be computed from the Taylor coefficients of φ at the local centers λ_j . So the computations for f_j 's are done with power series and it makes sense to move to $p(z)^n = w^n$, because the expansions are done for that variable. It is then proved the following result.

Theorem 2.2.1. *Suppose p has simple zeros and assume φ is holomorphic in a neighbourhood of $V(p, \rho) = \{z \in \mathbb{C} : |p(z)| \leq \rho\}$ and given in the multicentric representation (2.1.2). Then*

$$\varphi(z) = \sum_{j=1}^d \delta_j(z) [f_{j,0}(w^n) + \cdots + w^{n-1} f_{j,n-1}(w^n)],$$

where $f_{j,k}$ are holomorphic in a neighbourhood of the disc $|w| \leq \rho$ and given by

$$w^k f_{j,k}(w^n) = \frac{1}{n} \{f_j(w) + e^{-2\pi i k/n} f_j(e^{2\pi i/n} w) + \cdots + e^{-2\pi i(n-1)k/n} f_j(e^{2\pi i(n-1)/n} w)\}.$$

The proof uses the next result.

Proposition 2.2.2. *If f_j is given for $|p(z)| \leq \rho$, then $f_{j,k}$, $k = 1, \dots, n-1$, are defined for $|p(z)^n| \leq \rho^n$ and*

$$f_j(p(z)) = \sum_{k=0}^{n-1} p(z)^k f_{j,k}(p(z)^n), \quad \text{for } |p(z)| \leq \rho. \quad (2.2.2)$$

Further, if $f_j(p(z))$ is analytic for $|p(z)| \leq \rho$ then so are $f_{j,k}(p(z)^n)$.

Now let A be a bounded operator in a Hilbert space such that

$$V_{p^n}(A) = \{z : |p(z)^n| \leq \|p(A)^n\|\}.$$

To compute the Riesz projection we take $\varphi = 1$ in one of the components and $\varphi = 0$ in the other components of $V_{p^n}(A)$ and $f_j(p(z))$ as in (2.2.2).

We apply the fact that if φ is holomorphic in a neighbourhood of the unit disk $\mathbb{D}$ and $A \in \mathcal{B}(H)$, then

$$\|\varphi(A)\| \leq \sup_{\mathbb{D}} |\varphi|, \quad (2.2.3)$$

see e.g. [23].

We are then able to represent the Riesz projection with the help of multicentric representation.

Theorem 2.2.3. *Given a polynomial p of degree d , with distinct roots and a bounded operator A in a Hilbert space, we assume that the "expression"*

$$V_{p^n}(A) = \{z : |p(z)^n| \leq \|p(A)^n\|\}$$

has at least two components. Set $\rho = \|p(A)^n\|^{1/n}$ and let φ be a function such that $\varphi = 1$ in one component of $V_{p^n}(A)$ and $\varphi = 0$ in the others. Let s be the distance from the nearest outside critical point to the boundary of $V_{p^n}(A)$.

Then, considering $f_{j,k}$ as given by Theorem 2.2.1, one has

$$\varphi(A) = \sum_{j=1}^d \delta_j(A) \sum_{k=0}^{n-1} p(A)^k f_{j,k}(p(A)^n),$$

which is the Riesz spectral projection onto the invariant subspace corresponding to the spectrum inside the component where $\varphi = 1$.

The bound for the norm of $\varphi(A)$ is given by

$$\|\varphi(A)\| \leq \left[\left(1 + \frac{C}{s^{d-1}} \right) \frac{\|p(A)^k\|}{\rho^k} \sum_{j=1}^d \|\delta_j(A)\| \right] \sup_{|p(z)| \leq \rho} |\varphi(z)|. \quad (2.2.4)$$

The proof uses few other results and statements that are presented in details in Publication I.

Remark 2.2.4. In applications we will consider $\varphi = 1$ in the components of $V_{p^n}(A)$ that are on the right complex half-plane and $\varphi = -1$ in the components on the left complex half-plane. This way the computations are more symmetrical. In this situation there is only one critical point at the origin, which is simple, so we will have $\left(1 + \frac{C}{s} \right)$ in the formula for the bound of the Riesz projection.

2.3 Separation of the spectrum

The spectrum separation issues by the lemniscate is another topic explored in Publication I. Given a polynomial p denote by $V(p, \rho)$ the set

$$V(p, \rho) = \{z \in \mathbb{C} : |p(z)| \leq \rho\}.$$

Let $K \subset \mathbb{C}$ be compact and such that $\mathbb{C} \setminus K$ is connected. For $\delta > 0$ denote further

$$K(\delta) = \{z : \text{dist}(z, K) < \delta\}.$$

Then there exists a polynomial p and $\rho > 0$ such that

$$K \subset V(p, \rho) \subset K(\delta).$$

In particular, if $K = K_1 \cup K_2$ and $K_1 \cap K_2 = \emptyset$ since K is compact, then for small enough δ

$$K_1(\delta) \cap K_2(\delta) = \emptyset$$

as well. Thus, $V(p, \rho)$ separates the components K_1 and K_2 respectively.

If we assume that we have two analytic functions φ_j , each analytic in $K_j(\delta)$, we can view them as just one analytic function $\varphi : K(\delta) \rightarrow \mathbb{C}$, where φ agrees to φ_j on $K_j(\delta)$. We are interested in particular in the case where φ_j is constant. Multicentric representation then gives a power series which is simultaneously valid in both components.

For such a separation task, we can ask what is the smallest degree of a polynomial achieving this and, to answer this, we model the problem as follows: let

$$K_1(\delta) = \{z = 1 + iy : |y| \leq \tan(\delta)\}$$

and $K_2(\delta)$ symmetrically on the other side of the imaginary axis:

$$K_2(\delta) = \{z = -1 + iy : |y| \leq \tan(\delta)\}$$

Our first discussed problem in Publication I concerns the minimal degree of a polynomial p such that $K_j(\delta) \subset V(p, \rho)$ and $V(p, \rho) \cap i\mathbb{R} = \emptyset$.

Another natural task is related to existence of a logarithm. Let M be a compact and we assume $0 \notin M$. Now there exists a single valued logarithm in M if and only if 0 is in the unbounded component of the complement of M . That is, the set M does not separate origin from infinity. So the natural task here is to find a polynomial p such that $M \subset V(p, \rho)$ and $0 \notin V(p, \rho)$. As $V(p, \rho)$ is polynomially convex, this suffices for representing the logarithm in $V(p, \rho)$.

Remark 2.3.1. The separation by lemniscate is done by slightly perturbing the roots of the polynomial p and decreasing the level of the lemniscate to below 1 such that the spectrum lies inside the lemniscate. A few practical examples are worked out in Publication I and estimates for the smallest lemniscate level are given such that we get the desired separation for those particular cases.

3. Extension of multicentric holomorphic calculus to multiple commuting operators

In Publication II the multicentric holomorphic representation (2.1.2) is extended to n -tuple of commuting operators, first by working out the multicentric representation for two variables, then replacing the variables with a pair of commuting operators on a Hilbert space H , and finally moving to n variables and n -tuple of commuting operators.

Let $\mathbf{z} \in \mathbb{C}^n$ and let $\mathbf{p}$ be a vector of polynomials mapping

$$\mathbf{z} = \begin{pmatrix} z_1 \\ z_2 \\ \dots \\ z_n \end{pmatrix} \mapsto \begin{pmatrix} p_1(z_1) \\ p_2(z_2) \\ \dots \\ p_n(z_n) \end{pmatrix} = \begin{pmatrix} w_1 \\ w_2 \\ \dots \\ w_n \end{pmatrix} = \mathbf{w} \in \mathbb{C}^n,$$

where for every $j = 1, 2, \dots, n$, we have that $p_j(z_j)$ is a monic polynomial of degree d_j with distinct roots $\lambda_{1,j}, \lambda_{2,j}, \dots, \lambda_{d_j,j}$.

3.1 On a pair of commuting operators

We start with the particular case when $n = 2$. Recall that in multicentric holomorphic calculus one deals with polynomials p of level below 1, thus for $\mathbf{w} = (w_1, w_2) \in \mathbb{C}^2$ we have $|w_j| \leq 1$, for $j = 1, 2$, i.e. $\mathbb{D}^2 = \{\mathbf{w} \in \mathbb{C}^2 : |w_j| \leq 1, \text{ for } j = 1, 2\}$ is a unit bidisc. Using the change of variable $w_j = p_j(z_j)$ one works on unit discs which are simple sets, and then gets results on the rather more complicated sets

$$\Omega = \{\mathbf{z} = (z_1, z_2) \in \mathbb{C}^2 : z_j \in p_j^{-1}(\mathbb{D}^2), j = 1, 2\}.$$

According to [18] this multicentric representation of functions gives a simple way to generalize the von Neumann result, i.e. the unit disc is a spectral set for contractions in Hilbert spaces, or in other words, this calculus allows the preimage of the disc to be a K -spectral set.

Useful results in moving to operators are the spectral mapping theorem and the von Neumann inequality for a pair of commuting contractions. The

spectral mapping theorem states that if p is a polynomial with complex coefficients and $T \in \mathcal{B}(H)$ (space of bounded operators on a Hilbert space H), then $\sigma(p(T)) = p(\sigma(T))$.

When working with commuting n -tuples of Banach space operators the multicentric calculus is more involved. There are many possible definitions for a joint spectrum, but the one introduced by J.L. Taylor has the property that there exists a functional calculus for functions analytic on a neighbourhood of this spectrum.

Recall that the *Taylor spectrum* $\sigma_T(\mathbf{T})$ is the set of all $\lambda = (\lambda_1, \dots, \lambda_n) \in \mathbb{C}^n$ such that the n -tuple $\mathbf{T} - \lambda = (T_1 - \lambda_1, \dots, T_n - \lambda_n)$ is not Taylor invertible, [14].

Theorem 3.1.1. (Spectral Mapping Property). *Let $\mathbf{T} = (T_1, \dots, T_n) \in \mathcal{B}(H)^n$ be a commuting n -tuple of operators and let p be a polynomial in n variables. Then*

$$\sigma_T(p(\mathbf{T})) = p(\sigma_T(\mathbf{T})).$$

This is a special case of Theorem 2 in [15] where a vector-valued polynomial p is used. The von Neumann's inequality [16] says that for a contraction T on a Hilbert space and a polynomial p one has $\|p(T)\| \leq \sup\{|p(z)| : |z| \leq 1\}$.

Andô showed in [1] that the von Neumann's inequality holds for a pair of commuting contractions $\mathbf{T} = (T_1, T_2)$ and any polynomial p in two variables, i.e.

$$\|p(T_1, T_2)\| \leq \|p\|_\infty \quad (3.1.1)$$

where by $\|p\|_\infty$ one means the supremum norm over the bidisc $\mathbb{D}^2$ in $\mathbb{C}^2$.

The multicentric representation for $n = 2$, is worked out in Publication II and it is given by

$$\varphi(z_1, z_2) = \sum_{k=1}^{d_2} \sum_{j=1}^{d_1} \delta_{j,1}(z_1) \delta_{k,2}(z_2) f_{j,k}(w_1, w_2) \quad (3.1.2)$$

with $w_1 = p_1(z_1), w_2 = p_2(z_2)$.

Replacing the variables (z_1, z_2) with a pair of commuting operators on a Hilbert space H , (T_1, T_2) , we use the fact that the von Neumann inequality holds for a pair of commuting contractions and we are able to show the following,

$$\|\varphi(T_1, T_2)\| \leq \sum_{k=1}^{d_2} \sum_{j=1}^{d_1} \|\delta_{j,1}(T_1)\| \|\delta_{k,2}(T_2)\| C(\rho_1) C(\rho_2) \sup_{\mathbf{z} \in V_{\mathbf{p}}} |\varphi(z_1, z_2)|, \quad (3.1.3)$$

where $\mathbf{z} = (z_1, z_2)$ and ρ_1 and ρ_2 are the levels of p_1 and p_2 , respectively, such that they do not pass through any critical points and $V_{\mathbf{p}} = \{\mathbf{z} \in \mathbb{C}^2 : |p_1(z_1)| \leq \rho_1 \text{ and } |p_2(z_2)| \leq \rho_2\}$.

The constants $C(\rho_1), C(\rho_2)$ are given in Theorem 2.1.1.

3.2 On the n -tuples of commuting operators

When moving to n variables, i.e. for $\mathbf{z} \in \mathbb{C}^n$ one represents the holomorphic function φ using the new variables $\mathbf{w}$, and with the help of a vector $\mathbf{f}$ containing vector-valued functions of variables w_j , mapping $\mathbf{w} \mapsto \mathbf{f}(\mathbf{w}) \in \mathbb{C}^d$ (by $\mathbf{d}$ we mean $d_1 \times d_2 \times \cdots \times d_n$). The multicentric representation will then be of the form

$$\varphi(\mathbf{z}) = \sum_{\mathbf{j}} \delta_{\mathbf{j}}(\mathbf{z}) \mathbf{f}(\mathbf{w}) \text{ with } \mathbf{w} = \mathbf{p}(\mathbf{z}), \quad (3.2.1)$$

where $\mathbf{j}$ is a multi-index $(j_1, \dots, j_n)$, with $j_r = 1, \dots, d_r$ for all $r = 1, 2, \dots, n$.

To replace now the scalar variables with n -tuple of commuting operators, $\mathbf{T} = (T_1, \dots, T_n)$, we need to have the von Neumann inequality holding true for $n > 2$ commuting contractions. It is known that this is not the case, but from [6] we learn that for a small class of polynomials and some constant C_n the von Neumann inequality holds when $n > 2$.

This class is called the k -homogeneous polynomials defined as follows.

Definition 3.2.1. A k -homogeneous polynomial in n variables is a function $P : \mathbb{C}^n \rightarrow \mathbb{C}$ of the form

$$P(z_1, \dots, z_n) = \sum_{\alpha \in \Lambda(k, n)} a_{\alpha} z^{\alpha},$$

where $\Lambda(k, n) := \{\alpha \in \mathbb{N}_0^n : |\alpha| := \alpha_1 + \cdots + \alpha_n = k\}$, $z^{\alpha} = z_1^{\alpha_1} \cdots z_n^{\alpha_n}$ and $a_{\alpha} \in \mathbb{C}$.

The author in [6] studied the asymptotic behaviour (as n tends to infinity) of the smallest constant $C_{k, \infty}(n)$ such that

$$\|P(T_1, \dots, T_n)\|_{\mathcal{B}(H)} \leq C_{k, \infty}(n) \sup\{|P(z_1, \dots, z_n)| : |z_i| \leq 1\}, \quad (3.2.2)$$

for every k -homogeneous polynomial P in n variables and every n -tuple of commuting contractions $\mathbf{T}$. The estimate of $C_{k, \infty}(n)$ according to [6] is

$$n^{\frac{1}{2}[\frac{k-1}{2}]} \ll C_{k, \infty}(n) \ll n^{\frac{k-2}{2}},$$

where $[z]$ denotes the integer part of z .

Further work was done in [12] where the authors consider $C_{k, q}(n)$, for $1 \leq q < \infty$, the smallest constant such that

$$\|P(T_1, \dots, T_n)\|_{\mathcal{B}(H)} \leq C_{k, q}(n) \sup\{|P(z_1, \dots, z_n)| : \sum_{i=1}^n |z_i|^q \leq 1\},$$

for every k -homogeneous polynomial P in n variables and every n -tuple of commuting contractions $\mathbf{T}$ with $\sum_{i=1}^n \|T_i\|^q \leq 1$. The upper and lower estimates for the growth of $C_{k, q}(n)$ are the following

$$\begin{aligned} n^{\frac{k-1}{q'} - \frac{1}{2}[\frac{k}{2}]} &\ll C_{k, q}(n) \ll n^{\frac{k-2}{q'}} \text{ for } 1 \leq q \leq 2, \\ n^{\frac{k}{2} - \frac{1}{2}([\frac{k}{2}] + 1)} &\ll C_{k, q}(n) \ll n^{\frac{k-2}{2}} \text{ for } 2 \leq q < \infty, \end{aligned}$$

where q' denotes the conjugate of q , i.e., $\frac{1}{q} + \frac{1}{q'} = 1$. In all the estimates k is considered to be fixed.

Based on the combinatorial methods from [6], the authors in [9] change the construction of the Hilbert space and the operators given there to find the exact asymptotic growth of $C_{k,\infty}(n)$, answering a question posed by Dixon [6]. Their main result in [9] is the following theorem.

Theorem 3.2.2. *For $k \geq 3$, and $1 \leq q < \infty$, let $C_{k,q}(n)$ be the smallest constant such that*

$$\|Q(T_1, \dots, T_n)\|_{\mathcal{B}(H)} \leq C_{k,q}(n) \sup\{|Q(z_1, \dots, z_n)| : \|(z_i)_i\|_q \leq 1\},$$

for every k -homogeneous polynomial Q in n variables and every n -tuple of commuting contractions $\mathbf{T}$ with $\sum_{i=1}^n \|T_i\|^q \leq 1$. Then

$$(i) \ C_{k,\infty} \sim n^{\frac{k-2}{2}}$$

(ii) *for $2 \leq q < \infty$ we have*

$$\log^{-3/q}(n) n^{\frac{k-2}{2}} \ll C_{k,q}(n) \ll n^{\frac{k-2}{2}}.$$

In particular, $n^{\frac{k-2}{2}-\varepsilon} \ll C_{k,q}(n) \ll n^{\frac{k-2}{2}}$ for every $\varepsilon > 0$.

Considering Dixon's estimate (3.2.2) we extend in Publication II the multicentric calculus for an n -tuple of commuting contractions $\mathbf{T} = (T_1, \dots, T_n)$ by writing the components of the vector $\mathbf{f}(\mathbf{w})$ in the multicentric representation (3.2.1) as a sum of m k -homogeneous polynomials, and thus $\varphi(\mathbf{z})$ becomes a sum of m multicentric decompositions φ_k , consisting of k -homogeneous polynomials in variable $\mathbf{w}$,

$$\varphi(\mathbf{z}) = \sum_{k=0}^m \varphi_k(\mathbf{z}). \quad (3.2.3)$$

For these we can apply the functional calculus, since the von Neumann's inequality holds. Then the main estimate is obtained by summing up all estimates for φ_k . Details are presented in Publication II.

Theorem 3.2.3. *Let $\mathbf{z} \in \mathbb{C}^n$, $\mathbf{p}(\mathbf{z}) = (p_1, \dots, p_n)$ be a vector of polynomials such that for $j = 1, \dots, n$, p_j is a monic polynomial of degree d_j with distinct roots and let $\rho_j \geq 0$ satisfying $|p_j(z_j)| \leq \rho_j$ be such that the lemniscate contains no critical points of p_j . Let $k \geq 3$ and let $\mathbf{T} = (T_1, \dots, T_n)$ be an n -tuple of commuting contractions. Denote*

$$V_{\mathbf{p}} = \{\mathbf{z} \in \mathbb{C}^n : |p_j(z_j)| \leq \rho_j, \text{ for } j = 1, \dots, n\}.$$

Then for all φ defined by (3.2.3) there holds

$$\|\varphi(\mathbf{T})\| \leq K \sup_{\mathbf{z} \in V_{\mathbf{p}}} |\varphi(\mathbf{z})|,$$

with K being a constant that depends on the levels ρ_j 's of the polynomials p_j 's, on δ_j at the operator $\mathbf{T}$ and on the degree of φ but not on φ itself.

Remark 3.2.4. Note that the above estimates from Theorem 3.2.3 work for any q such that $1 \leq q < \infty$ due to Theorem 3.2.2.

To summarize, we were able to extend the holomorphic multicentric representation to a pair of commuting operators and we managed to find a bound for the representation, knowing that the von Neumann inequality holds for a pair of commuting contractions. Furthermore, considering a smaller class of polynomials, i.e. k -homogeneous polynomials and a suitable constant, we could extend the calculus even more, to n -tuple of commuting operators and write the multicentric representation as a sum of multicentric representations consisting of k -homogeneous polynomials.

Remark 3.2.5. We only considered the commuting case but the calculus can also be carried out for non-commuting operators. To this end, a natural functional calculus for a pair (A, B) of operators arises from associating the function $\varphi(A, B)$ with a scalar function $\varphi(z_1, z_2) = \sum_{i,j} \alpha_{ij} z_1^i z_2^j$, that is,

$$\varphi(A, B)(C) = \sum_{i,j} \alpha_{ij} A^i C B^j.$$

Here A would map in a Banach space X , B in Y while C is the “variable” operator mapping Y to X .

It is easy to do the similar extension to multicentric but this was not pushed in the thesis. However, in Publication IV, in particular, the Sylvester equation is of this nature. For a glimpse of the non-commuting case and the value it may bring for the multicentric representation in general, one can check [21].

4. A Banach algebra with a multicentric calculus

4.1 Construction of the algebra

In [20] Olavi Nevanlinna extended the single variable multicentric holomorphic calculus (2.1.2) to functions that need not be differentiable, by constructing a Banach algebra for $\mathbb{C}^d$ -valued continuous functions in such a way that the original functions φ appear as Gelfand transforms. To this end, a multiplication was defined, called 'polyproduct' and denoted by $\odot$, and an operator norm, $\|\cdot\|$, which is equivalent with the max norm.

In other words, after defining an operator norm, $\|\cdot\|$, the vector space $C(M)^d$ of continuous functions f from a compact $M \subset \mathbb{C}$ into $\mathbb{C}^d$, with the operator norm $\|f\|$ and the product $\odot$ becomes a Banach algebra denoted by $C_\Lambda(M)$. Then the function φ in (2.1.2) appears as the Gelfand transform in the new algebra $C_\Lambda(M)$.

To extend the multicentric non-holomorphic functional calculus, shortly described above, to pair of commuting operators, a multiplication and an operator norm are defined. We use the multicentric representation of a pair of commuting variables, (3.1.2), and then replace the variables by operators.

The extension in our study only considers the commuting case, but there can also be value in the non-commuting case and for this one can check [21]. The difficulties in moving to a pair of variables is not in lifting commuting operators, but finding a Banach algebra which can then be identified with the space of continuous two variable functions. We work this out by finding a suitable product and operator norm which make the vector space of continuous functions a Banach algebra. Then another difficulty is finding the suitable cross norm so that the Banach algebra will be identified with the tensor product of single variable Banach algebras. That is, being able to identify the Banach space of continuous two variable functions with the tensor product of continuous one variable Banach spaces. Finally, because of these identifications we are able to apply the Gelfand theory and to

find the characters of the Banach algebra. It then turns out that this represents, like in [20], the original functions as Gelfand transforms.

Let M_1, M_2 be two compact sets, hence $M_1 \times M_2$ is a compact set. Given continuous vector-valued functions f mapping a compact $M_1 \times M_2 \subset \mathbb{C}^2$ into $\mathbb{C}^{d_1 \times d_2}$ determines a unique function φ on $K_1 \times K_2 = p_1^{-1}(M_1) \times p_2^{-1}(M_2)$ given in (3.1.2) The Lagrange interpolation polynomials at λ_j and μ_k , respectively, $\delta_{1,j} \in \mathbb{P}^{d_1-1}$, $\delta_{2,k} \in \mathbb{P}^{d_2-1}$, are given by

$$\delta_{1,j}(z_1) = \frac{w_1}{p'_1(\lambda_j)(z_1 - \lambda_j)} \quad \text{and} \quad \delta_{2,k}(z_2) = \frac{w_2}{p'_2(\mu_k)(z_2 - \mu_k)}.$$

From the single variable case we know that the space of continuous functions from M_i to $\mathbb{C}^{d_i}$, $i = 1, 2$, that is, $C(M_1, \mathbb{C}^{d_1})$, and $C(M_2, \mathbb{C}^{d_2})$, together with the multiplication and operator norm are the algebras $C_{\Lambda_1}(M_1)$ and $C_{\Lambda_2}(M_2)$. Then the tensor product $C_{\Lambda_1}(M_1) \otimes C_{\Lambda_2}(M_2)$ will give the new algebra structure for the two variable case, which is denoted $C_{\Lambda_1 \times \Lambda_2}(M_1 \times M_2)$.

The multiplication, $\odot$, in the vector space $C(M_1 \times M_2, \mathbb{C}^{d_1 \times d_2})$ and the operator norm are below presented after few notations are in place. Set

$$L_1 \text{ a matrix with zero diagonal and } L_{1,jl} = \frac{1}{\lambda_j - \lambda_l}$$

$$L_2 \text{ a matrix with zero diagonal and } L_{2,km} = \frac{1}{\mu_k - \mu_m}$$

$$l_1 \text{ a vector in } \mathbb{C}^{d_1} \text{ with elements } l_{1,l} = \frac{1}{p'_1(\lambda_l)}$$

$$l_2 \text{ a vector in } \mathbb{C}^{d_2} \text{ with elements } l_{2,m} = \frac{1}{p'_2(\mu_m)}.$$

We use L_1^T, L_2^T, l_1^T and l_2^T for the transpose of L_1, L_2, l_1 and l_2 respectively. Then define for $f \in \mathbb{C}^{d_1 \times d_2}$

$$\square_k : f \mapsto \square_k f = \begin{pmatrix} 0 & f_{1,k} - f_{2,k} & \cdots & f_{1,k} - f_{d_1,k} \\ f_{2,k} - f_{1,k} & 0 & \cdots & f_{2,k} - f_{d_1,k} \\ \cdots & \cdots & \cdots & \cdots \\ f_{d_1,k} - f_{1,k} & \cdots & f_{d_1,k} - f_{d_1-1,k} & 0 \end{pmatrix}$$

called the boxing of the k^{th} column of $f(w_1, w_2)$, and

$${}_j\square : f \mapsto {}_j\square f = \begin{pmatrix} 0 & f_{j,1} - f_{j,2} & \cdots & f_{j,1} - f_{j,d_2} \\ f_{j,2} - f_{j,1} & 0 & \cdots & f_{j,2} - f_{j,d_2} \\ \cdots & \cdots & \cdots & \cdots \\ f_{j,d_2} - f_{j,1} & \cdots & f_{j,d_2} - f_{j,d_2-1} & 0 \end{pmatrix}$$

called the boxing of the transpose of the j^{th} row of $f(w_1, w_2)$. The boxing notion was defined in [20] and here we use the same notion since the formula agrees with the one in [20] if defined for one variable functions.

Definition 4.1.1. Let f and g be pointwisely defined functions from $M_1 \times M_2 \subset \mathbb{C}^2$ into $\mathbb{C}^{d_1 \times d_2}$. Then the 'polyproduct' $f \odot g$ is a function defined on $M_1 \times M_2$ taking values in $\mathbb{C}^{d_1 \times d_2}$ such that elementwise we have

$$\begin{aligned}
 (f \odot g)_{jk}(w_1, w_2) &= (f_{j,k} \circ g_{j,k})(w_1, w_2) \\
 &\quad - w_1(L_1 \circ \square_k f(w_1, w_2) \circ \square_k g(w_1, w_2))l_1 \\
 &\quad - w_2 l_2^T (L_2^T \circ {}_j \square f^T(w_1, w_2) \circ {}_j \square g^T(w_1, w_2)) \\
 &\quad + w_1 w_2 (l_2 \otimes l_1)^T ((L_2^T \otimes {}_j L_1^T) \circ \boxtimes_j f^T(w_1, w_2) \circ \boxtimes_j g^T(w_1, w_2))
 \end{aligned} \tag{4.1.1}$$

where by ${}_j L_1^T$ we mean the transpose of the j -th line in L_1 , i.e. a vector in $\mathbb{C}^{d_1}$, and for $f \in \mathbb{C}^{d_1 \times d_2}$ we define $\boxtimes_j f^T$ as the matrix in $\mathbb{C}^{d_1^T \times d_2}$ with d_2 columns, denoted col_k , for $k = 1, \dots, d_2$.

Note that this definition does not present all formulas. They are long and can be found in Publication III.

Let us, for short, denote $X = C(M_1 \times M_2, \mathbb{C}^{d_1 \times d_2})$ and $f \in X$ having the norm

$$|f|_\infty = \max_{j,k} \max_{M_1 \times M_2} |f_{jk}(w_1, w_2)|,$$

for $j = 1, 2, \dots, d_1$, $k = 1, 2, \dots, d_2$. Thus X is a Banach space. Consider the multiplication in the X being the one defined above. We proceed by looking at how the product $\odot$ is compatible with the norm in this Banach space X . Thus is defined an operator norm that is equivalent with the norm on X .

Definition 4.1.2. For $f \in X$ we set

$$\|f\| := \sup_{|g|_\infty \leq 1} |f \odot g|_\infty.$$

This is a norm in X and it is equivalent with $|\cdot|_\infty$.

Proposition 4.1.3. There is a C , depending only on $M_1 \times M_2$ and Λ_i , for $i = 1, 2$ such that

$$\|f \odot g\| \leq \|f\| \|g\|, \tag{4.1.2}$$

$$|f|_\infty \leq \|f\| \leq C|f|_\infty. \tag{4.1.3}$$

We have now all ingredients for defining the Banach algebra.

Definition 4.1.4. The Banach space $C(M_1 \times M_2, \mathbb{C}^{d_1 \times d_2})$ of continuous functions from compact $M_1 \times M_2$ into $\mathbb{C}^{d_1 \times d_2}$ with the operator norm and product $\odot$ becomes a Banach algebra and is denoted by $C_{\Lambda_1 \times \Lambda_2}(M_1 \times M_2)$.

Moreover, $C_{\Lambda_1 \times \Lambda_2}(M_1 \times M_2)$ is a unital Banach algebra.

In the algebra setup the next step was to find a cross norm, say α , on the completed tensor algebras $C_{\Lambda_1}(M_1) \hat{\otimes}_\alpha C_{\Lambda_2}(M_2)$ which will help in canonically identify this completed tensor algebra with the algebra $C_{\Lambda_1 \times \Lambda_2}(M_1 \times M_2)$.

Now recall from [7] the following.

Definition 4.1.5. Let X and Y be Banach spaces and $X \otimes_a Y$ be the algebraic tensor product of X and Y as linear vector spaces over $\mathbb{C}$. If for $u \in X \otimes_a Y$ one defines

$$\|u\|_\gamma = \inf \left\{ \sum_{i=1}^n \|x_i\| \|y_i\| : x_1, \dots, x_n \in X, y_1, \dots, y_n \in Y, u = \sum_{i=1}^n x_i \otimes y_i \right\},$$

then $\|\cdot\|_\gamma$ is a norm on $X \otimes_a Y$. The completion of $X \otimes_a Y$ is the projective tensor product of X and Y and is denoted $X \widehat{\otimes}_\gamma Y$.

Definition 4.1.6. Let X and Y be Banach spaces and $X \otimes_a Y$ be the algebraic tensor product of X and Y as linear vector spaces over $\mathbb{C}$. Let X^* and Y^* be the dual of X and Y , respectively. If one defines for $u \in X \otimes_a Y$ the norm given by

$$\|u\|_\varepsilon = \sup \left\{ \left| \sum_{i=1}^n \varphi(x_i) \psi(y_i) \right| : x_1, \dots, x_n \in X, y_1, \dots, y_n \in Y, \right. \\ \left. \varphi \in (X^*)_1, \psi \in (Y^*)_1, u = \sum_{i=1}^n x_i \otimes y_i \right\},$$

where $(X^*)_1$ and $(Y^*)_1$ denote the unit balls of the dual Banach spaces X^* and Y^* , respectively, then $\|\cdot\|_\varepsilon$ is a norm on $X \otimes_a Y$. The completion of $X \otimes_a Y$ is the injective tensor product of X and Y and is denoted $X \widehat{\otimes}_\varepsilon Y$.

Then for X and Y compact Hausdorff spaces

$$C(X) \widehat{\otimes}_\varepsilon C(Y) = C(X \times Y); \quad (4.1.4)$$

and

$$C(X) \widehat{\otimes}_\gamma C(Y) = C(X \times Y) \text{ iff } X \text{ or } Y \text{ is finite,} \quad (4.1.5)$$

where $C(X)$ means the space of continuous functions on X . The statement given by (4.1.4) appears in [24] as well.

Furthermore, according to [7], if A_1 and A_2 are Banach algebras, then $A_1 \widehat{\otimes}_\varepsilon A_2$ and $A_1 \widehat{\otimes}_\gamma A_2$ are Banach algebras. And, if A_1 and A_2 are commutative with maximal ideal spaces $\mathfrak{M}_1$ and $\mathfrak{M}_2$, respectively, then $\mathfrak{M}_1 \times \mathfrak{M}_2$ is the maximal ideal space of both $A_1 \widehat{\otimes}_\varepsilon A_2$ and $A_1 \widehat{\otimes}_\gamma A_2$.

From [24] we learn that for a Banach space Y and a compact set $M \subset \mathbb{C}$ the injective tensor product $C(K) \widehat{\otimes}_\varepsilon Y$ can be identified with the Banach space $C(M, Y)$ and the norm on this space is given by $\|f\|_\infty = \sup \{ \|f(t)\| : t \in M \}$. Moreover, this identification can be applied to get a representation of a space of continuous functions of two variables as an injective tensor product of two spaces of continuous functions, that is, for M_1, M_2 compact spaces,

$$C(M_1) \widehat{\otimes}_\varepsilon C(M_2) = C(M_1 \times M_2),$$

since $C(M_1 \times M_2)$ can be identified with $C(M_1, C(M_2))$.

It was then easily shown in Publication III that the following holds.

Theorem 4.1.7. *The Banach space $C(M_1 \times M_2, \mathbb{C}^{d_1 \times d_2})$ can be canonically identified with the injective tensor product $C(M_1, \mathbb{C}^{d_1}) \widehat{\otimes}_\varepsilon C(M_2, \mathbb{C}^{d_2})$, in short*

$$C(M_1 \times M_2, \mathbb{C}^{d_1 \times d_2}) = C(M_1, \mathbb{C}^{d_1}) \widehat{\otimes}_\varepsilon C(M_2, \mathbb{C}^{d_2}).$$

4.2 Application to the Gelfand theory

Next recall from [13] that given A_1 and A_2 two function algebras on compact Hausdorff spaces M_1 and M_2 , respectively, and setting $A_1 \widehat{\otimes}_\varepsilon A_2$ to be the completion of $A_1 \otimes_a A_2$ (the algebraic tensor product of A_1 and A_2) under the ε -norm, then the ε -norm is identical with the uniform norm on $M_1 \times M_2$ and $C(M_1) \widehat{\otimes}_\varepsilon C(M_2) = C(M_1 \times M_2)$. Thus $A_1 \widehat{\otimes}_\varepsilon A_2$ is a Banach algebra. It is easily seen that $A_1 \widehat{\otimes}_\varepsilon A_2$ becomes a function algebra on $M_1 \times M_2$, denoted by $\mathcal{U}$.

Let $\mathfrak{M}(A)$ denote the maximal ideal space of an algebra A . It is known that $\mathfrak{M}(\mathcal{U})$ is homeomorphic to $\mathfrak{M}(A_1) \times \mathfrak{M}(A_2)$, see Theorem 2 in [26].

Remark 4.2.1. For a commutative Banach algebra with unit, all characters - complex homomorphisms - are automatically bounded and of norm 1. The focus is then on maximal ideals since these are kernels of characters.

If we denote by A_1 the commutative Banach algebra with unit $C_{\Lambda_1}(M_1)$ and by A_2 the commutative Banach algebra with unit $C_{\Lambda_2}(M_2)$, we have that $X = A_1 \widehat{\otimes}_\varepsilon A_2$, where $X = C_{\Lambda_1 \times \Lambda_2}(M_1 \times M_2)$, is a Banach algebra with unit as defined above. Hence it is easily seen that we have the identification

$$\mathfrak{M}(A_1) \times \mathfrak{M}(A_2) = \mathfrak{M}(X).$$

In other words, one gets all the characters of the Banach algebra X out of the characters of the Banach algebras A_1 and A_2 . This is just Theorem 2 in [26].

Therefore, in the multicentric setup, the characters of X are given by

$$\chi_{(z_1, z_2)} : f \mapsto \sum_{j=1}^{d_1} \sum_{k=1}^{d_2} \delta_{j,1}(z_1) \delta_{k,2}(z_2) f_{j,k}(p_1(z_1), p_2(z_2)) \quad (4.2.1)$$

and the space of all characters of X will be given by the set

$$\mathcal{X}_X = \{\chi_{(z_1, z_2)} : (z_1, z_2) \in p_1^{-1}(M_1) \times p_2^{-1}(M_2)\}.$$

Further work in Publication III shows that the multicentric representation operator acts as the Gelfand transformation.

Definition 4.2.2. *Given $f \in C_{\Lambda_1 \times \Lambda_2}(M_1 \times M_2)$ we set*

$$\hat{f} : p_1^{-1}(M_1) \times p_2^{-1}(M_2) \rightarrow \mathbb{C}^2$$

$$\hat{f} : (z_1, z_2) \mapsto \hat{f}((z_1, z_2)) = \sum_{j=1}^{d_1} \sum_{k=1}^{d_2} \delta_{j,1}(z_1) \delta_{k,2}(z_2) f_{j,k}(p_1(z_1), p_2(z_2)).$$

Let $K_1 \times K_2 = p_1^{-1}(M_1) \times p_2^{-1}(M_2)$. For $f \in C_{\Lambda_1 \times \Lambda_2}(M_1 \times M_2)$, since the algebra $C_{\Lambda_1 \times \Lambda_2}(M_1 \times M_2)$ is commutative, then the spectrum $\sigma(f)$ can be expressed in terms of the Gelfand transform

$$\sigma(f) = \{\hat{f}((z_1, z_2)) : (z_1, z_2) \in K_1 \times K_2\}.$$

Lastly, the semi-simplicity property of the algebra X is worked out. Recall that an algebra $\mathcal{A}$ is *semi-simple* if $\text{rad}\mathcal{A} = \{0\}$.

Theorem 4 in [26] is saying that for given A_1 and A_2 commutative Banach algebras the following holds:

1. If either A_1 or A_2 satisfy the condition of approximation then $A_1 \hat{\otimes}_{\gamma} A_2$ is semi-simple iff A_1 and A_2 are semi-simple.
2. Suppose that $A_1 \hat{\otimes}_{\varepsilon} A_2$ becomes a Banach algebra, then $A_1 \hat{\otimes}_{\varepsilon} A_2$ is semi-simple if and only if A_1 and A_2 are semi-simple.

Therefore, since

$$X = C_{\Lambda_1 \times \Lambda_2}(M_1 \times M_2)$$

is a Banach algebra, we get that X is semi-simple iff $A_1 = C_{\Lambda_1}(M_1)$ and $A_2 = C_{\Lambda_2}(M_2)$ are semi-simple. In [20] it is shown that $C_{\Lambda}(M)$ semi-simple iff M contains no isolated critical values of p .

To summarize, we have defined a product and an operator norm so that the constructed tensor product Banach algebra can be associated with the tensor product of one variable algebras. Then, with a suitable cross norm, the space of continuous functions $C(M_1 \times M_2)$ is identified with the completed tensor product of one variable continuous functions, $C(M_1), C(M_2)$. Hence we have identified the tensor Banach algebra with the space of continuous functions, we extended the multicentric calculus to a class of functions φ for which the constructed Banach algebra structure give the Gelfand transform, we have showed that all the characters in that algebra are determined by the characters in the single variable algebras and we formulated the condition for the algebra to be semi-simple.

5. Change of variables using rational functions

The single variable multicentric calculus is further developed in Publication IV where the authors studied the necessary modifications needed if rational function $r = p/q$ are taken as new variables. This allows one to have a representation for functions in sets that need not be polynomially convex, by considering $r = p/q$ with q of lower degree than p . In such case, much of the multicentric calculus carries over with minor changes.

5.1 Representation using rational functions as variables

We start by formulating a result equivalent to Hilbert lemniscate lemma, approximating any compact set arbitrarily well in a neighbourhood of it. Given a compact K such that $\mathbb{C} \setminus K$ is *simply connected* and $\varepsilon > 0$ there exists a polynomial p such that if

$$V_p = \{z \in \mathbb{C} : |p(z)| \leq 1\}$$

and $K_\varepsilon = \{z : \text{dist}(z, K) \leq \varepsilon\}$, then

$$K \Subset V_p \Subset K_\varepsilon.$$

Here $\Subset$ means that the smaller compact is included in the interior of the larger compact.

Suppose $r = p/q$ is a rational function, with p and q having no common roots. Set

$$V_r = \{z \in \mathbb{C} : |r(z)| \leq 1\}. \quad (5.1.1)$$

As we do not have control of the size of r globally we restrict V_r into a compact set. Denote by Γ_ε the following compact set surrounding K :

$$\Gamma_\varepsilon := K_\varepsilon \setminus \text{int } K_{\varepsilon/2}. \quad (5.1.2)$$

Theorem 5.1.1. *Given a compact $K \subset \mathbb{C}$ and $\varepsilon > 0$ let Γ_ε be as in (5.1.2). Then there exists a rational function r such that*

$$K \Subset V_r \text{ and } V_r \cap \Gamma_\varepsilon = \emptyset. \quad (5.1.3)$$

Further, the rational function $r = p/q$ can be so chosen that $\deg q < \deg p$.

Then, another result is formulated as a corollary, where the spectrum of operators play the role of the compact set. Given a bounded operator A in a Banach space $\mathcal{X}$, $A \in B(\mathcal{X})$, fix $\varepsilon > 0$ and let $r = p/q$ be as in the above Theorem when the spectrum $\sigma(A)$ is taken as the compact set K . In particular, r is holomorphic in the spectrum and $r(A)$ is a well defined bounded operator. Then

$$\|r(A)^m\|^{1/m} \rightarrow \rho(r(A)) = \sup_{z \in \sigma(A)} |r(z)|.$$

Since r is not a constant, and $\sigma(A) \in V_r$, we have by maximum principle

$$\sup_{z \in \sigma(A)} |r(z)| < 1.$$

But then there exists n such that $\|r(A)^n\| < 1$. Denote by $\tilde{p}$ a tiny perturbation of p^n so that all roots of $\tilde{r} = \tilde{p}/q^n$ are simple and we still have $\|\tilde{r}(A)\| < 1$. In order to formulate the corollary, let us denote by Γ_ε the set surrounding the spectrum as in (5.1.2) with $\sigma(A) = K$.

Corollary 5.1.2. *Given a bounded operator $A \in B(\mathcal{X})$, fix an $\varepsilon > 0$ and denote by Γ_ε the set around the spectrum $\sigma(A)$ as above. Then there exists a rational function $r = p/q$, such that $\deg q < \deg p$ where p has simple roots and $\|r(A)\| < 1$, while $|r(z)| > 1$ for $z \in \Gamma_\varepsilon$.*

Let $r = p/q$ with p having d simple roots $\Lambda = \{\lambda_j\}$ so that $q(\lambda_j) \neq 0$ and such that $d = \deg p > \deg q$. Denoting by δ_j the rational functions

$$\delta_j(z) = \frac{r(z)}{r'(\lambda_j)(z - \lambda_j)} \quad (5.1.4)$$

we consider representations of scalar functions φ in the form

$$\varphi(z) = \sum_{j=1}^d \delta_j(z) f_j(w) \quad \text{where } w = r(z). \quad (5.1.5)$$

The assumptions that the roots of p are simple and that q is of lower degree than p are not necessary, but made for simplifying the discussion.

Modifying the discussion in section 2.1 in [19] we easily show the existence and uniqueness of the new representation, (5.1.5).

Assume now that $M \subset \mathbb{C}$ is compact and let $K = r^{-1}(M)$. Denote by $M_0 = M \setminus W_c$ where W_c denotes the set of critical values of r and put $K_0 = r^{-1}(M_0)$.

Proposition 5.1.3. *Given a function φ mapping $K_0 \rightarrow \mathbb{C}$, with K_0 as above. Then there exists a unique f mapping $M_0 \rightarrow \mathbb{C}^d$ such that*

$$\varphi(z) = \sum_{j=1}^d \delta_j(z) f_j(r(z)) \quad (5.1.6)$$

holds for $z \in K_0$. The function f inherits the smoothness of φ . In particular, if φ is continuous or holomorphic in K_0 , then f is continuous or holomorphic in M_0 .

At a critical value w_c there are fewer equations and $f_k(w_c)$ s do exist, but they are not unique. It is therefore of interest to study what continuity conditions on φ guarantee continuity of f_k s at critical values. Using the Cauchy integral we see that if φ is holomorphic in K , then f can be extended from M_0 to M so that it is holomorphic also at the critical values.

Lemma 5.1.4. *Let $\deg q < \deg p$ and denote by Z_q the zeros of q . Assume $z \notin Z_q$. Then*

$$\sum_{j=1}^d \delta_j(z) = 1. \quad (5.1.7)$$

In the non-holomorphic case things are worked out following [20]. Let $\{e_i\}$ denote the standard basis of $\mathbb{C}^d$. Assume we are given a $d \times d$ multiplication table $\Sigma = \{\sigma_{ij}\}$ and a frozen $w \in M$.

Definition 5.1.5. *Let p be monic of degree d with simple roots $\{\lambda_j\}$ and q of degree at most $d-1$, with $q(\lambda_j) \neq 0$ and denote $r = p/q$. If the multiplication table satisfies*

$$\sigma_{ij} = \frac{1}{r'(\lambda_j)} \frac{1}{\lambda_i - \lambda_j}, \quad (5.1.8)$$

then we say that the product $\odot$ in $C_\Sigma(M)$ is determined by the rational function r .

Theorem 5.1.6. *In the unital commutative Banach algebra $C_\Sigma(M)$ with Σ generated by a rational function as in Definition 5.1.5, all characters are of the form*

$$\chi_z : f \mapsto \sum_{i=1}^d \delta_i(z) f_i(w). \quad (5.1.9)$$

In particular the Gelfand transformation $f \mapsto \hat{f}$ is given by $\hat{f}(z) = \chi_z(f)$.

5.2 Applications and remarks

Publication IV presents two applications, one to the Sylvester equation and one to K-spectral sets.

Let A and B be bounded operators in Banach spaces $\mathcal{X}, \mathcal{Y}$ respectively. Then $AX - XB = C$ is the related Sylvester equation, where C is a given operator and X the unknown, both mapping $\mathcal{Y}$ to $\mathcal{X}$. It is well known [2] that a unique bounded X exists for every bounded C if and only if A and B have disjoint spectra: $\sigma(A) \cap \sigma(B) = \emptyset$. In Section 5 of [21], the multicentric calculus was applied to the case where the polynomial convex hulls of the

spectra were disjoint: $\widehat{\sigma(A)} \cap \widehat{\sigma(B)} = \emptyset$. The solution was constructed as a convergent power series provided that one has a polynomial lemniscate which separates the spectra into different components. When using rational lemniscates we can remove the need to assume that the polynomially convex hulls do not intersect.

It was shown in [18] that polynomial lemniscate sets are K -spectral sets, provided that the boundaries are smooth, meaning that they do not contain critical points. As a second application, it is shown that this extends to rational lemniscates as well. Details can be checked in Publication IV.

Remark 5.2.1. We did not consider a general rational function, but if we think of the case where the degree of q is greater than the degree of p , then we can not have (5.1.7) holding true. But it does not mean that this would not be useful. Doing that, then one could represent $1/\text{variable}$. In the complement of compact sets such can be useful.

On the other hand, if p and q are of the same degree, it can be done but for a special reason. It would allow one to look at half plane. Simple Möbius transformation can map inside a disc. Taking a polynomial, then a Möbius transformation and have a rational with same degrees is possible, but the need must be already known.

OR (last paragraph): On the other hand, we can consider p and q having the same degree and work out the calculus for such a case, but the reason must be known. It would allow one to look at half plane. For this case, we could take a polynomial and then a Möbius transformation, since simple Möbius transformation can map inside a disc.

References

- [1] Andô T. On a pair of commuting contractions: Acta Sci. Math. (Szeged); 1963; Vol. 24; p. 88-90.
Andrei D. Multicentric holomorphic calculus for n -tuples of commuting operators: Adv. Oper. Theory; 2019; Vol. 4 (2): p. 447–461.
- [2] Bhatia R, Rosenthal P. How and Why to Solve the Operator Equation $AX - XB = Y$: Bull. London Math. Soc.; 1997; Vol. 29; p. 1-21.
- [3] Cauchy AL. Résumé d'un mémoire sur la Mécanique céleste et sur un nouveau calcul appelé calcul des limites: Ouvres; 1825; Vol. 2; no.12; p. 48–112.
- [4] Conway JB. A Course in Functional Analysis: Springer; 1990; New York.
- [5] Curtiss JH. On the Jacobi Series: Trans. Amer. Math. Soc.; 1941; Vol. 39; no. 3.
- [6] Dixon PG. The von Neumann inequality for polynomials of degree greater than two: J. London Math. Soc. (2); 1976; Vol. 14(2); p. 369-375.
- [7] Douglas RG. Banach Algebra Techniques in Operator Theory: Academic Press; 1972; New York and London.
- [8] Fekete M. Über die Verallgemeinerung der Picard-Landauschen und Picard-Schottkyschen Sätze auf Reihen, die nach Potenzen eines Polynoms fortschreiten und Polynome niedrigeren Grades zu Koeffizienten haben: Math. Ann. 106; 1932.
- [9] Galicer D. Muro S. Sevilla-Peris P. Asymptotic estimates on the von Neumann inequality for homogeneous polynomials: J. Reine Angew. Math.; 2016.
- [10] Jacobi CGJ. Über Reihenentwicklungen, welche nach den Potenzen eines gegebenen Polynoms fortschreiten, und zu Coeffizienten Polynome eines niederen Grades haben: J. Reine Angew. Math. 53; 1856.

- [11] Kienast A. Über die Darstellung der analytischen Funktionen durch Reihen, die nach Potenzen eines Polynomes fortschreiten und Polynomes eines niedrigeren Grades zu Koeffizienten haben: Dissertation; Zürich; 1906.
- [12] Mantero AM. Tonge A. Banach algebras and von Neumann's inequality: Proc. London Math. Soc. (3); 1979; Vol. 38(2); p. 309-334.
- [13] Mochizuki N. The tensor product of function algebras: Tohoku Math. J.; 1965; (2) 17 no.2; p. 139–146.
- [14] Müller V. Spectral theory of linear operators and spectral systems in Banach algebras: Operator Theory: Advances and Applications; Birkhäuser Basel; 2003.
- [15] Müller V. Taylor functional calculus: Springer Basel; 2015; p. 1181-1215.
- [16] von Neumann J. Eine Spektraltheorie für allgemeine Operatoren eines unitären Raumes: Math. Nachr. 4; 1951; p. 258-281.
- [17] Nevanlinna O. Computing the spectrum and representing the resolvent: Numer. Funct. Anal. Optim. 30; 2009; p. 1025-1047.
- [18] Nevanlinna O. Lemniscates and K-spectral sets: J. Funct. Anal.; 2012; 262: p. 1728–1741.
- [19] Nevanlinna O. Multicentric holomorphic calculus: Comput. Methods Funct. Theory 2012; 12 (1): p. 45–65.
- [20] Nevanlinna O. Polynomial as a new variable ? A Banach algebra with a functional calculus: Oper. Matrices; 2016; vol. 10; no. 3: p. 567–592.
- [21] Nevanlinna O. Sylvester equations and polynomial separation by spectrum: Oper. Matrices; 2019; vol. 13; no. 3; p.867–885.
- [22] O'Meara KC, Vinsonhaler C. On approximately simultaneously diagonalizable matrices: Linear Algebra Appl.; 2006; 412: p. 39–74.
- [23] Paulsen V. Completely Bounded Maps and Operator Algebras: Cambridge University Press; 2002;
- [24] Ryan RA. Introduction to Tensor Products of Banach Spaces: Springer; 2002; London.
- [25] Stewart J. Calculus: Cengage Learning; 2015; USA.
- [26] Tomiyama J. Tensor products of commutative Banach algebras: Tohoku Math. J.; 1960; (2) 12 no. 1: p. 147–154.
- [27] Turumaru T. On the direct-product of operator algebras: IV. Tohoku Math. J.; 1956; (2) 8 no. 3: p. 281–285.

ISBN 978-952-64-0624-4 (printed)

ISBN 978-952-64-0625-1 (pdf)

ISSN 1799-4934 (printed)

ISSN 1799-4942 (pdf)

Aalto University

School of Science

Department of Mathematics and Systems Analysis

www.aalto.fi

**BUSINESS +
ECONOMY**

**ART +
DESIGN +
ARCHITECTURE**

**SCIENCE +
TECHNOLOGY**

CROSSOVER

**DOCTORAL
DISSERTATIONS**