Work and energy fluctuations in open quantum systems

Rui Sampaio

Work and energy fluctuations in open quantum systems

Rui Sampaio

A doctoral dissertation completed for the degree of Doctor of Science (Technology) to be defended, with the permission of the Aalto University School of Science, at a public examination held at the lecture hall M1 (Otakaari 1, Espoo) of the school on 15 February 2019 at noon.

Aalto University
School of Science
Department of Applied Physics
Multiscale Statistical and Quantum Physics group

Supervising professor

Prof. Tapio Ala-Nissilä

Thesis advisor

Prof. Tapio Ala-Nissilä

Preliminary examiners

Prof. Dr. Joachim Ankerhold, University of Ulm, Germany

Prof. Juan Garrahan, University of Nottingham, United Kingdom

Opponent

Dr. Guillermo Albareda, Max Planck Institute for the Structure and Dynamics of Matter, Germany

Aalto University publication series

DOCTORAL DISSERTATIONS 25/2019

© 2019 Rui Sampaio

ISBN 978-952-60-8414-5 (printed)

ISBN 978-952-60-8415-2 (pdf)

ISSN 1799-4934 (printed)

ISSN 1799-4942 (pdf)

<http://urn.fi/URN:ISBN:978-952-60-8415-2>

Unigrafia Oy

Helsinki 2019

Finland

Author

Rui Sampaio

Name of the doctoral dissertation

Work and energy fluctuations in open quantum systems

Publisher School of Science

Unit Department of Applied Physics

Series Aalto University publication series DOCTORAL DISSERTATIONS 25/2019

Field of research Engineering Physics

Manuscript submitted 4 December 2018

Date of the defence 15 February 2019

Permission to publish granted (date) 23 January 2019

Language English

☐ **Monograph**

☒ **Article dissertation**

☐ **Essay dissertation**

Abstract

Technological advances over the past few decades have made it possible to address quantum systems down to individual quanta of energy. Artificial atoms are now routinely fabricated and manipulated using solid-state technology. Exciting new applications are now within grasp, such as quantum heat engines, which have been theorised to achieve energy efficiencies not possible for their classical counterparts. Energy fluctuations are an inherent part of these devices and a detailed understanding of them is required. However, the proper theoretical framework under which such fluctuations are to be understood is still debated.

In this thesis, we study energy fluctuations in open quantum systems at the level of individual runs of an experimental protocol. This work has been carried out in two parts.

In the first part, we focus on calorimetric measurements – a promising new experimental protocol to continuously measure energy fluctuations of a qubit interacting with a thermal environment. Applying the protocol to more complex systems is not straightforward. Using a simple two-level approximation, we show that for some experimental parameters this protocol can be extended to a driven harmonic oscillator. Another aspect of the calorimetric protocol is that it can induce non-Markovian effects which affect how work and heat are defined. We quantify the degree of non-Markovianity induced by this measurement protocol, when applied to a qubit, based on a previously developed theoretical model. Importantly, we show that non-Markovianity does not necessarily decrease monotonically with the environment size.

The attempt to extend the calorimetric protocol to general systems has led to the second part of this thesis. The calorimetric detection is a particular case of a continuously monitored system. However, other schemes exist that lead to different definitions of work and heat. Reconciling these approaches is not straightforward, as evidenced by the many work definitions present in the literature. The difficulty is centred at the fact that, in the orthodox interpretation of quantum mechanics, energy is not always a well-defined quantity.

Here, based on the Bohmian interpretation of quantum mechanics, we put forward a general framework to remove this problem in a clear conceptual manner. In particular, we show that the classical definition of work for closed systems can be straightforwardly generalised to the quantum case. For an open system, we used the Bohmian conditional wave function which provides a stochastic state evolution for the open system. We show that this approach can lead to new insights regarding the role of entanglement in energy fluctuations and has the potential to provide a general and observer-independent framework of energy fluctuations in open quantum systems.

The results presented in this thesis give a small contribution to the developing field of quantum thermodynamics.

Keywords open quantum systems, single-shot energy fluctuations, calorimetric measurements, bohmian mechanics, conditional wave function

ISBN (printed) 978-952-60-8414-5

ISBN (pdf) 978-952-60-8415-2

ISSN (printed) 1799-4934

ISSN (pdf) 1799-4942

Location of publisher Helsinki

Location of printing Helsinki **Year** 2019

Pages 113

urn <http://urn.fi/URN:ISBN:978-952-60-8415-2>

Preface

The work presented in this dissertation was carried out during the years 2015-19 in the Mutliscale Statistical and Quantum Physics (MSP) group at Aalto University with support from the Magnus Ehrnrooth Foundation, the Education Network in Condensed Matter and Materials Physics (CMMP), the Centre of Excellence in Computational Nanoscience (COMP) and the Centre of Excellence in Quantum Technology, Finland (QTF).

First and foremost, I'd like to acknowledge my supervisor, Prof. Tapio Ala-Nissilä. Tapio was one of the first people I met when I first came to Finland, and without whom this dissertation had not been possible. I've learned much from Tapio and I've always appreciated the freedom given to me to explore different viewpoints and approaches, some less popular than others, to the problems at hand.

Secondly, I'd like to thank Dr. Thomas Philbin and Dr. Janet Anders, from Exeter University, for their contribution to the second half of this dissertation. I haven't immensely enjoyed the discussions with Tom on fundamental problems in modern physics, from interpretations to quantum mechanics to the arrow of time, and I am deeply thankful to Janet for her patience in molding my incoherent thoughts and ideas into proper scientific contributions.

It is my pleasure to also acknowledge Dr. Samu Suomela and Dr. Rebecca Schmidt. I have enjoyed our many discussions on the white board and their company throughout conferences, schools and workshops. They have played a fantastic role as opponents in my ramblings about quantum mechanics and its interpretations.

These four years and the work presented here were also shaped by good moments outside the office. I'd like to thank Miguel for his troubles organizing poker games, Vili for entertaining us with some blues on the piano, Cristian for some good chess games, Brecht for his hopeless Many-Worlds adoration, Milica for her sushi lessons and Asta, Jussi, Tapio and Sakari for all the Rock n' Roll. I'd also like to thank all those in the MSP group and the fourth floor for all the small moments that made coming to work always a joy.

On the family front, I'd like to thank my mom and dad for always supporting and encouraging me on this little adventure. This dissertation marks the end of a

chapter for me and I'm grateful to have had you both with me along the way. I'd also like to thank Jorma and Leila, for welcoming me in their home and making my trips to Oulu and around Finland always a pleasure.

Last, but no least, thank you Kristiina, for keeping me sane, giving me perspective and listening to my ramblings. One can easily get lost in the storm, but you have always been my light.

Espoo, January 25, 2019,

Rui Sampaio

Contents

Preface	1
Contents	3
List of Publications	5
Author's Contribution	7
1. Introduction	9
1.1 Motivation	9
1.2 Overview	10
2. Open quantum systems	13
2.1 The reduced density operator of the system	13
2.2 The Markovian master equation	14
2.3 Stochastic Schrödinger equations	14
2.4 Continuously monitored Markovian environment	15
2.5 Non-Markovianity measures	17
3. Calorimetric measurements of a driven harmonic oscillator	19
3.1 Qubit	19
3.2 Harmonic Oscillator	20
3.2.1 The calorimeter two level approximation	20
3.2.2 Results	21
4. Non-Markovianity due to driving and finite-size environment	27
4.1 The qubit-calorimeter model	27
4.2 BLP measure and initial conditions	28
4.3 Trace distance	29
4.4 Influence of the system parameters on $\mathcal{N}(\Phi)$	30
4.5 Conclusions	32
5. Energy fluctuations in closed quantum systems	35

5.1	Motivation	35
5.2	Two-point measurement protocol	36
5.3	Other approaches	36
5.4	A “no-go” theorem from quantum work	37
6.	Quantum Work in the Bohmian Framework	39
6.1	Bohmian Mechanics	39
6.1.1	Description of experiments	39
6.1.2	Dynamics	40
6.2	Connection to the Aharonov-Albert-Vaidman weak values . . .	41
6.3	Work in closed systems	42
6.4	Example: Driven harmonic oscillator	43
6.4.1	Thermal state as a mixture of energy eigenstates . .	44
6.4.2	Thermal state as a mixture of coherent states	45
6.4.3	Comparison of work averages	45
6.5	Conclusions	46
7.	Energy exchange and the Conditional Wave Function of open quantum systems	47
7.1	Motivation	47
7.2	Conditional Wave Functions	48
7.3	Conditional energy	49
7.4	Driving with entanglement	51
7.5	Conclusions	53
8.	Summary and Conclusions	55
	References	59
	Publications	65

List of Publications

This thesis consists of an overview and of the following publications which are referred to in the text by their Roman numerals.

- I** Rui Sampaio, Samu Suomela, and Tapio Ala-Nissila. Calorimetric measurement of work for a driven harmonic oscillator. *Physical Review E*, Volume 94, Issue 6, 062122 (9 pages), December 2016.
- II** Rui Sampaio, Samu Suomela, Rebecca Schmidt, and Tapio Ala-Nissila. Quantifying non-Markovianity due to driving and a finite-size environment in an open quantum system. *Physical Review A*, Volume 95, Issue 2, 022120 (7 pages), February 2017.
- III** R. Sampaio, S. Suomela, T. Ala-Nissila, J. Anders, and T. G. Philbin. Quantum work in the Bohmian framework. *Physical Review A*, Volume 97, Issue 1, 012131 (8 pages), January 2018.
- IV** R. Sampaio, S. Suomela, T. Ala-Nissila, J. Anders, and T. G. Philbin. Contributions to single-shot energy exchanges in open quantum systems. Submitted to *Physical Review E*, 11 pages, September 2018.

Author's Contribution

Publication I: “Calorimetric measurement of work for a driven harmonic oscillator”

The author derived the analytical results and carried out the numerical simulations. The author wrote the first version of the manuscript.

Publication II: “Quantifying non-Markovianity due to driving and a finite-size environment in an open quantum system”

The author derived part of the analytical results and carried out the numerical simulations. The author wrote the first version of the manuscript.

Publication III: “Quantum work in the Bohmian framework”

The author derived part of the analytical results. The author wrote the first version of the manuscript.

Publication IV: “Contributions to single-shot energy exchanges in open quantum systems”

The author derived the analytical results and carried out the numerical simulations. The author wrote the first version of the manuscript.

Author's Contribution

During his doctoral studies, the author has also contributed to other related articles [1, 2], excluded from this thesis.

1. Introduction

1.1 Motivation

The intersection of thermodynamics and quantum theory goes back to the very inception of the latter. The photon or “quantum of light”, hinted by Planck and later coined by Einstein, is born out of thermodynamic considerations [3, 4, 5, 6]. The field of quantum thermodynamics, as it is known today, is primarily focused on extending thermodynamic concepts, such as work, heat, and entropy, to the description of small quantum systems [7]. This effort has been motivated by the enormous technological advances in manipulating and measuring individual atoms and particles as well as the creation of artificial atoms in superconducting circuits [8, 9]. There is also a growing interest in understating and taking advantage of quantum effects in small machines such as heat engines or refrigerators [10, 11, 12, 13, 14, 15].

At first sight, this extension may seem troubling since thermodynamics, as presented in textbooks, describes systems with a large number of particles in (quantum or classical) thermal equilibrium [16]. There is no reason *a priori* why it should be valid outside the so-called thermodynamic limit. However, during the past 20 years theoretical and experimental developments have been providing mounting evidence to the contrary. The extension of thermodynamics to small systems comes in the form of fluctuation relations [17, 18, 19, 20] and sits within a general framework known as (classical) stochastic thermodynamics [21]. Despite some criticism [22], this approach has proven incredibly successful. In particular, it has provided a more detailed understating of the universal second law of thermodynamics at the level of individual microscopic processes in and out of equilibrium. Underlying this framework is the assumption that all the physical properties of the system, such as position, velocity or energy, have well-defined values at all times and that they are, in principle, measurable along individual runs of an experiment. However, for quantum systems, this assumption cannot be made, at least not in an obvious way. As a result, this problem has triggered a diverse number of contrasting attempts to extend the

successful picture of classical stochastic thermodynamic to the quantum regime. It is the author's opinion that one of the central questions is thus *how to correctly estimate the energy of an open quantum system for a single run of an experiment*. The results presented in this thesis give a small contribution towards answering this question.

1.2 Overview

In the first part of this dissertation, the focus is set on a promising new experimental protocol to measure work and heat dissipation in superconducting qubits¹ along individual runs of an experiment. The protocol, known as calorimetric detection, consists in measuring single quanta of energy exchanged between the qubit and its environment. It provides a way of directly and continuously measuring heat exchange. But to infer the work done on the qubit due to an external driving source, one has to rely on the fact that there are only two available energy levels. To which extent the method can be extended to more complex systems remains an open question. In Publication I this question is addressed for the particular case of a quantum harmonic oscillator². We show that under certain conditions the average work can be directly measured or inferred as if the system was a qubit. Another aspect of the calorimetric detection is that its low heat capacity comes from its small size which may induce non-Markovian evolution for the qubit. In Publication II, the amount of non-Markovianity induced by the calorimetric detection was studied using a previously developed model for the qubit-environment evolution. Some surprising results are reported, most notably that non-Markovianity does not necessarily decrease monotonically with increasing environment size.

The calorimetric detection scheme is an example of a continuously monitored system. But different experimental designs yield different answers. How heat and work are identified depends on how the experiment is designed and, more importantly, what is being measured [24]. The second part of this dissertation focuses on a more detailed study of energy fluctuations in quantum systems. A review of the different approaches on how to identify work in closed systems³ is first given. A recently reported no-go theorem is also reviewed which nicely encapsulates all these approaches and serves as a way of categorising them.

In Publication III we put forward a different approach based on the intersection of the Bohmian interpretation of quantum mechanics and the quantum Hamilton-Jacobi equation [25, 26, 27, 28]. Bohmian mechanics describes quantum phenomena based on actual point-particle trajectories and the quantum

¹A qubit is an idealised model of any system whose description is effectively captured by only two energy levels [23].

²A quantum harmonic oscillator is another idealised model for any system whose energy spectrum is composed of an infinite number of equidistant levels.

³In this context a *closed system* refers to a system evolving unitarily under a possibly time-dependent Hamiltonian according to the Schrödinger equation.

Hamilton-Jacobi equation tells us the Hamiltonian⁴ that generates them. With these two ingredients, the classical definition of work can be straightforwardly generalised to any closed quantum system. This generalisation retains two key features, namely, 1) it generates a proper (positive and normalised) probability distribution and 2) it respects the conservation of the average energy. However, it also provides new insights. In particular, if one considers proper mixtures of wave functions, the work distribution depends on the mixture itself. This fact may be somewhat surprising from the standard interpretation perspective, but it is in complete accordance with empirical facts. Moreover, since this approach is observer-independent [29], one can also treat measurements explicitly. This opens a new path to investigate the thermodynamic cost of quantum measurements [30, 31, 32, 33, 34, 35].

Finally, in Publication IV we study the contributions to the energy fluctuations of open quantum systems in single runs of an experiment. The description of individual runs of experiments dealing with open systems can be described in terms of stochastic wave functions resulting from the unravelling of the corresponding master equation [36]. However, what the stochastic wave functions represent is not entirely clear. The open system is, in general, entangled with its environment. Thus, in the standard interpretation, it cannot be described by a wave function of its own [29]. However, the Bohmian interpretation provides the concept of a conditional wave function as a way of ascribing a wave function to an open system in a clear conceptual manner [27, 37]. We leverage this approach to introduce the concept of conditional energy and identify three distinct contributions to its fluctuations — 1) an external contribution coming from the explicit time dependence of the Hamiltonian, 2) an interaction contribution that depends only on the interaction Hamiltonian and 3) an entanglement contribution which depends only on the entanglement between the system and its environment. The entanglement contribution is particularly interesting as it provides a direct link between entanglement and energy fluctuations at the level of individual runs of an experiment. These quantities are experimentally accessible via weak measurement schemes [38, 39], opening a possible new way of quantifying entanglement.

⁴The Hamiltonian here refers to a function of phase-space coordinates and time, not the Hamiltonian operator.

2. Open quantum systems

In this chapter, the basic concepts of open quantum systems underlying this work are outlined.

2.1 The reduced density operator of the system

In the theory of open quantum system one typically starts by considering the combined evolution of the (open) system and its environment. The environment contains all the degrees of freedom that we are not interested in but that are relevant to the evolution of the system. Let $\mathcal{H} = \mathcal{H}_S \otimes \mathcal{H}_E$ be the Hilbert space associated with the combined system and environment, where $\mathcal{H}_S$ is the Hilbert space of the system and $\mathcal{H}_E$ is the Hilbert space of the environment. The combined system is assumed to be described by a pure state $|\psi(t)\rangle \in \mathcal{H}$ and evolves according to the Schrödinger equation (SE) $i\partial_t|\psi(t)\rangle = \hat{H}(t)|\psi(t)\rangle$, where $\hat{H}(t)$ is the relevant Hamiltonian. Let $\hat{A} = \sum_k a_k |a_k\rangle\langle a_k|$ be an observable of the system where $\{|a_k\rangle\}_k$ are the eigenvectors of $\hat{A}$ and $\{a_k\}_k$ are the corresponding eigenvalues (assume no degeneracies for simplicity). If a measurement of $\hat{A}$ is performed at time t , the probability of obtaining the result a_k is given by [36],

$$p_k = \text{Tr} \left\{ \hat{\sigma}(t) (|a_k\rangle\langle a_k| \otimes \hat{1}_E) \right\} = \text{Tr}_S \left\{ \hat{\rho}(t) |a_k\rangle\langle a_k| \right\}, \quad (2.1)$$

where $\hat{1}_E$ is the identity operator in $\mathcal{H}_E$, $\hat{\sigma}(t) = |\psi(t)\rangle\langle\psi(t)|$ and $\hat{\rho}(t) = \text{Tr}_E \{\hat{\sigma}(t)\}$ is the *reduced density matrix* of the system. Equation (2.1) shows that all the information relevant to determine the probabilities of measurement outcomes of system observables is contained in the reduced density matrix. Rather than determining the evolution of $|\psi(t)\rangle$, which in general is not possible, we can instead find *approximate* solutions for the evolution of $\hat{\rho}(t)$. Finding and studying these solutions for a diversity of systems is one of the main objective in the field of open quantum systems.

2.2 The Markovian master equation

Let the Hamiltonian be of the form $\hat{H}(t) = \hat{H}_S(t) \otimes 1_E + 1_S \otimes \hat{H}_E + \hat{H}_{\text{int}}$, where $\hat{H}_S(t)$ is the bare system Hamiltonian acting only on the system Hilbert space, $\hat{H}_E$ is the environment bare Hamiltonian, acting only on the environment Hilbert space, and $\hat{H}_{\text{int}}$ is an interaction Hamiltonian that acts on both the system and the environment Hilbert spaces. It follows from the Liouville-von Neumann equation that the most general form for the evolution of the reduced density matrix is,

$$d_t \hat{\rho}(t) = -\frac{i}{\hbar} [\hat{H}_S(t), \hat{\rho}(t)] + \mathcal{D}[\hat{\sigma}(t)], \quad (2.2)$$

where $\mathcal{D}[\hat{\sigma}(t)] = -i \text{Tr}_E \{ [1_S \otimes \hat{H}_E + \hat{H}_{\text{int}}, \hat{\sigma}(t)] \} / \hbar$. In general, this equation cannot be solved exactly apart from some very specific cases. Thus, one needs to use effective descriptions that are approximations of Eq. (2.2), usually referred to as *master equations*. When dealing with a large thermal environment under weak coupling and a clear separation of time-scales, a Markovian master equation can be derived [36], namely,

$$d_t \hat{\rho}(t) = -\frac{i}{\hbar} [\hat{H}_{\text{eff}}(t), \hat{\rho}(t)] + \sum_{k=1}^N \Gamma_k \left(\hat{L}_k \hat{\rho}(t) \hat{L}_k^\dagger - \frac{1}{2} \{ \hat{L}_k^\dagger \hat{L}_k, \hat{\rho}(t) \} \right), \quad (2.3)$$

where $\Gamma_k \geq 0$ are dissipation rates, $\hat{L}_k$ are known as Lindblad or dissipation operators and $\hat{H}_{\text{eff}}$ is the effective Hamiltonian of the system and N is the total number of Lindblad operators. Eq. (2.3) is known as the *Gorini Kossakowski Sudarshan Lindblad* (GKSL) master equation. An effective Hamiltonian rather than the bare Hamiltonian $\hat{H}_S(t)$ is used because the interaction with the environment typically leads to a re-normalization of the bare Hamiltonian. It has been shown [40, 41] that any Markovian and linear dissipative system must necessarily be of this form. Because of this, this equation can be applied to effectively describe a variety of experimental scenarios and plays a central role in the study of open quantum systems.

2.3 Stochastic Schrödinger equations

Master equations such as Eq. (2.3) can be unravelled into *Stochastic Schrödinger equations* (SSE) of the form [36],

$$d|\phi(t)\rangle = -\frac{i}{\hbar} \left(\hat{H}_{\text{eff}}(t) + \frac{i}{2} \sum_{k=1}^N \Gamma_k \hat{L}_k^\dagger \hat{L}_k \right) |\phi(t)\rangle dt + \sum_{k=1}^N \left(\frac{\hat{L}_k |\phi(t)\rangle}{\|\hat{L}_k |\phi(t)\rangle\|} - |\psi(t)\rangle \right) dN_k(t), \quad (2.4)$$

where $dN_k(t)$ are independent Poisson increments satisfying $dN_k dN_l = dN_k \delta_{k,l}$ and $E[dN_k(t)] = \Gamma_k \|\hat{L}_k |\phi(t)\rangle\|^2 dt$, where $E[dN_k(t)]$ is the expectation value of

$dN(t)$. This is simply a multi-variable stochastic differential equation with Poisson noise. One can solve it numerically according to the following scheme. Consider a uniform time grid with an interval Δt . At each time step t , either

- (1) The state is updated smoothly according to the first term, with probability $p_0 = 1 - \sum_k \Gamma_k \|\hat{L}_k |\phi(t)\rangle\|^2 \Delta t$, such that the new state is given by $|\phi(t + \Delta t)\rangle = [1 - (i/\hbar)(\hat{H}_{\text{eff}}(t) + (i/2) \sum_{k=1}^N \Gamma_k \hat{L}_k^\dagger \hat{L}_k)] |\phi(t)\rangle \Delta t$, up to a normalization constant; or
- (2) The state undergoes a jump according to the second term, with probability $\Gamma_k \|\hat{L}_k |\phi(t)\rangle\|^2 \Delta t$, such that, the new state is given by $|\phi(t + \Delta t)\rangle = \hat{L}_k |\phi(t)\rangle / \|\hat{L}_k |\phi(t)\rangle\|$. Note that the condition $dN_k dN_l = dN_k \delta_{k,l}$ guarantees that only one $dN_k(t)$ is non-zero (and equal to one), at most.

Using this iteration scheme one can generate random trajectories of $|\phi(t)\rangle$ within a certain time period, say, for $t \in [0, T]$. Let $|\phi^{(j)}(t)\rangle$ denote one such trajectory. The reduced density matrix is then recovered as the statistical average of the trajectories $|\phi^{(j)}(t)\rangle$, i.e.,

$$\hat{\rho}(t) = E [|\phi(t)\rangle \langle \phi(t)|] := \lim_{M \rightarrow \infty} \frac{1}{M} \sum_{j=1}^M |\phi^{(j)}(t)\rangle \langle \phi^{(j)}(t)|, \quad (2.5)$$

where $E[\dots]$ represents the expectation value. In particular, one can show that the expectation value of Eq. (2.4) (converted to its density matrix representation) yields Eq. (2.3). Thus, Eq. (2.4) can be seen as a simple numerical tool to calculate the reduced density operator. For large systems this is more efficient than solving the Lindblad equation because it scales with N , rather than N^2 , where N is the number of degrees of freedom [42]. However, SSEs contain more information than the Lindblad equation. In particular, they can be used to interpret individual runs of experiments where either the system or the environment are being continuously monitored [23, 36].

2.4 Continuously monitored Markovian environment

A general and rigorous discussion on the use of SSEs to describe continuously monitored systems can be found in Refs. [23, 36]. Here, we outline how an SSE naturally captures the description of a continuously monitored environment. We note first that the word “continuously” is to be interpreted in the sense of a coarse-grained time-scale. Time coarse-graining is required to justify the Markovian assumption that at each point in time the system and environment are uncorrelated and their evolution depends only on the state at that time. In other words, the environment is measured at a finite rate $f = 1/\Delta t$, where the time between the measurements Δt is assumed to be large enough that the Markovian approximation is justified but much shorter than the timescale

in which the system evolves [36]. Now suppose that at time t the state of the system-environment is $|\phi(t)\rangle|\chi(t)\rangle$ and that at time $t + \Delta t$ a measurement of the environment is performed. Assume that there are $N + 1$ possible measurement outcomes. Of those, N correspond to the cases where the environment state has changed, *i.e.*, $\chi(t + \Delta t) \neq \chi(t)$, and one corresponds to the case where nothing happens, *i.e.*, $|\chi(t + \Delta t)\rangle = |\chi(t)\rangle$. In other words, there are N possible *events* that can be detected in addition to the *null event* where nothing happens to the environment. To give a concrete example, in the case of the calorimetric detection (see Chapter 3), there are two possible events, a photon was either absorbed or emitted by the calorimeter, in addition to the null event where neither absorption nor emission is observed. Further, we assume that when outcome k is observed at time $t + \Delta t$, the state is projected according to $|\phi(t + \Delta t)\rangle = \hat{L}_k |\phi(t)\rangle / \|\hat{L}_k |\phi(t)\rangle\|$. Then, the SSE in Eq. (2.4) provides the following interpretation. If the null outcome is observed, the state evolves smoothly according to the case (1) in the previous section. Otherwise, if event k is observed, the state jumps according to case (2) in the previous section. A trajectory with N jumps for $t \in [0, T]$ can be specified by a set of jump operators $\{\hat{L}_{k_j}\}_{j=1}^N$, and times $\{t_j\}_{j=1}^N$, with $t_j > t_{j-1}$, $t_1 > 0$ and $t_N < T$. The wave function at time T is then given by,

$$\begin{aligned} |\phi(t)\rangle &= \hat{U}_{\text{nh}}(T, t_N) \hat{L}_{k_N} \hat{U}_{\text{nh}}(t_N, t_{N-1}) \hat{L}_{k_{N-1}} \dots \hat{L}_{k_1} \hat{U}_{\text{nh}}(t_1, 0) |\phi(0)\rangle \\ &= \hat{V}_N(t_1, k_1; \dots; t_N, k_N) |\phi(0)\rangle \end{aligned} \quad (2.6)$$

where $\hat{U}_{\text{nh}}(t, t') = \mathcal{T}_> \exp((i/\hbar) \int_{t'}^t (\hat{H}_{\text{eff}}(s) - \iota \sum_{k=1}^N \Gamma_k \hat{L}_k^\dagger \hat{L}_k) ds)$ is the non-Hermitian evolution operator that dictates the evolution in between jumps. The operator $\hat{V}(t_1, k_1; \dots; t_N, k_N) := \hat{U}_{\text{nh}}(T, t_N) \hat{L}_{k_N} \hat{U}_{\text{nh}}(t_N, t_{N-1}) \hat{L}_{k_{N-1}} \dots \hat{L}_{k_1} \hat{U}_{\text{nh}}(t_1, 0)$ is a non-unitary evolution operator that can be used to compactly specify the N jump trajectory. Due to its non-unitary character, the squared norm of the wave function is not conserved. However, it encodes the probability for that particular trajectory to occur. That is, the probability for a particular trajectory specified by $\hat{V}(t_1, k_1; \dots; t_N, k_N)$ to occur is given by,

$$p(t_1, k_1; \dots; t_N, k_N) := \|\hat{V}(t_1, k_1; \dots; t_N, k_N) |\phi(0)\rangle\|^2. \quad (2.7)$$

Finally, note that Eq. (2.3) can equally well describe a scenario where no continuous monitoring is performed. One can still simulate the trajectories $|\phi^{(j)}(t)\rangle$ and recover the reduced density matrix. However, since no measurements are *actually* being performed, the system and environment will, in general, be entangled throughout their evolution. According to the standard interpretation, the system does not have a wave function of its own in this case and the proper way to interpret these trajectories becomes more opaque [29]. In Chapter 7, we review a possible approach to tackle these cases.

2.5 Non-Markovianity measures

At very low temperatures or strong coupling, the assumption that the environment relaxes much faster than the system cannot always be made and non-Markovian effects become important [43, 44, 45]. In general, the evolution of the system can no longer be captured by Eq. (2.3), but it has been recently shown that even the general evolution in Eq. (2.2) can be written in a “Lindblad-like” form, with the $\hat{L}_k$ operators now dependent on the evolution of the combined system and environment [46]. The study of non-Markovian open systems is not new (see, *e.g.*, Ref. [47]), but in more recent years considerable effort has been put into addressing non-Markovian systems from a quantum information perspective. This has led to the development of new research topics such as *reservoir engineering*, where the goal is to design environments to precisely tune dissipation and decoherence rates of the system of interest [48, 49, 50, 51, 52].

One significant problem that arises is how to determine the degree of non-Markovianity of the evolution, and different authors have proposed different *measures of non-Markovianity* [53, 54, 55, 56, 57, 58, 59]. One such measure is the Breuer-Laine-Pilo (BLP) one [58]. Let Φ_{t_2, t_1} be a completely positive, trace preserving (CPTP) map that determines the evolution of system, *i.e.*, $\hat{\rho}(t_2) = \Phi_{t_2, t_1}[\hat{\rho}(t_1)]$. Note that, in general, $\Phi_{t_2, t_1} \neq \Phi_{t_2, t_3} \Phi_{t_3, t_1}$, for $t_1 < t_3 < t_2$, *i.e.*, the map is not divisible. The BLP measure is then formally defined as,

$$\mathcal{N}(\Phi) = \max_{\rho_0^1, \rho_0^2} \int_{\mathcal{J} > 0} dt \mathcal{J}(t), \quad (2.8)$$

where $\hat{\rho}_0^1$ and $\hat{\rho}_0^2$ are two initial states and $\mathcal{J}(t) := D(\hat{\rho}^1(t), \hat{\rho}^2(t)) = \|\Phi_t(\hat{\rho}_0^1) - \Phi_t(\hat{\rho}_0^2)\|/2$ is the distinguishability of the two states defined as the trace distance between the two solutions at time t^1 and $\Phi_t := \Phi_{t, 0}$. The integral is taken over all regions where $\mathcal{J}(t) > 0$ and the maximisation is taken over all possible pairs of initial states. The BLP measure offers a physical interpretation regarding the information lost to the environment. For Markovian evolution, the trace distance between any two solutions always decreases. Thus, the ability to distinguish between the two solutions steadily decreases with time. In other words, information about the initial state is continuously lost to the environment. On the other hand, for non-Markovian evolution part of this information can flow back from the environment to the system. This backflow of information is what can be quantified by the BLP measure [48, 58]. Note that other measures can yield different results *e.g.* a particular map can be considered non-Markovian according to one measure but not by another. One advantage of the BLP measure is that, contrary to other measures, it allows an interpretation in terms of information flow between the system and the environment and it can be related to thermodynamic properties [48].

¹ The trace distance between two states $\hat{\rho}_1$ and $\hat{\rho}_2$ is defined as $\|\hat{\rho}_1 - \hat{\rho}_2\| := \text{Tr} \{ \sqrt{(\hat{\rho}_1 - \hat{\rho}_2)^2} \} / 2 = \sum_i |\lambda_i| / 2$, where λ_i are the eigenvalues of $\hat{\rho}_1 - \hat{\rho}_2$.

3. Calorimetric measurements of a driven harmonic oscillator

Calorimetric measurements are a promising experimental setup to study energy fluctuation in a continuously monitored open quantum system. In this Chapter, we first review how work and heat are measured when the protocol is applied to a qubit. We then summarize the results on a possible approach to measure fluctuations in an harmonic oscillator.

3.1 Qubit

Calorimetric measurements are a particular case of continuously monitored environments where the quantity being monitored is the temperature of the environment (hereafter referred to as the *calorimeter*). If both the temperature and heat capacity of the calorimeter are low enough, single quanta of energy exchanged with the qubit should be observable. The evolution of the system is modelled by a stochastic Schrödinger equation as described in Section 2.4. For the case of qubit with energy splitting $\hbar\omega$, there are two Lindblad operators, $\hat{L}_0 = \sqrt{\Gamma_0}\hat{\sigma}^{(-)}$ and $\hat{L}_1 = \sqrt{\Gamma_1}\hat{\sigma}^{(+)}$, where $\hat{\sigma}^{(-)}$ ($\hat{\sigma}^{(+)}$) is the lowering (raising) operator. If the calorimeter is in thermal equilibrium at inverse temperature β , the transition rates obey detailed balance and are given by $\Gamma_0 = \gamma(1 + N(\beta))$ and $\Gamma_1 = \gamma N(\beta)$, where γ is a parameter that determines the coupling strength between the system and calorimeter and $N(\beta) = 1/(\exp[\beta\hbar\omega] - 1)$ is the average thermal occupation number of the calorimeter¹. The operator $\hat{L}_0$ ($\hat{L}_1$) is associated with a photon absorbed (emitted) by the calorimeter and leaves the qubit in its ground (excited) state. In between jumps, the state evolves according to the non-Hermitian Hamiltonian $\hat{H}_{\text{nh}} = \hat{H}_{\text{eff}}(t) + \hat{D}$, where $\hat{H}_{\text{eff}}(t) = \hat{H}_S + \hat{H}_D(t)$, $\hat{H}_S = \hbar\omega\hat{\sigma}^{(+)}\hat{\sigma}^{(-)}$, $\hat{H}_D(t)$ is a time dependent driving Hamiltonian, assumed to be weak compared to $\hat{H}_S$, and $\hat{D} = \hat{L}_0\hat{L}_0^\dagger + \hat{L}_1\hat{L}_1^\dagger = \gamma(2N(\beta) + 1)\hat{\sigma}^{(+)}\hat{\sigma}^{(-)} + \gamma N(\beta)\hat{1}$.

Consider now a trajectory of N jumps from $t = 0$ to $t = T$ described by the set of jumps $\{i_1, \dots, i_N\}$ at times $\{t_1, \dots, t_N\}$, as described in Section 2.4. Heat is defined

¹The calorimeter is modelled as an infinite bath of harmonic oscillators.

as the net sum of quanta exchanged with the environment, namely,

$$Q^{(i)} = \hbar\omega \sum_{j=1}^{N+1} (-1)^{i_j}, \quad (3.1)$$

with the convention that positive heat corresponds to energy absorbed by the calorimeter. The change in energy $\Delta E^{(i)}$ is inferred from the the *guardian photons*, i.e., by looking at the last jump prior to the driving period and first jump after the driving period. This is possible because after each jump the system remains in the corresponding eigenstate if the system is not being driven. Let $\ell_i^{(i)} \in \{0, 1\}$ be the initial guardian photon for the i^{th} trajectory and $\ell_f^{(i)} \in \{0, 1\}$ the final guardian photon for the same trajectory. Then, $\Delta E^{(i)} = \hbar\omega(\ell_f^{(i)} - \ell_i^{(i)})$ and the work for i^{th} trajectory is defined as

$$W^{(i)} = \Delta E^{(i)} + Q^{(i)}. \quad (3.2)$$

Note that in Eq. (3.1), the last guardian photon is identified as the $N + 1$ jump ($i_{N+1} := \ell_f^{(i)}$), which must be included as heat.

3.2 Harmonic Oscillator

Consider now the case of a driven harmonic oscillator with Hamiltonian $\hat{H}_{\text{eff}}(t) = \hbar\omega \hat{a}^\dagger \hat{a} + \lambda_0 \sin(\omega t)(\hat{a}^\dagger + \hat{a})$, with $\lambda_0 \ll \hbar\omega$, where $\hat{a}$ ($\hat{a}^\dagger$) is the annihilation (creation) operator. Similarly to the qubit case, the evolution is described by the jump operators $\hat{L}_0 = \sqrt{\Gamma_0} \hat{a}$ and $\hat{L}_1 = \sqrt{\Gamma_1} \hat{a}^\dagger$ and the non-hermitian Hamiltonian $\hat{H}_{\text{nh}}(t) = \hat{H}_{\text{eff}}(t) + \hat{D}$, where $\hat{D} = \hat{L}_0 \hat{L}_0^\dagger + \hat{L}_1 \hat{L}_1^\dagger = \gamma(2N(\beta) + 1) \hat{a}^\dagger \hat{a} + \gamma N(\beta) \hat{1}$. While the heat can be measured in the same way, the change in internal energy can no longer be inferred from the guardian photons. After a jump, the state is not necessarily projected to an energy eigenstate if it wasn't in one before the jump. One way to remedy this situation is to apply the TPM protocol do have a well-defined initial and final energy. Let $E_i^{(i)} = \hbar\omega n_i^{(i)}$ and $E_f^{(i)} = \hbar\omega n_f^{(i)}$ be the energy outcomes of a projective energy measurement before and after the driving period for the i^{th} trajectory, respectively, where $n_i^{(i)}, n_f^{(i)} \in \{0, 1, 2, \dots\}$. Then, the work done under the two measurement protocol is defined as,

$$W_p^{(i)} = \Delta E_p^{(i)} + Q^{(i)}, \quad (3.3)$$

where $Q^{(i)}$ is defined as above (without any guardian photons) and $\Delta E_p^{(i)} = E_f^{(i)} - E_i^{(i)}$. We refer to this quantity as the *projective work*. However, projective measurements can be difficult to implement, especially if the system becomes more complex. Although the change in energy cannot be inferred from the guardian photons, they still contain relevant information.

3.2.1 The calorimeter two level approximation

The calorimeter protocol operates at low temperature. A possible approach is to consider a *two-level-approximation* (TLA) where we assume that the harmonic

Figure 3.1. The average and variance of the projective and calorimeter work for temperatures $\beta \hbar \omega = 1, 2$ and 5 . The parameters used are $\lambda_0 = 0.01 \hbar \omega$, $\hbar \gamma = 0.01 \lambda_0$ and $T = \hbar \pi / \lambda_0$.

oscillator is well approximated by its two lowest levels. Thus, we define,

$$W_c^{(i)} := \hbar \omega (\ell_f^{(i)} - \ell_i^{(i)}) + Q^{(i)}, \quad (3.4)$$

as the work measured in the calorimetric protocol under the two-level approximation. The last term is the contribution from the last guardian photon, which must be included as heat. We refer to $W_c^{(i)}$ as the *calorimeter work* to distinguish it from the projective work. We can now compare the moments of the distribution of W_c and W_p to determine to which extent this two level approximation can be used.

3.2.2 Results

Unitary limit

In the unitary limit, the driving period is short enough that no jumps occur and we can approximate the evolution as unitary. Thus, no heat is exchanged during the driving period and the work is determined solely by the guardian photons. At low temperature where only the two lowest levels of the harmonic oscillator are relevant in equilibrium we can write k^{th} moment of the calorimeter work as,

$$\langle W_c^k \rangle_u \approx (\hbar\omega)^k \frac{w_{0k} + w_{1k} e^{\beta\hbar\omega}}{1 + e^{-\beta\hbar\omega}}, \quad (3.5)$$

where the coefficients w_{nk} can be expressed in terms of hypergeometric functions (see Appendix A of Publication I). For the projective work, we find for the average and variance,

$$\begin{aligned} \langle W_p(t) \rangle_u &= \hbar\omega\mu(t) \\ [\sigma_{W_p}^2(t)]_u &= 2\hbar\omega(N(\beta) + 1/2)\langle W_p(t) \rangle_u, \end{aligned} \quad (3.6)$$

where $\mu(t) = \lambda_0^2 t^2/4$. Figure ?? shows the projective and calorimeter work at different temperatures.

We see that at low temperature and short driving periods both the average and variance of projective work and well approximated by the calorimeter work. For longer driving periods, higher levels of the harmonic oscillator will be populated and the TLA is no longer justified.

For higher temperatures, the average still follows a similar behavior to the low temperature case. However, the variance has a more subtle behavior. If the temperature is too high, a non-zero variance is present even in the limit of vanishing driving times. Naturally, this is an artifact coming from the wrong inference of the energy change. The TLA is no longer justified in this limit.

Single-jump correction

For longer driving times, jump correction have to be included. We can treat U_{nh} perturbatively and find corrections for trajectory with at most one jump. Using second-order perturbation theory, we write,

$$\hat{U}_{nh}(t) \approx e^{-\frac{i}{\hbar} \frac{\lambda_0 t}{\sqrt{2}} \hat{p}} \left[1 - \frac{i}{\hbar} \int_0^t dt_1 \hat{D}(t_1) - \frac{1}{\hbar^2} \int_0^t dt_1 \int_0^{t_1} dt_2 \hat{D}(t_1) \hat{D}(t_2) \right] \quad (3.7)$$

where $\hat{D}(t) = -i\gamma_\Sigma/2[\hat{H}'(t) + x_0(t)\hat{X}]$, $x_0(t) = \sqrt{2\mu(t)}$, $\hat{H}'(t) = \hat{a}^\dagger \hat{a} + \Gamma_1/(\Gamma_0 + \Gamma_1) + \mu(t)$, and $\hat{X} = (\hat{a}^\dagger + \hat{a})/\sqrt{2}$. We then use this expansion to calculate the operator $\hat{V}_0(T)$ and $\hat{V}_1(t_1, k_1)$ up to second-order in $\Gamma_0 + \Gamma_1$. Fig. 3.2 shows one example for $\beta\hbar\omega = 2$. We plot in solid lines the analytical results coming from the perturbation scheme and the markers show numerical results without approximation. As expected, the longer the system is driven, the worst the approximation. Note, however, that the error between the calorimeter and projective work for both the average and variance decreases, compared to the unitary limit.

Overdamped limit

For $\hbar\gamma \gg \lambda_0$ the perturbative method fails and we have to resort to numerical solutions. We note first that the average calorimeter work will correctly reproduce the average projective work. This is because the relaxation time to equilibrium is faster than the time required for the driving to push the system out of equilibrium. The system remains close to its equilibrium state with a negligible change in its internal energy. Thus, $\langle W_p \rangle \approx \langle W_c \rangle \approx \langle Q \rangle$. However, this

Figure 3.2. One-jump correction to the average and variance of the projective and calorimeter work for temperature $\beta \hbar \omega = 2$. The dashed line shows the unitary limit results and the solid lines show the analytical results using the perturbation technique. The symbols show the numerical solution without any approximation.

Figure 3.3. Numerical solutions for average and variance of the projective and calorimeter work for different couplings $\gamma/\omega = 0.01, 0.05$ and 0.1 with $\lambda_0/\hbar\omega = 0.01$ and $\beta\hbar\omega = 2$.

is no longer true for the variance and, in general, higher moments. We illustrate this in Fig. 3.3.

The results presented here show that, in certain limits, it is possible to use the calorimeter protocol to infer the distribution of work of a driven open harmonic oscillator.

4. Non-Markovianity due to driving and finite-size environment

The small size of the calorimeter required to detect single quanta events may lead to non-Markovian effects. Understanding to which extent one may need to take non-Markovian effects into account is important. For example, the calorimeter may no longer stay in thermal equilibrium with a well-defined temperature which would call into question how to correctly identify the heat.

4.1 The qubit-calorimeter model

Consider again a qubit with Hamiltonian $\hat{H}_c = \hbar\omega_0\hat{a}^\dagger\hat{a} + \lambda_0\sin(\omega_0t)(\hat{a}^\dagger + \hat{a})$ as our system of interest where $\hbar\omega_0$ is the energy gap, $\hat{a}$ is the lowering operator and λ_0 is the external driving strength. Recently, a finite environment master equation (FEME) was introduced to explicitly model the finite size of the environment in the calorimetric detection [1, 60]. In the FEME model, the calorimeter is modeled as a collection of N qubits with Hamiltonian $\hat{H}_c = \sum_{k=1}^N \hbar\omega_0\hat{a}_k^\dagger\hat{a}_k$, where $\hat{a}_k$ is the lowering operator. The model assumes that the calorimeter decoheres quickly into a microcanonical ensemble such that the state of the combined system and environment $\hat{\rho}_{qc}$ is given at all times by,

$$\hat{\sigma}_{qc}(t) = \sum_{n=0}^N \hat{\sigma}_q(n, t) \otimes \hat{\sigma}_c(E_n), \quad (4.1)$$

where n is the number of excited qubits in the calorimeter and $E_n = \hbar\omega n$ is the corresponding energy. The state $\sigma_c(E_n)$ is the microstate ensemble of the calorimeter eigenstates with energy E_n , i.e., $\hat{\sigma}_c(E_n) = \sum_k |\Psi_k\rangle\langle\Psi_k| \delta_{\epsilon_k, E_n} / N(E_n)$, where $|\Psi_k\rangle$ is an energy eigenstate of the calorimeter with energy ϵ_k and $N(E_n)$ is the total number of microstates with energy E_n . The reduced density operator of the system $\hat{\rho}(t)$ is given by tracing $\hat{\sigma}_{qc}(t)$ over the calorimeter, namely,

$$\hat{\rho}(t) = \sum_{n=0}^N \hat{\sigma}_q(n, t). \quad (4.2)$$

Note that $\hat{\sigma}_{extq}(n, t)$ are not reduced density operators because they are not normalized.

Treating the drive and interaction with the calorimeter perturbatively, it was shown that the operators $\hat{\rho}(n, t)$ obey the following set of equations,

$$\begin{aligned} \dot{\hat{\sigma}}_q(n, t) = & \frac{i}{\hbar} [\hat{\sigma}_q(n, t), \hat{H}_q(t)] - \frac{\Gamma_{\uparrow}(n)}{2} [\hat{\sigma}_q(n, t), \hat{a}\hat{a}^\dagger] + \Gamma_{\downarrow}(n-1) \hat{a}\hat{\sigma}_q(n-1, t) \hat{a}^\dagger \\ & + \Gamma_{\uparrow}(n+1) \hat{a}^\dagger \hat{\sigma}_q(n+1, t) \hat{a} - \frac{\Gamma_{\downarrow}(n)}{2} [\hat{\sigma}_q(n, t), \hat{a}^\dagger \hat{a}], \end{aligned} \quad (4.3)$$

where $\Gamma_{\uparrow}(n)$ and $\Gamma_{\downarrow}(n)$ are the transition rates associated with the Lindblad operators $\hat{a}^\dagger$ and $\hat{a}$, respectively. The transition rates depend on the calorimeter energy E_n and are expressed as

$$\Gamma_{\downarrow}(n) = g(1-n)/N; \quad (4.4)$$

$$\Gamma_{\uparrow}(n) = gn/N, \quad (4.5)$$

where g is a tunable parameter representing the interaction strength between the system and calorimeter. Because the transition rates depend on the state of the calorimeter, the evolution of the qubit depends on its history, which leads to non-Markovian behaviour. Note that the evolution of the combined system and calorimeter is Markovian [60].

4.2 BLP measure and initial conditions

To evaluate the degree of non-Markovianity we use the BLP measure introduced in Section 2.5. To have a well-defined map for the qubit, the initial state of the combined system and calorimeter has to be factorised, *i.e.*,

$$\hat{\sigma}_{qc}(0) = \hat{\rho}(0) \otimes \hat{\sigma}_c, \quad (4.6)$$

and the calorimeter state $\hat{\sigma}_c$ fixed. We take the initial state of the calorimeter as a thermal state at inverse temperature β , such that, $\hat{\sigma}_c := \hat{\sigma}_c^\beta = \sum_{n=0}^N p_n \hat{\sigma}_c(E_n)$, where $p_n = \binom{N}{n} \exp\{E_n - F\}$ and F is the free energy. This implies $\hat{\rho}(n, t) = p_n \hat{\rho}(0)$.

Let Φ_t be the map acting on the reduced density operator of the system, fixed by the above-choice of initial conditions. The set of equations in Eq. (4.3) and the trace operation in Eq. (4.2) are linear. Thus, it follows that the map Φ_t is also linear, *i.e.*, for any two initial states $\hat{\rho}^1(0)$ and $\hat{\rho}^2(0)$ and any two complex numbers c_1 and c_2 , we have that $c_1 \Phi_t(\hat{\rho}^1(0)) + c_2 \Phi_t(\hat{\rho}^2(0)) = \Phi_t(c_1 \hat{\rho}^1(0) + c_2 \hat{\rho}^2(0))$. Then, to evaluate the BLP measure, we only need to evolve the difference between pairs of initial states $\tilde{\rho}_0 = \hat{\rho}^1(0) - \hat{\rho}^2(0)$. It has been shown that for a qubit the pairs that maximizes the BLP measure are pure orthogonal states [61]. As such, we can restrict the search to pair differences of the form,

$$\tilde{\rho}_0 = \begin{pmatrix} \cos\theta & e^{i\phi} \sin\theta \\ e^{-i\phi} \sin\theta & -\cos\theta \end{pmatrix}, \quad (4.7)$$

where $\theta \in [0, \pi)$ and $\phi \in [0, \pi)$ are the usual Bloch sphere angles. The initial conditions to solve the set of equations in Eq. (4.3) are then $\rho(n, 0) = p_n \tilde{\rho}_0$.

Figure 4.1. Information flow of the qubit $\mathcal{I}_{\text{int}}(t)$, calorimeter $\mathcal{I}_{\text{ext}}(t)$ and their sum as a function of time. The parameters used are $\beta\hbar\omega = 2$, $\lambda_0/\hbar\omega = 0.08$, $g/\hbar\omega = 0.066$, $N = 20$, $\theta = 1.69$ and $\phi = 0$. Inset: Region of the trace distance time derivative that contributes to the BLP measure.

4.3 Trace distance

It is instructive to first look at the behavior of the information flow described by $\dot{\mathcal{I}}(t)$. Let $\hat{\sigma}_{\text{qc}}^1(t)$ and $\hat{\sigma}_{\text{qc}}^2(t)$ be two solutions with initial conditions $\hat{\rho}^1(0)$ and $\hat{\rho}^2(0)$, respectively, in accordance with Eq. (4.6). We now define two quantities, the *internal information* as $\mathcal{I}_{\text{int}}(t) := \mathcal{I}(t)$ and the *external information* as $\mathcal{I}_{\text{ext}}(t) := D(\hat{\sigma}_{\text{qc}}^1(t), \hat{\sigma}_{\text{qc}}^2(t)) - \mathcal{I}(t)$. Figure 4.1 shows these two quantities and their sum for a particular choice of initial conditions (see the caption for details).

We first note that if the combined qubit and calorimeter were isolated and their joint evolution unitary, the sum of the internal and external information would be constant. In other words, the information gained (lost) by the qubit must equal the information lost (gained) by the environment. However, in the FEME model, the combined calorimeter and qubit evolution is dissipative and Markovian. Thus, the total information $\mathcal{I}_{\text{int}}(t) + \mathcal{I}_{\text{ext}}(t)$ is constantly lost. We also see that there are discontinuities in the external information flow. These discontinuities appear because the trace distance is, in general, not differentiable everywhere. The same doesn't happen for the internal information because for a two-level system is a particular case where the trace distance is differentiable everywhere. The inset in Fig. 4.1 highlights one of the regions that contribute to the BLP measure. One can say that, during those periods of time, information flows back into the qubit. The BLP measure is calculated by maximising this

area over all possible initial qubit state pairs.

From an experimental perspective, it is important to know which parameters maximise or minimise the degree of non-Markovianity. In the next section, we look at the interplay between the calorimeter size N , the driving strength λ_0 and the coupling constant γ .

4.4 Influence of the system parameters on $\mathcal{N}(\Phi)$

Recall from Section 2.5 that we have introduced the quantity

$$\mathcal{N}(\Phi) = \max_{\rho_0^1, \rho_0^2} \int_{\dot{\mathcal{J}} > 0} dt \dot{\mathcal{J}}(t), \quad (4.8)$$

as the degree of non-Markovianity according to the BLP measure. We calculate $\mathcal{N}(\Phi)$ for three different calorimeter sizes $N = 5, 50$ and 100 and for $\lambda_0, g \in [0.01, 0.1]$, shown in Fig. 4.2. All results are calculated with $\beta\hbar\omega = 2$. Surprisingly, a non-Markovianity structure emerges characterized by “rays” of constant $\mathcal{N}(\Phi)$. This non-trivial structure shows that one cannot deduce that the degree of non-Markovianity will decrease by increasing the calorimeter size at constant λ_0 and γ . Indeed, one can go from Markovian behaviour to non-Markovian behaviour by increasing N . Another aspect is the fact that $\mathcal{N}(\Phi)$ vanishes when there is no driving. In this case, the set of equations in Eq. (4.3) can be solved exactly and we find,

$$\dot{\mathcal{J}}(t) = \sqrt{f^2(t)\cos^2\theta + g^2(t)\sin^2\theta}, \quad (4.9)$$

where $f(t) = (N \exp\{-g(1 + 1/N)t\} + 1)/(N + 1)$ and $g(t) = \exp\{-gt/2\}$. Since $f(t)$ and $g(t)$ are monotonically decreasing functions of time, $\dot{\mathcal{J}}(t) < 0$ for all t and $\mathcal{N}(\Phi) = 0$.

There are two figures of merit that might be useful in practical experimental scenarios, namely, 1) the maximum amount of non-Markovianity $\mathcal{N}_{\max}(N)$ at fixed size N and the parameters associated with it, and 2) the time at which information starts to flow back t_R (see Fig. 4.1). For the first case, $\mathcal{N}_{\max}(N)$ is shown with a dotted line in Fig. 4.2 and it's determined by the ratio $a_N = \lambda_0/g$ (see Fig. 4.3). From Fig. 4.3 we see that the maximum decreases monotonically with increasing N . It is also noticeable that the maximum amount of information flowing back into the system is not significant, even at very small calorimeter sizes $N \sim \mathcal{O}(10)$.

In the second case, the behavior of t_R is also not a monotonic function of the calorimeter size. However, if we focus on the parameters that maximize $\mathcal{N}(\Phi)$ a clear pattern emerges (see Fig. 4.4) and we can write,

$$t_R \approx \frac{\hbar}{\lambda_0}. \quad (4.10)$$

This relation shows that the recurrence time is independent of the calorimeter size when the parameters are tuned according to the ratio a_N . Thus, one

Figure 4.2. BLP measure ($10^2 \mathcal{N}(\Phi)$, color scale) as a function of λ_0 and g for $N = 5$ (top), 50 (middle) and 100 (bottom). The dashed lines indicate the line of maximum non-Markovianity determined by the ratio a_N , see the main text for details.

Figure 4.3. The ratio $a_N = \lambda_0/g$ that determines the maximum $\mathcal{N}_{\max}(N)$ as a function of N (top) and the dependence of $\mathcal{N}_{\max}(N)$ on the calorimeter size N (bottom).

can easily control t_R by adjusting the driving strength and coupling to the calorimeter.

4.5 Conclusions

In this Chapter, it was shown how the finite size of the calorimeter induces non-Markovian evolution for the qubit. The degree of non-Markovianity was quantified using the BLP measure applied to a previously developed model for the qubit-calorimeter system. Most noticeably, it was found that the degree of non-Markovianity doesn't necessarily decrease monotonically with the calorimeter size. Thus, by adjusting the driving strength or coupling between the qubit and the calorimeter, it is possible to mitigate or enhance non-Markovianity.

Figure 4.4. Recurrence time t_R for three different calorimeter sizes $N = 5, 50$ and 100 as a function of λ_0 . The coupling g is implicitly fixed by the ratio a_N , see the main text for details.

5. Energy fluctuations in closed quantum systems

In trying to extend the calorimetric protocol to a driven harmonic oscillator, we saw one of the difficulties in addressing energy fluctuations in quantum systems. The system can start and end in a state where its energy is not well-defined, according to the orthodox interpretation of quantum mechanics. At the same time, the energy fluctuations of its environment are continuously measured and have a well-defined value. It thus seem paradoxical to state that the system has exchanged energy with its environment without being able to even formally define how much energy was lost or gained by the system. This problem was circumvented by employing the two-point-measurement (TPM) protocol. Despite its versatility, however, this solution also has its drawbacks. In this Chapter, we expand on this topic by highlighting the difficulties in defining work in a closed quantum system. Some of the different approaches in the literature are briefly revised together with the no-go theorem for quantum work.

5.1 Motivation

To motivate the topic, consider a system at time $t = 0$ described by a state $\hat{\rho}$ which can be pure or mixed. During a period T , the system is driven unitarily by an external source according to a time-dependent Hamiltonian $\hat{H}_t$ such that the state at time T is given by $\hat{\rho}' = \hat{U} \hat{\rho} \hat{U}^\dagger$ where $\hat{U} = \mathcal{T}_> \exp\{i \int_0^T \hat{H}_t dt / \hbar\}$ and $\mathcal{T}_>$ is the time-ordering operator arranging the terms from right to left with increasing time. One then asks: How much work was done on the system by the external driving source in this *single run* of the experiment? The problem is, of course, that if $\hat{\rho}$ and $\hat{\rho}'$ do not correspond to eigenstates of $\hat{H}_0$ and $\hat{H}_T$, respectively, the system does not have a well-defined energy change according to the orthodox interpretation of quantum mechanics. This fact has lead to different approaches on how to answer this question.

5.2 Two-point measurement protocol

The two-point-measurement (TPM) protocol was first introduced by Tasaki [62] and Kurchan [63] and has been extensively used in the context of, *e.g.*, fluctuation relations [62, 63, 64, 65, 66, 67], heat engines [10], many-body systems [68], relativistic systems [69] and open quantum systems [70, 71, 72]. It consists of the following procedure: 1) Measure $\hat{H}_0 = \sum_i e_i |e_i\rangle\langle e_i|$ at time 0, resulting in the eigenstate $|e_i\rangle$ with energy e_i , with probability $p_i = \langle e_i | \hat{\rho} | e_i \rangle$. 2) Evolve $|e_i\rangle$ using the unitary $\hat{U}$. 3) Measure $\hat{H}_T = \sum_j e'_j |e'_j\rangle\langle e'_j|$ at time T , resulting in the eigenstate $|e'_j\rangle$ with energy e'_j , with probability $p_{j|i} = \langle e'_j | \hat{\rho}' | e'_j \rangle = |\langle e'_j | \hat{U} | e_i \rangle|^2$. The work is then defined to be the difference in the energy outcomes, *i.e.*,

$$w_{i,j}^{\text{tpm}} := e'_j - e_i.$$

This corresponds to a random variable W with distribution $P^{\text{tpm}}(W = w) = \sum_{i,j} w \delta_{w, w_{i,j}^{\text{tpm}}}$. The TPM protocol is particularly attractive because of its simplicity of implementation and interpretation. Further, it satisfies both Jarzynski's and Crooks' fluctuation relations [67]. This makes it a very useful quantity to study out of equilibrium fluctuations, in particular, if the initial state is thermal. This protocol, however, induces an evolution different from the undisturbed process described in Section 5.1. It is the state $|e_i\rangle$ resulting from the first measurement that undergoes the unitary evolution, rather than the state $\hat{\rho}$. A consequence of this can be seen in the average work. In general, the average work given by the TPM protocol differs from the change in the average energy of the undisturbed process, *i.e.*,

$$\langle W \rangle_{\text{tpm}} = \sum_w P^{\text{tpm}}(W = w) w \neq \text{Tr} \{ \hat{\rho} \hat{H}_0 \} - \text{Tr} \{ \hat{\rho}' \hat{H}_T \}. \quad (5.1)$$

This situation arises when energy coherences are present in the initial state, *i.e.*, $[\hat{\rho}, \hat{H}_0] \neq 0$. Different authors have shown that one can extract useful work from these initial coherences [31, 73, 74, 75, 76]. In this sense, the TPM seems inadequate to characterize these processes. Several authors have also noted that during the measurement steps the system becomes open and the evolution is not unitary [31, 77, 78]. Thus, the identification of the change in energy $\Delta e = e'_j - e_i$ with work is not straightforward since one should account for the energy exchanged with the measuring device.

5.3 Other approaches

An earlier approach to study work fluctuation was based on an operator of work [79, 80, 81, 82], defined as $\hat{W} := \hat{U}^\dagger \hat{H}_T \hat{U} - \hat{H}_0 = \sum_w w |w\rangle\langle w|$. The work distribution is then associated with the spectrum of $\hat{W}$, *i.e.*, $P^{\hat{W}}(W = w) = \text{Tr} \{ \rho |w\rangle\langle w| \}$. This produces a positive distribution, by construction, and respects the change in average energy of the undisturbed process. The spectrum

of $\hat{W}$ is interpreted as the possible work values along individual runs. This, however, leads to somewhat contradictory results [82]. Consider the case of a cyclic driving protocol, such that, $\hat{H}_T = \hat{H}_0$ but $[\hat{U}^\dagger \hat{H}_T \hat{U}, \hat{H}_0] \neq 0$. Then, it is possible to find an eigenstate $|0\rangle$ of $\hat{W}$ with $w = 0$ such that

$$\langle 0 | (\hat{U}^\dagger \hat{H}_T \hat{U})^k | 0 \rangle \neq \langle 0 | \hat{H}_0^k | 0 \rangle, \quad k > 2. \quad (5.2)$$

On one hand, according to the operator of work, if starting from the state $|0\rangle$, the work done is zero with probability one. But on the other hand, moments of the energy distribution higher than the second have changed, implying that the work distribution should have non-zero variance. The operator of work has also been criticised on the fact that work should describe a process, not a state, and thus it cannot be described by an observable [64].

A different class of approaches involves quasi-probabilities and are based on, *e.g.*, full-counting-statistics [83], the consistent-histories interpretation [84] or weak values [85]. While these approaches can be useful in identifying purely quantum features of certain protocols [86], the fact that they do not form a probability distribution means that it is mathematically impossible to construct an underlying random variable. Thus, it is not clear how to interpret the “moments” of the distribution nor how to assign a definitive value to an individual run of an experiment. A thorough discussion on the different approaches can be found in Ref. [2].

5.4 A “no-go” theorem from quantum work

Recently, a “no-go” theorem has been proven showing that no work distribution in closed quantum systems can simultaneously satisfy the following three requirements [87], namely:

1. The work distribution is described by a positive operator valued measurement (POVM), *i.e.*, the probability of having done an amount of work w given an initial state $\hat{\rho}$ and driving protocol $\hat{U}$ is given by $p(w|\hat{\rho}, \hat{U}) = \text{Tr} \{ \hat{\rho} \hat{M}_w \}$, with the condition $\sum_{\{w\}} \hat{M}_w = \hat{1}$, where the set $\{w\}$ represents all possible work values.
2. For initial states that commute with the initial Hamiltonian the work distribution reduces to the TPM work distribution, *i.e.*, given an initial state $\hat{\rho}_D$ such that $[\hat{\rho}_D, \hat{H}_0] = 0$, then $p(w|\hat{\rho}_D, \hat{U}) = \sum_{i,j} p_i p_{j|i} \delta_{e'_j - e_i, w}$, where $p_i = \langle e_i | \hat{\rho}_D | e_i \rangle$ and $p_{j|i} = |\langle e'_j | \hat{U} | e_i \rangle|^2$ (see Section 5.2).
3. The work distribution respects the change in the Hamiltonian expectation value, *i.e.*, $\sum_{\{w\}} w p(w|\hat{\rho}, \hat{U}) = \text{Tr} \{ \hat{\rho}' \hat{H}_T \} - \text{Tr} \{ \hat{\rho} \hat{H}_0 \}$.

As an example, the TPM protocol satisfies 1 and 2 but not 3, the operator of work satisfies 1 and 3 but not 2 and quasi-probabilities don't satisfy 1. Note that there can be a set of POVM operators $\{\hat{M}_w\}$ that satisfy all three properties for some specific choices of $\hat{\rho}$ and $\hat{U}$. What is proven is that there is no such POVM that satisfies all three properties for all $\hat{\rho}$, $\hat{U}$, $\hat{H}_T$ and $\hat{H}_0$.

Like all no-go theorems, this result can be read in different ways [88]. On one hand, we can accept all three requirements and conclude that no work distribution can be assigned to a closed system evolution. On the other hand, one can re-think what should be the requirements for the work distribution in a closed quantum system. For example, the first requirement implies a positive distribution but the converse is not true. One can have a positive distribution that is not described by a POVM. In Chapter 6 and Publication III we present a new approach where this is the case.

6. Quantum Work in the Bohmian Framework

One of the difficulties encountered in defining work in quantum systems is that the energy of the system is not always well-defined. One is then forced to consider operational approaches, where the measurement protocol itself is used to define work. However, in the Bohmian interpretation of quantum mechanics, one can construct a well-defined and observer-independent energy for the system based on the so-called quantum Hamilton-Jacobi equation. This formulation leads to a natural extension of the classical work definition to the quantum regime.

6.1 Bohmian Mechanics

Bohmian Mechanics (BM) is a non-local, realist interpretation of quantum experiments [89]. Its inception dates back to de Broglie [90, 91], who has put forward the idea that quantum experiments could be explained by point-particles guided by a pilot-wave. This idea was later generalised by Bohm and others [25, 26, 89] into what is commonly known today as *Bohmian Mechanics*. This theory yields the same predictions as the orthodox formulation, as far as outcomes of measurements are concerned, but provides an objective, observer-independent description of quantum phenomena. Bohmian Mechanics has been applied to a variety of different problems [92, 93], from cosmology [94] to chemistry (see, *e.g.*, Ref [95]), with different motivations, such as, the development of accurate and efficient numerical methods [96]. In the next sections, a brief introduction to the mathematical formulation of BM is provided, before discussing its application to define work in a thermodynamics context.

6.1.1 Description of experiments

In the non-relativistic formulation of BM, systems are modelled by a collection of N point-particle positions $\vec{z}_1, \dots, \vec{z}_N$, $\vec{z}_i \in \mathbb{R}^3$ and a wave function $\Psi(\vec{z}, t)$, where $\vec{z} := (\vec{z}_1, \dots, \vec{z}_N) \in \mathbb{R}^{3N}$ is the configuration vector. One typically works with the

wave function in the position representation, written in polar form,

$$\Psi(\vec{z}, t) = \langle \vec{z} | \Psi(t) \rangle = R^{(\Psi)}(\vec{z}, t) \exp\{i S^{(\Psi)}(\vec{z}, t)/\hbar\}, \quad (6.1)$$

where $|\vec{z}\rangle$ is an element of the position basis and $R^{(\Psi)}(\vec{z}, t) \in \mathbb{R}_0^+$ and $S^{(\Psi)}(\vec{z}, t) \in \mathbb{R}$ are the modulus and phase of the wave function at point $\vec{z}$ and t , respectively.

In BM, experiments are described in the following way. One begins by preparing an initial state $\Psi(\vec{z}, 0)$ for which the system is in a well-defined, but random, initial configuration $Z := (Z_1, \dots, Z_N)^1$. The probability that the system is in a configuration $\vec{z}$ inside an infinitesimal volume element $d\mathbf{z}$ is given by $\mathbb{P}_0(\vec{z}) d\mathbf{z} := (R^{(\Psi)})^2(\vec{z}, 0) d\mathbf{z}$, i.e., the Born rule. A given protocol described by a Hamiltonian $\hat{H}_t$ is then implemented and the system configuration evolves deterministically. If measurements are carried out, they are treated like any other interaction by including the configuration of the measuring apparatus itself in the description. The combined system and measuring apparatus evolve deterministically and the random outcome is traced back to our lack of knowledge regarding the initial configuration Z , now both of the system and measuring apparatus. When the experiment is repeated, a new initial configuration Z' is drawn from $\mathbb{P}_0$ which may lead to a different measurement outcome. A detailed discussion showing how the standard description of measurements and the notion of *effective collapse* and *effective wave functions* arises from this picture can be found in Refs. [27, 98].

6.1.2 Dynamics

Suppose that the system Hamiltonian is of the form,

$$\hat{H}(t) = \sum_i \frac{\hat{P}_i^2}{2m_i} + \hat{V}(\hat{Z}, t), \quad (6.2)$$

where m_i is the mass of particle i , $\hat{P}_i$ is the three-dimensional momentum operator of particle i , defined by $\langle \vec{z} | \hat{P}_i | \Psi(t) \rangle = (-i \nabla_i \Psi)(\vec{z}, t)$ and $\hat{V}(\hat{Z}, t)$ is the potential energy operator dependent only on the position of the particles and time t , such that, $\langle \vec{z} | \hat{V}(\hat{Z}, t) | \Psi(t) \rangle = V(\vec{z}, t) \Psi(\vec{z}, t)$, $V(\vec{z}, t) \in \mathbb{R}$. Plugging Eq. (6.1) into the Schrödinger equation (SE) $i \partial_t \Psi(\vec{z}, t) = \hat{H}(t) \Psi(\vec{z}, t)$ and equating the real and imaginary parts, we find,

$$\partial_t \rho^{(\Psi)}(\vec{z}, t) = - \sum_{i=1}^N \nabla_i (\rho^{(\Psi)}(\vec{z}, t) \vec{v}_i(\vec{z}, t)); \quad (6.3)$$

$$-\partial_t S^{(\Psi)}(\vec{z}, t) = \sum_{i=1}^N \frac{[\nabla_i S^{(\Psi)}(\vec{z}, t)]^2}{2m_i} + V(\vec{z}, t) + \mathcal{Q}^{(\Psi)}(\vec{z}, t), \quad (6.4)$$

¹It may also be the case that one does not know the initial wave function but only has access to the reduced density operator, corresponding to a mixed state. In this case, a similar description can be provided using an appropriate coarse-graining scheme [97].

² ∇_i is the gradient operator with respect to particle i position.

where $\rho^{(\Psi)}(\vec{z}, t) := (R^{(\Psi)})^2(\vec{z}, t)$,

$$\mathcal{Q}^{(\Psi)}(\vec{z}, t) = - \sum_{i=1}^N \frac{\hbar^2}{2m_i} \frac{\nabla_i^2 R^{(\Psi)}(\vec{z}, t)}{R^{(\Psi)}(\vec{z}, t)} \quad (6.5)$$

is the *quantum potential* and

$$\vec{v}_i^{(\Psi)}(\vec{z}, t) = \frac{\nabla_i S^{(\Psi)}(\vec{z}, t)}{m_i} = \frac{\hbar}{m_i} \text{Im} \frac{(\nabla_i \Psi)(\vec{z}, t)}{\Psi(\vec{z}, t)} \quad (6.6)$$

is a velocity field, also known as the *guidance equation*. Equation (6.3) is a continuity equation and Eq. (6.4) is the so-called quantum Hamilton Jacobi (HJ) equation. The decomposition into Eqs. (6.3) and (6.4) shows that the SE equally well describes the evolution of an ensemble of point particles —Eq. (6.3) describes the evolution of the ensemble distribution and Eq. (6.4) describes the motion of the individual point particles in the ensemble. According to HJ theory, the energy of the system for a configuration $\vec{z}$ at time t is given by,

$$E^{(\Psi)}(\vec{z}, t) = -\partial_t S^{(\Psi)}(\vec{z}, t), \quad (6.7)$$

and the momentum of the particle i by,

$$\vec{p}_i^{(\Psi)}(\vec{z}, t) = \nabla_i S^{(\Psi)}(\vec{z}, t). \quad (6.8)$$

Formally, the trajectories of each particle $\vec{Z}_i^{(\Psi)}(t|Z)$ given an initial condition Z are given by integrating the velocity field in Eq. (6.6), namely,

$$\vec{Z}_i^{(\Psi)}(t|Z) = Z_i + \int_0^t ds \vec{v}_i^{(\Psi)}(\vec{Z}^{(\Psi)}(s|Z), s). \quad (6.9)$$

Note that these trajectories depend implicitly on the distribution of the ensemble via the quantum potential. In Fig. 6.1 we illustrate three possible trajectories for a driven harmonic oscillator, initially in its ground state (see the example in Section 6.4).

6.2 Connection to the Aharonov-Albert-Vaidman weak values

It is interesting to note that there is a close relationship between the Bohmian trajectories and the Aharonov-Albert-Vaidman (AAV) weak values and measurements [38, 39]. The AAV weak value (hereafter simply referred to as weak value) of an operator $\hat{O}$ with pre-selection on $|\psi\rangle$ and post-selection on $|\phi\rangle$ is defined by,

$$\phi\langle\hat{O}\rangle_\psi := \frac{\langle\phi|\hat{O}|\psi\rangle}{\langle\phi|\psi\rangle}. \quad (6.10)$$

Taking $|\phi\rangle = |\vec{z}\rangle$ and $|\psi\rangle = |\Psi\rangle$, we see that the momentum given by Eq. (6.8) is the real part of the weak value of momentum post-selected on position, *i.e.*,

$$\vec{p}_i^{(\Psi)}(\vec{z}, t) = \Re \left\{ \frac{\langle\vec{z}|\hat{P}_i|\Psi\rangle}{\langle\vec{z}|\Psi\rangle} \right\} = \Re \{ \vec{z}\langle\hat{P}\rangle_\psi \}. \quad (6.11)$$

Figure 6.1. Illustration of three possible trajectories for a driven harmonic oscillator, initially in the ground state (see the example in Section 6.4 for more details). For a given run of the experiment, the particle starts at z_0 with initial energy $E^{(\Psi)}(z_0, 0)$ and ends up at $z_t := Z^{(\Psi)}(t|z_0)$ with energy $E^{(\Psi)}(z_t, t)$. At any time point s during the evolution, its energy is always well-defined and given by $E^{(\Psi)}(z_s, s)$.

A similar relation exist for the HJ energy in Eq. (6.7), namely,

$$E^{(\Psi)}(\vec{z}, t) = \Re \left\{ \frac{\langle \vec{z} | \hat{H}(t) | \Psi \rangle}{\langle \vec{z} | \Psi \rangle} \right\} = \Re \left\{ \vec{z} \langle \hat{H}(t) \rangle_{\Psi} \right\}. \quad (6.12)$$

All these quantities are thus experimentally accessible.

Note that, for an energy eigenstate $|e\rangle$, $E^{(e)}(\vec{z}, t) = e$ such that the system has energy e , independent of its configuration. Thus, from a Bohmian perspective, when a strong energy measurement is performed on a state $|\Psi\rangle$, resulting in an energy eigenstate $|e\rangle$, what is revealed is the energy of the system *after* the measurement took place, as opposed to the energy $E^{(\Psi)}(\vec{z}, t)$ prior to the measurement [27, 99]. The latter can only be inferred through weak measurements. An analogous argument is valid for a momentum measurement.

6.3 Work in closed systems

The HJ theory also provides a *Hamiltonian function* in phase space $\mathcal{H}^{(\Psi)}(\vec{z}, \vec{p}, t)$ that generates the trajectories through Hamilton's equation of motion, namely,

$$\mathcal{H}^{(\Psi)}(\vec{z}, \vec{p}, t) = \sum \frac{p_i^2}{2m_i} + V(\vec{z}, t) + \mathcal{Q}^{(\Psi)}(\vec{z}, t). \quad (6.13)$$

The exact same trajectories in Eq. (6.9) will be generated by this Hamiltonian, provided the same initial conditions $(Z, P) = (Z, \nabla S^{(\Psi)}(Z, 0))$ are given, where $\nabla = (\nabla_1, \dots, \nabla_N)$. Accordingly, the same statistics concerning any observation are recovered if the system has the initial phase-space distribution,

$$f_0^{(\Psi)}(\vec{z}, \vec{p}) = |\Psi(\vec{z}, 0)|^2 \delta(\vec{p} - \nabla S^{(\Psi)}(\vec{z}, 0)). \quad (6.14)$$

Additionally, we can also explicitly prepare a different initial state for each run of the experiment, *i.e.*, a proper statistical mixture of pure states. For this case,

we would also have a distribution over the initial possible wave functions $\{\Psi_j\}_j$ with probabilities $\{p_j\}_j$ ($p_j \geq 0$, $\sum_j p_j = 1$)³. The initial phase space distribution is obtained by summing over this distribution, namely,

$$f_0^{\{\Psi_j\}}(\vec{z}, \vec{p}) = \sum_j p_j f_0^{(\Psi_j)}(\vec{z}, \vec{p}). \quad (6.15)$$

Note that we can have two different statistical mixtures $\{\Psi_j\}_j$ and $\{\Phi_i\}_i$ that correspond to two different initial phase space distributions $f_0^{\{\Psi_j\}}$ and $f_0^{\{\Phi_i\}}$ but to the same density operator $\hat{\rho} = \sum_j p_j |\Psi_j\rangle\langle\Psi_j| = \sum_i p_i |\Phi_i\rangle\langle\Phi_i|$. These two distributions are experimentally accessible as long as the two mixtures are explicitly prepared (see also Ref. [36], Chapter 5).

We now have a description of the evolution of pure states in terms of phase-space trajectories generated by a Hamiltonian function $\mathcal{H}^{(\Psi)}$. As such, we can extend the classical definition to the quantum case as,

$$\begin{aligned} W^{(\Psi)}[\vec{Z}^{(\Psi)}(s|Z)] &= \int_0^t \partial_s \mathcal{H}^{(\Psi)}|_{\vec{Z}_s, s} = \int_0^t \mathrm{d}s E^{(\Psi)}|_{\vec{Z}_s, s} \mathrm{d}s = E^{(\Psi)}(\vec{Z}_t, t) - E^{(\Psi)}(\vec{Z}_0, 0) \\ &:= W^{(\Psi)}(t, Z), \end{aligned} \quad (6.16)$$

where we have used the shorthand notation $\vec{Z}_s := \vec{Z}^{(\Psi)}(s|Z)$. Since $\vec{Z}_t$ depends only on Ψ and Z , the work is completely determined by the initial configuration and wave function. The work distribution is given by,

$$P^{(\Psi)}(w) = \int \mathrm{d}\mathbf{z} |\Psi(\vec{z}, 0)|^2 \delta(w - W^{(\Psi)}(t, z)), \quad (6.17)$$

and for the case of a proper mixture of initial states by,

$$P^{\{\Psi_j\}}(w) = \sum_j p_j P^{(\Psi_j)}(w). \quad (6.18)$$

This distribution is positive semi-definitive, normalized and respects the conservation of average energy, *i.e.*, $\langle W \rangle_{\{\Psi_j\}} := \int \mathrm{d}w P^{\{\Psi_j\}}(w) w = \mathrm{Tr} \{ \hat{\rho}(t) \hat{H}_t \} - \mathrm{Tr} \{ \hat{\rho}(0) \hat{H}_0 \}$, where $\hat{\rho}(\tau) = \sum_j p_j |\Psi_j(\tau)\rangle\langle\Psi_j(\tau)|$ is the density operator associated with the proper mixture. Note that the average is independent of the mixture but higher moments will, in general, depend on the mixture.

It is worth emphasize that the work distribution in Eq. (6.18) is not captured by a POVM. As such, there is no incompatibility with the no-go theorem presented in Section 5.4.

6.4 Example: Driven harmonic oscillator

Consider the case of a driven harmonic oscillator with Hamiltonian,

$$\hat{H}(t) = \frac{\hat{P}^2}{2m} + \frac{1}{2} m \omega^2 \hat{X}^2 - \hat{Z} f_1(t) + \hat{P} f_2(t), \quad (6.19)$$

³We take the distribution to be discrete for simplicity's sake.

where m and ω are the oscillator mass and angular frequency, respectively, $\hat{Z}$ ($\hat{P}$) is the position (momentum) operator and

$$f_1(t) = -A \sin(\omega t), \quad f_2(t) = -\frac{A}{m\omega} \cos(\omega t), \quad (6.20)$$

where A is the driving amplitude. The eigenstates of this Hamiltonian are given by $|\tilde{n}_\alpha\rangle := D(\alpha)|n\rangle$, where $|n\rangle$ is an eigenstate of the undriven Hamiltonian $\hat{H}_0 = \hat{P}^2/2m + m\omega^2\hat{Z}^2/2$, $D(\alpha) := \exp\{\alpha\hat{a}^\dagger - \alpha^*\hat{a}\}$ is the displacement operator and $\alpha = -iA/\sqrt{2\hbar m\omega^3}$. The energy levels are given by $E_n = \hbar\omega(n + 1/2 + |\alpha|^2)$. The evolution operator associated with this Hamiltonian is given by,

$$\hat{T}(t, 0) = \exp\left\{\frac{At}{\sqrt{2\hbar m\omega}}[-\hat{a}e^{i\omega t} + \hat{a}^\dagger e^{-i\omega t}]\right\} e^{i\hat{H}_0 t/\hbar}, \quad (6.21)$$

where $\hat{a}$ ($\hat{a}^\dagger$) is the annihilation (creation) operator. Decomposing the SE into the continuity and HJ equations, we find that the velocity field is given by,

$$v(z, t) = \frac{1}{m} \partial_x S^{(\Psi)}(z, t) - f_2(t), \quad (6.22)$$

where $S^{(\Psi)}$ is again the global phase of the wave function.

6.4.1 Thermal state as a mixture of energy eigenstates

Consider an initially prepared mixture of energy eigenstates,

$$\{|\tilde{n}_\alpha\rangle_{n=0}^\infty, \quad p_n = \exp\{-\beta\hbar\omega n\}(1 - \exp\{-\beta\hbar\omega\}), \quad (6.23)$$

such that $\hat{\rho}_\beta = \exp\{-\beta\hat{H}(0)\}/Z_\beta = \sum_{n=0}^\infty p_n |\tilde{n}_\alpha\rangle\langle\tilde{n}_\alpha|$, where β is the inverse temperature and Z_β the partition function. The S -function for each eigenstate is given by,

$$\begin{aligned} S^{(\tilde{n}_\alpha)}(z, t) = & -\hbar\omega \left(n + \frac{1}{2}\right) t + \frac{A^2 t}{2m\omega^2} - \frac{A}{\omega} z [\cos(\omega t) + \omega t \sin(\omega t)] \\ & + \frac{A^2}{4m\omega^3} [2\omega t \cos(2\omega t) + (\omega^2 t^2 - 1) \sin(2\omega t)]. \end{aligned} \quad (6.24)$$

From Eq. (6.22) we can determine the trajectories $Z^{(\tilde{n}_\alpha)}(t|Z)$ and, plugging the result into Eq. (6.16), we find the work along and individual trajectories to be

$$W^{(\tilde{n}_\alpha)}(t, Z) = \frac{A\tau[A\tau + 2mZ\omega \cos(\omega\tau)]}{2m}. \quad (6.25)$$

Thus, the work distribution *given* that the initial state is $|\tilde{n}_\alpha\rangle$ is

$$P^{(\tilde{n}_\alpha)}(w) = \int dZ |\langle Z|\tilde{n}_\alpha\rangle|^2 \delta(w - W^{(\tilde{n}_\alpha)}(t, Z)). \quad (6.26)$$

The distribution associated with the eigenstate mixture is then given by

$$P^{(\tilde{n}_\alpha)}(w) = \sum_{n=0}^\infty p_n P^{(\tilde{n}_\alpha)}(w). \quad (6.27)$$

6.4.2 Thermal state as a mixture of coherent states

Consider now an initially prepared mixture of coherent states (eigenstates of the annihilation operator),

$$\{\eta_\alpha\}_\eta, \quad p_\alpha(\eta) = \frac{(e^{\beta\hbar\omega} - 1)}{\pi} \exp\left\{-|\eta - \alpha|^2 [e^{\beta\hbar\omega} - 1]\right\}, \quad (6.28)$$

such that $\hat{\rho}_\beta = \exp\{-\beta\hat{H}(0)\}/Z_\beta = \int d^2\eta p_\alpha(\eta)|\eta\rangle\langle\eta|$. The S -function for each coherent state is given by,

$$\begin{aligned} S^{(\eta)}(z, t) = & -\frac{1}{2}\hbar\omega t + \frac{\hbar A t \eta_I}{\sqrt{2\hbar m \omega}} - z \left[(A t + \eta_R \sqrt{2\hbar m \omega}) \sin(\omega t) - \eta_I \sqrt{2\hbar m \omega} \cos(\omega t) \right] \\ & - \hbar \left(\eta_R + \frac{A t}{\sqrt{2\hbar m \omega}} \right) \eta_I \cos(2\omega t) \\ & + \frac{\hbar}{2} \left[\left(\eta_R + \frac{A t}{\sqrt{2\hbar m \omega}} \right)^2 - [\eta_I]^2 \right] \sin(2\omega t), \end{aligned} \quad (6.29)$$

where η_R (η_I) is the real (imaginary) part of η . Similarly to the previous case, we determine the trajectories $Z^{(\eta)}(t|Z)$ from Eq. (6.22), and find the work to be given by

$$\begin{aligned} W(t, Z) = & A\tau \left[\frac{A\tau}{2m} + \sqrt{\frac{2\hbar\omega}{m}} \eta_R \right] + \{A[\omega\tau \cos(\omega\tau) + \sin(\omega\tau)] \\ & + \sqrt{2\hbar m \omega^3} [\eta_R(\cos(\omega\tau) - 1) + \eta_I \sin(\omega\tau)]\} \\ & \times \left[Z - \eta_R \sqrt{\frac{2\hbar}{m\omega}} \right]. \end{aligned} \quad (6.30)$$

Note that for the coherent states, we can have non-zero work even when $A = 0$. This is a purely quantum effect that comes from the fact that for non energy eigenstate the quantum potential is explicitly time dependent. However, keep in mind that the average work always vanishes when the system is undriven. The work distribution *given* that the initial state is $|\eta\rangle$ is then given by

$$P^{(\eta)}(w) = \int dZ |\langle Z|\eta\rangle|^2 \delta(w - W^{(\eta)}(t, Z)), \quad (6.31)$$

and the distribution associated with the coherent state mixture by

$$P^{(\eta)}(w) = \int d^2\eta p_\alpha(\eta) P^{(\eta)}(w). \quad (6.32)$$

6.4.3 Comparison of work averages

As described above, the work average is independent of the mixture and in this case given by $\langle W \rangle = (At)^2/(2m)$. But higher moments will differ. To illustrate

this, consider the “exponentiated work” average $\langle \exp\{-\beta W\} \rangle$. To show that they are different it suffices to look at the high temperature limit $\beta \ll 1$, for which we find,

$$\langle e^{-\beta W} \rangle_{\{\eta\}} = 1 + \beta \hbar \omega \sin^2\left(\frac{\omega \tau}{2}\right) + O(\beta^2), \quad (6.33)$$

for the coherent state mixture and

$$\langle e^{-\beta W} \rangle_{\{n\}} = 1 - \frac{\beta (A\tau)^2}{2m} \sin^2(\omega \tau) + O(\beta^2), \quad (6.34)$$

for the energy eigenstates mixture.

We see that the Jarzynski equality does not apply in general for the Bohmian work. The reason for this is that each wave function contributes with a different ensemble of trajectories and Hamiltonian function. Moreover, the initial phase-space distribution is not of Boltzmann form, even if the mixtures’ density operator is. For this particular driving, the free energy remains constant and the Jarzynski equality reads $\langle \exp\{-\beta W\} \rangle = 1$. We see that this is correctly recovered in the high-temperature limit $\beta \rightarrow 0$, which is the relevant classical limit here.

6.5 Conclusions

In this Chapter and Publication III we have explored a new approach to quantum work. Based on the Bohmian interpretation and the Hamilton-Jacobi theory we generalised the classical definition of work to any closed quantum process in a straightforward manner. This approach has the advantage of providing well-defined results for any such process, with a proper probability distribution. Furthermore, it respects the conservation of average energy. It also introduces new and unexplored facets of quantum work, such as its possible dependence on the statistical mixture of states, when the mixture is known. This approach can also provide a clear answer to the quantum-classical transition by studying centre-of-mass dynamics [100], the high-temperature limit or large quantum numbers.

7. Energy exchange and the Conditional Wave Function of open quantum systems

In the previous chapter, we have used quantum phase-space trajectories to study work in closed quantum systems, based on the quantum Hamilton-Jacobi equation. In this Chapter, we take a step towards open quantum systems by exploring the concept of the *conditional wave function* from Bohmian mechanics.

7.1 Motivation

In the limit of Markovian evolution and continuous monitoring, stochastic Schrödinger Equations (SSE) can be derived to model the evolution of the state in a single run. However, taking this approach beyond this limit is not entirely straightforward. Under strong coupling or non-Markovian evolution, the system, environment and measuring apparatus can be in a complicated entangled state. In this case, according to the standard interpretation, the system does not have a wave function of its own, *i.e.*, there is no wave function that depends only on the degrees of freedom of the system. Thus, while it is still possible to unravel non-Markovian master equations into SSE [101, 102, 103], what the stochastic state represents becomes unclear [29, 104, 105].

However, in the Bohmian interpretation, the concept of *conditional wave functions* (CWF) provides a unique and conceptually clean way of defining a wave function for any system entangled with other degrees of freedom [27, 37]. To study the fluctuation of energy, we introduce the concept of *conditional energy* as the expectation value of the system Hamiltonian with respect to its CWF. We show that this quantity naturally partitions into three distinct contributions, namely, an *external* contribution due to the explicit time dependence of the system Hamiltonian, an *interaction* contribution due to the presence of an interaction Hamiltonian and an *entanglement* contribution which can be solely attributed to the presence of entanglement. The entanglement contribution provides a direct link between the energy fluctuation of an open system and the entanglement between the system and its environment.

7.2 Conditional Wave Functions

Consider again the case described in Section 6.1 of N point particles with wave function $\Psi(\vec{z}, t)$ and the Hamiltonian in Eq. (6.2). We now take particle $i = 1$ as our system of interest and the remaining $N - 1$ as its environment. Let $\vec{x}$ be the configuration of the system and $\vec{y}$ the configuration of the environment, such that, $\vec{z} = (\vec{x}, \vec{y})$ and $\Psi(\vec{z}, t) = \Psi(\vec{x}, \vec{y}, t)$. Also, let $\vec{Y}(t|Z) := (\vec{Z}_2^{(\Psi)}(t|Z), \dots, \vec{Z}_N^{(\Psi)}(t|Z))$ be the environment trajectory given the initial condition Z , as per Eq. (6.9). In the position representation the CWF $\tilde{\phi}^{(\Psi)}(\vec{x}, t|Z)$ is formally defined as,

$$\tilde{\phi}^{(\Psi)}(\vec{x}, t|Z) := \Psi(\vec{x}, \vec{Y}(t|Z), t) = \langle \vec{x}, \vec{Y}(t|Z) | \Psi(t) \rangle = \langle \vec{x} | \tilde{\phi}^{(\Psi)}(t|Z) \rangle, \quad (7.1)$$

where $|\vec{x}\rangle$ is an element of the system position basis, $|\vec{x}, \vec{Y}(t|Z)\rangle$ an element of the combined system and environment configuration basis and $|\tilde{\phi}^{(\Psi)}(t|Z)\rangle$ is the ket representation of the CWF. The crucial idea behind the CWF is that, given the same initial condition Z , it generates the exact same system trajectory that the wave function $\Psi(\vec{z}, t)$ generates through Eqs. (6.6) and (6.9). If the same trajectories are generated, the same statistical predictions are obtained. Note that the CWF is conditioned on the trajectory of the environment and thus an explicit function of $\vec{x}$ and t only. In this sense, it is a single-particle wave function. A detailed discussion on CWFs can be found in Ref. [27].

While $\Psi(\vec{z}, t)$ is a function of the combined system and environment configuration and evolves according to a Schrödinger equation (SE), the CWF is a function of the system configuration alone and evolves according to a non-linear Schrödinger equation (NSE). If we split the Hamiltonian as $\hat{H}(t) = \hat{H}_S(t) \otimes 1_E + 1_S \otimes \hat{H}_E + \hat{H}_{\text{int}}$, the NSE that governs the evolution of the CWF is given by,

$$\begin{aligned} \partial_t \tilde{\phi}^{(\Psi)}(\vec{x}, t|Z) = & -\frac{i}{\hbar} \left\{ \langle \vec{x} | \hat{H}_S(t) | \tilde{\phi}(t|Z) \rangle + \langle \vec{x}, \vec{Y}_t | \hat{H}_{\text{int}} | \Psi(t) \rangle \right. \\ & \left. + \langle \vec{x}, \vec{Y}_t | 1_S \otimes \hat{H}_E | \Psi(t) \rangle - \vec{v}_{\vec{Y}_t}^{(\Psi)} \cdot \langle \vec{x}, \vec{Y}_t | 1_S \otimes \hat{P}_E | \Psi(t) \rangle \right\}, \end{aligned} \quad (7.2)$$

where we have used the shorthand notation $\vec{Y}_t := \vec{Y}(t|Z)$ and

$$\vec{v}_{\vec{Y}_t}^{(\Psi)} \cdot \langle \vec{x}, \vec{Y}_t | 1_S \otimes \hat{P}_E | \Psi(t) \rangle := -i\hbar \sum_{i=2}^N \vec{v}_i^{(\Psi)}(\vec{Z}(t|Z), t) \cdot (\nabla_i \Psi)(\vec{x}, \vec{Y}_t, t). \quad (7.3)$$

Note that Eq. (7.2) is stochastic in nature despite having no explicit noise term. Every time an experimental run is repeated the system and environment start in a new configuration Z' (as described in the previous Chapter), which leads to a different trajectory of the environment and thus, in general, a different CWF evolution.

For the purpose of calculations and presentation, it is more convenient to express Eq. (7.2) in terms of the unnormalized density operator $\hat{\rho}^{(\Psi)}(t|Z) = |\tilde{\phi}^{(\Psi)}(t|Z)\rangle \langle \tilde{\phi}^{(\Psi)}(t|Z)|$. The evolution of $\hat{\rho}^{(\Psi)}$ is then given by

$$d_t \hat{\rho}^{(\Psi)}(t|Z) = d_t \hat{\rho}_S^{(\Psi)}(t|Z) + d_t \hat{\rho}_{\text{int}}^{(\Psi)}(t|Z) + d_t \hat{\rho}_{\text{ent}}^{(\Psi)}(t|Z), \quad (7.4)$$

where

$$d_t \hat{\rho}_S^{(\Psi)} = -\frac{i}{\hbar} [\hat{H}_S, \hat{\rho}_S^{(\Psi)}], \quad (7.5)$$

$$d_t \hat{\rho}_{\text{int}}^{(\Psi)} = -\frac{i}{\hbar} \langle \vec{Y}_t | [\hat{H}_{\text{int}}, \hat{\sigma}^{(\Psi)}(t)] | \vec{Y}_t \rangle_E, \quad (7.6)$$

$$d_t \hat{\rho}_{\text{ent}}^{(\Psi)} = -\frac{i}{\hbar} \left[\langle \vec{Y}_t | [\hat{H}_E, \hat{\sigma}^{(\Psi)}(t)] | \vec{Y}_t \rangle_E - \vec{v}_{\vec{Y}_t}^{(\Psi)} \cdot \langle \vec{Y}_t | [\hat{P}_E, \hat{\sigma}^{(\Psi)}(t)] | \vec{Y}_t \rangle_E \right], \quad (7.7)$$

and $\hat{\sigma}^{(\Psi)}(t) = |\Psi(t)\rangle\langle\Psi(t)|$. We have used the shorthand notation $\langle \vec{Y}_t | \hat{A} | \vec{Y}_t \rangle_E := \text{Tr}_E \{ \mathbb{1}_S \otimes |\vec{Y}_t\rangle\langle\vec{Y}_t| \hat{A} \}$ for any operator $\hat{A}$ in the combined Hilbert space of the system and environment, and $|\vec{Y}_t\rangle\langle\vec{Y}_t|$ is a projection operator into the subspace spanned by the environment configuration basis element $|\vec{Y}_t\rangle$.

Finally, we point out that the CWF also admits an operational interpretation from weak values [39]. The CWF $\phi^{(\Psi)}(x, t|Z) = \Psi(\vec{x}, \vec{Y}_t, t)$ is the weak value of the projection operator $|\vec{x}\rangle\langle\vec{x}|$, pre-selected on Ψ and post-selected on the state $|\rho_S = 0\rangle \otimes |\vec{y} = \vec{Y}_t\rangle$, where $|\rho_S = 0\rangle$ is a zero-momentum eigenstate of the system (see Section 6.2).

7.3 Conditional energy

To study the energy fluctuations of an open system, one could extend the definition introduced in Chapter 6. This choice would, however, lead to some difficulties. On the one hand, when the system and environment are entangled, the quantum potential becomes a function of both the system and environment degrees of freedom. One would then have to decide whether the quantum potential energy is part of the system energy, the environment energy or both. While this would be more a matter of convention rather than a problem *per se*, it would introduce some ambiguity in the analysis. On the other hand, the energy of the system would be a function of the system configuration. In many experimental scenarios, this configuration is not available or is not relevant for the level of description sought. Thus, an approach at the level of the system CWF can be more easily generalised and applied to a variety of experimental scenarios¹.

We thus take a different route here. A common approach to study energy fluctuations in open systems described by stochastic states is to look at the expectation value of the system Hamiltonian. Following this approach, we define the *conditional energy* $u^{(\Psi)}(t|Z)$ as,

$$u^{(\Psi)}(t|Z) := \frac{\langle \tilde{\phi}^{(\Psi)}(t|Z) | \hat{H}_S(t) | \tilde{\phi}^{(\Psi)}(t|Z) \rangle}{\langle \tilde{\phi}^{(\Psi)}(t|Z) | \tilde{\phi}^{(\Psi)}(t|Z) \rangle} = \frac{\text{Tr} \{ \hat{\rho}^{(\Psi)}(t|Z) \hat{H}_S(t) \}}{\text{Tr} \{ \hat{\rho}^{(\Psi)}(t|Z) \}}. \quad (7.8)$$

To avoid confusion, we'll refer to $\text{Tr} \{ \hat{H}_S(t) \sigma^{(\Psi)}(t) \}$, *i.e.*, the expectation value of $\hat{H}_S(t)$ at the level of $\Psi(t)$, as the *canonical average energy*.

¹ The CWF is still a function of the environment configuration trajectory which might be impossible to track for large systems in practice. However, in some cases, suitable approximations can be found to model the environment trajectory [106, 107].

Using the NSE Eq. (7.4) we show that the conditional energy time derivative, or *conditional energy flow*, can be divided into three distinct terms as,

$$d_t u^{(\Psi)}(t|Z) = d_t u_{\text{ext}}^{(\Psi)}(t|Z) + d_t u_{\text{int}}^{(\Psi)}(t|Z) + d_t u_{\text{ent}}^{(\Psi)}(t|Z), \quad (7.9)$$

where

$$d_t u_{\text{ext}}^{(\Psi)}(t|Z) = \frac{\text{Tr} \{ \hat{\rho}^{(\Psi)}(t|Z) d_t \hat{H}_S(t) \}}{\text{Tr} \{ \hat{\rho}^{(\Psi)}(t|Z) \}}; \quad (7.10)$$

$$d_t u_{\text{int}}^{(\Psi)}(t|Z) = \frac{\text{Tr} \{ (\hat{H}_S(t) - u^{(\Psi)}(t|Z)) d_t \hat{\rho}_{\text{int}}^{(\Psi)}(t|Z) \}}{\text{Tr} \{ \hat{\rho}^{(\Psi)}(t|Z) \}}; \quad (7.11)$$

$$d_t u_{\text{ent}}^{(\Psi)}(t|Z) = \frac{\text{Tr} \{ (\hat{H}_S(t) - u^{(\Psi)}(t|Z)) d_t \hat{\rho}_{\text{ent}}^{(\Psi)}(t|Z) \}}{\text{Tr} \{ \hat{\rho}^{(\Psi)}(t|Z) \}}. \quad (7.12)$$

These three terms are identified as the *external*, *interaction* and *entanglement* contributions to the energy flow based on the following observations. If the system Hamiltonian has no explicit time dependency, usually associated with an external control, $\partial_t \hat{H}_S(t) = 0$ and the external term vanishes. If there is no interaction Hamiltonian between the system and environment at time t , $\hat{H}_{\text{int}} = 0$, $d_t \hat{\rho}_{\text{int}}^{(\Psi)} = 0$ and the interaction contribution vanishes. If there is no entanglement between the system and environment at time t , the entanglement contribution vanishes. To see this, consider a factorized state $|\Psi(t)\rangle = |\phi(t)\rangle \otimes |\chi(t)\rangle$, with $|\phi(t)\rangle$ the system wave function and $|\chi(t)\rangle$ the environmental wave function. Then, $d_t \hat{\rho}_{\text{ent}}^{(\Psi)}(t|Z) = f(t|Z) |\phi(t)\rangle \langle \phi(t)|$ with $f(t|Z)$ a complex-valued function that depends only on time and the initial conditions Z . Noting that the conditional wave function reduces to $|\tilde{\phi}^{(\phi \otimes \chi)}(t|Z)\rangle = \langle \tilde{Y}_t | \chi(t) \rangle |\phi(t)\rangle$, we see that $\text{Tr} \{ \hat{\rho}^{(\phi \otimes \chi)}(t|Z) \} = |\langle \tilde{Y}_t | \chi(t) \rangle|^2$ and the conditional energy reduces to $u^{(\phi \otimes \chi)}(t|Z) = |\langle \tilde{Y}_t | \chi(t) \rangle|^2 \langle \phi(t) | \hat{H}_S(t) | \phi(t) \rangle / |\langle \tilde{Y}_t | \chi(t) \rangle|^2 = \langle \phi(t) | \hat{H}_S(t) | \phi(t) \rangle$. Thus, it follows from Eq. (7.12) that the entanglement contribution vanishes as,

$$d_t u_{\text{ent}}^{(\phi \otimes \chi)}(t|Z) = \frac{f(t|Z)}{|\langle \tilde{Y}_t | \chi(t) \rangle|^2} (\langle \phi(t) | \hat{H}_S(t) | \phi(t) \rangle - u^{(\phi \otimes \chi)}(t|Z)) = 0. \quad (7.13)$$

Note that each of these terms can be non-zero while the others are zero. The entanglement term, in particular, provides a direct link between the energy fluctuations of an open quantum system and its entanglement with the environment.

To link these results with the canonical average energy, we must take the statistical average of the conditional energy over all possible initial conditions $\langle \langle u^{(\Psi)} \rangle \rangle(t) := \int dZ |\Psi(Z, 0)|^2 u^{(\Psi)}(t|Z)$. We find that,

$$\langle \langle u^{(\Psi)} \rangle \rangle(t) = \text{Tr} \{ \hat{H}_S(t) \sigma^{(\Psi)}(t) \}, \quad (7.14)$$

which shows that the average of the conditional energy over many runs of the protocol indeed reproduces the expected canonical average energy. Furthermore,

we can relate the statistical average of the conditional energy flows Eqs. (7.10)-(7.12) to that of the time derivative of the canonical average $d_t \text{Tr} \{ \hat{H}_S(t) \sigma^{(\Psi)}(t) \}$. For the external contribution, we find that

$$\langle \langle d_t u_{\text{ext}}^{(\Psi)} \rangle \rangle(t) = \text{Tr} \{ \hat{\sigma}^{(\Psi)}(t) d_t \hat{H}_S(t) \}, \quad (7.15)$$

which confirms that the statistical average of the external contribution is just the expectation value of the external work. From Eq. (7.14) it now follows that

$$\langle \langle d_t u_{\text{int}}^{(\Psi)} \rangle \rangle(t) + \langle \langle d_t u_{\text{ext}}^{(\Psi)} \rangle \rangle(t) = \text{Tr} \{ \hat{H}_S(t) d_t \hat{\sigma}^{(\Psi)}(t) \}, \quad (7.16)$$

and, in particular, we can show that,

$$\langle \langle d_t u_{\text{ent}}^{(\Psi)} \rangle \rangle(t) = -\frac{i}{\hbar} \text{Tr} \{ \hat{\sigma}^{(\Psi)}(t) [\hat{u}^{(\Psi)}(t), \hat{H}_{\text{int}}] \}, \quad (7.17)$$

where $\hat{u}^{(\Psi)}(t)$ is a state-dependent operator whose elements in an arbitrary basis are given by,

$$\langle i | \hat{u}^{(\Psi)}(t) | j \rangle = \int d\mathbf{y} \langle i | \vec{y} \rangle \langle \vec{y} | j \rangle u^{(\Psi)}(t, \vec{y}), \quad (7.18)$$

where $|i\rangle, |j\rangle$ are vectors from the basis set, and $u^{(\Psi)}(t, \vec{y})$ is the conditional energy of the system at time t given that the environment is in a configuration $\vec{y}$ at the same time². Equation (7.17) shows that the entanglement contribution makes a contribution to the canonical average energy flow only when an interaction is present. This is in line with the notion that, on average, no energy flow can come about from entanglement alone. In general, even in the presence of an interaction Hamiltonian, Eq. (7.17) can be zero if, for example, $[\hat{u}^{(\Psi)}(t), \hat{H}_{\text{int}}] = 0$. A few relevant examples can be found in Publication IV. Here, we present a case of “driving with entanglement” or *steering*.

7.4 Driving with entanglement

Consider two 1/2-spin particles, S and E , prepared in an entangled state,

$$\Psi(x, y, 0) = \frac{1}{\sqrt{2}} \begin{pmatrix} g(y)e^{ikx}f(x) \\ g(y)f(x) \\ 0 \\ 0 \end{pmatrix}, \quad (7.19)$$

where x is the position of particle S , y is the position of particle E , $f(x) = \sqrt[4]{1/(2\pi)} \exp\{-x^2/4\}$, $g(y) = \sqrt[4]{1/(2\pi)} \exp\{-y^2/4\}$, k is a wave number and the spin components are represented in the basis $\{ |\uparrow_E\rangle |\uparrow_S\rangle, |\uparrow_E\rangle |\downarrow_S\rangle, |\downarrow_E\rangle |\uparrow_S\rangle, |\downarrow_E\rangle |\downarrow_S\rangle \}$. We work in units of $\hbar = 1$ and assume both particles have mass one. An unitary

²Note that, like the CWF, the value of the conditional energy at time t depends only on the environment configuration at the same time. The initial conditions are needed only to determine its time evolution.

operation $\hat{U} = \exp(-ivt\hat{P}_E|\uparrow_E\rangle\langle\uparrow_E|\otimes 1_S)$ acting *only* on particle E is then applied, where v is a real-valued constant, t is the duration of the unitary and $\hat{P}_E$ is the momentum operator of particle E . At the end of this operation, the new state is,

$$\Psi(x, y, t) = \frac{1}{\sqrt{2}} \begin{pmatrix} g(y - vt)e^{ikx}f(x) \\ g(y)f(x) \\ 0 \\ 0 \end{pmatrix}. \quad (7.20)$$

For large enough vt , this operation splits the initial wave function into two wave packets with disjoint y -support, one centered at $(x, y) = (0, 0)$ and another one at $(x, y) = (0, vt)$. We take particle S as our system of interest and look at its kinetic energy by setting $\hat{H}_S = P_S^2/2$. Since there is no explicit time dependence on $\hat{H}_S$ and no interaction term, the only contribution comes from the entanglement Eq. (7.12).

The velocity field of particle E can be determined from the continuity equation in configuration space [108]. For particles with spin, the probability distribution in configuration space is given by taking the inner product in spin-space [99], that is,

$$\mathbb{P}(x, y, t) = \sum_{s, s'} \Psi_{s, s'}^*(x, y, t) \Psi_{s, s'}(x, y, t) = g^2(y + vt)f^2(x)/2 + g^2(y)f^2(x)/2, \quad (7.21)$$

where $\Psi_{s, s'}(x, y, t)$ is the $|s_S\rangle|s'_E\rangle$ component of $\Psi(x, y, t)$. Taking the partial time derivative we find the continuity equation,

$$\partial_t \mathbb{P}(x, y, t) + \partial_y [v_E^{(\Psi)}(y, t) \mathbb{P}(x, y, t)] = 0, \quad (7.22)$$

where $v_E^{(\Psi)}(y, t) = vg^2(y + vt)/[g^2(y + vt) + g^2(y)]$. With the assumption that the velocity vanishes at infinity, the velocity field of particle E is given by $v_E^{(\Psi)}(y, t)$. For the conditional energy we find the simple expression,

$$u^{(\Psi)}(t|Z) = \frac{1}{2} \left[\frac{1}{4} + k^2 \frac{g^2(Y_t - vt)}{g^2(Y_t - vt) + g^2(Y_t)} \right]. \quad (7.23)$$

The trajectories $Y(t|Z)$ and the respective conditional energy $u^{(\Psi)}(t|Z)$ are plotted in Fig. 7.1.

The results can be easily understood on physical grounds. Initially, the energy is independent of y and simply the average of the energy of a Gaussian wave packet with zero group velocity, $E^{(0)} = 1/8$ and that of a Gaussian wave packet with group velocity k , $E^{(k)} = E^{(0)} + k^2/2$. As time evolves and the wave packets begin to separate, the energy changes depending on the exact trajectory Y_t . Once the wave packets are well separated, only one term in Eq. (7.20) is relevant and the wave function behaves as if *effectively factorized* [27]. As a consequence, the energy converges either to $E^{(0)}$ or $E^{(k)}$.

Note that because there is no interaction, the canonical average energy remains constant. However, in each run of the experiment, the system state can be

Figure 7.1. (a) Particle trajectories Y_t (normalized with the standard deviation) for evenly separated initial positions for the case of two entangled spin 1/2 particles. (b) The conditional energy change corresponding to the trajectories in (a) normalized with the splitting parameter $\Delta = k^2/2$.

“collapsed” to either one of the two wave packets by measuring first the position of particle E. Thus, for each run of the experiment, the particle energy can increase or decrease. These fluctuations can be directly attributed to entanglement and inferred experimentally via weak measurements. A similar experiment was reported in Ref. [109], where a photon is remotely steered via entanglement and the trajectories reconstructed using weak measurement.

7.5 Conclusions

In this Chapter, we have presented a general way of describing energy fluctuations at the level of individual runs of experiments based on the Bohmian CWF. The main advantage of using the CWF is that it provides a conceptually clean way of assigning a wave function to an open system. From single runs to the ensemble average over many runs, different experimental scenarios can be analysed and understood on general grounds.

The CWF reveals three distinct contributions to the energy fluctuations of the open quantum system. They are, 1) an external contribution coming from any explicit time dependence of the system bare Hamiltonian, also typically identified as the external work [7], 2) an interaction contribution coming from the presence

of an interaction Hamiltonian with its environment and 3) an entanglement contribution, coming from the presence of entanglement between the open system and its environment. We highlight the entanglement contribution since it provides a direct link between energy fluctuations and entanglement without evoking any information-theoretic arguments.

The close connection to weak values provides an operational interpretation of the CWF and puts it within reach of experimental verification. Aside from its theoretical interest, it can offer new ways to experimentally quantify the role of entanglement in energy fluctuations and identify genuine quantum advantages in novel devices such as quantum heat engines.

8. Summary and Conclusions

The work presented in this thesis gives a small contribution to the understanding of energy fluctuations in quantum systems. The thesis focuses primarily on how to address energy fluctuations along individual runs of an experiment, an essential step to fully import the successful picture of classical stochastic thermodynamics to the quantum regime. This work was conducted in two parts.

In the first part, the experimental calorimetric detection protocol was analysed for a quantum harmonic oscillator. This protocol is designed to work at low temperatures where single quanta of energy can be observed. Since only the lowest levels of the system will be initially occupied, a simple “two-level approximation” (TLA) was proposed and its validity compared to the results of the two-point-measurement (TPM) protocol. Different ratios between the driving and dissipation strength were analysed. In the limit of unitary driving, where dissipation is negligible, analytic expressions are reported that allow recovering the two-point measurement result. When the dissipation characteristic time is of the same order as the driving time, the results for the unitary regime were expanded based on second-order time-dependent perturbation theory. The results are in good agreement with numerical simulation and show that the error between the TLA and TPM decreases. Finally, numerical simulations were carried out to study the case where the dissipation dominates. It was found that the average work is in excellent agreement with the TPM result, even for more extended driving periods.

Analysis of the calorimetric setup was carried out further by exploring possible non-Markovian effects due to the finite size of the environment. Using a previously developed finite-environment master equation (FEME), the induced non-Markovianity was quantified using the Breuer-Laine-Piilo (BLP) measure. In particular, the dependence of the BLP measure on the environment size, system driving strength and coupling to the environment were studied. A non-trivial non-Markovian structure in the parameter-space was revealed. A remarkable feature is that non-Markovianity does not decrease monotonically with the environment size. By adjusting the driving strength and coupling to the environment, one can induce or mitigate non-Markovian effects.

In the second part of this thesis, a more detailed and general analysis of

energy fluctuations in quantum systems is undertaken. The main difficulty in addressing energy fluctuations in quantum systems is that energy is not always well-defined according to the standard interpretation of quantum mechanics. While the TPM protocol is very versatile, it cannot capture processes with initial energy superpositions. However, in the Bohmian interpretation of quantum mechanics, it is possible to construct a well-defined energy for general quantum processes. Together with the quantum Hamilton-Jacobi equation, this is leveraged to introduce a different approach to quantum energy fluctuation. First, it is shown how the quantum Hamilton-Jacobi equation can be used to provide a Hamiltonian function in phase-space that generates Bohmian trajectories. With this crucial ingredient, any closed quantum process can be described in terms of ensembles of trajectories generated by a Hamiltonian function in phase space. One can then generalise the classical definition of work, as power integrated along the system trajectory, to the quantum case. This approach leads to a proper probability distribution of work that respects the conservation of average energy. It also reveals an interesting new contribution coming from the quantum potential, inducing purely quantum fluctuations. Another distinct feature arises when explicitly prepared statistical mixtures of pure states are considered. In that case, the work distribution is, in general, dependent on the mixture itself, even when they correspond to the same density operator. These different distributions can be reconstructed from weak measurements and provide a new perspective into the fluctuations of energy in closed quantum systems.

To carry the analysis to open systems, the Bohmian conditional wave function was employed to define the conditional energy of the open system. It was shown that there are three distinct contributions to the fluctuations of the conditional energy: 1) An external contribution coming from the explicit time dependence of the system bare Hamiltonian, typically associated with an external control parameter; 2) an interaction contribution due to the presence of an interaction Hamiltonian and; 3) an entanglement contribution due to the presence of entanglement between the system and environment. The entanglement contribution is perhaps the most interesting as it provides a direct link between energy fluctuations and entanglement. Such a connection is not readily available in other stochastic approaches based on the standard interpretation.

The work developed here opens a new way to understand fluctuations of energy in open quantum systems through an observer-independent lens. The close connection between Bohmian trajectories and weak measurements makes these quantities experimentally accessible, offering a new form of experimental characterisation. In recent years, Bohmian trajectories have been applied to tackle different problems, from transport in realistic nanoscale electronic systems [96, 107, 110] and chemical reactions [92] to the study of the foundations of quantum mechanics [109, 111, 112, 113, 114]. Developing a general framework of stochastic quantum thermodynamic based on CWF is undoubtedly a future possibility. What we have identified as the external contribution is already related to what is commonly known as the external work. In the limit

of weak interaction, continuous monitoring of the environment and Markovian evolution, we expect the entanglement contribution to be negligible. Thus, heat can be identified with the interaction contribution. For an entangled system and environment, an exciting prospect is to determine some form of “entanglement heat”, that is, the part of the energy exchanged that can be attributed directly to entanglement. Future work is thus likely to be linked with developing these ideas further, either through new numerical implementations or by inspiring new experiments to test the limits of our current understanding of quantum mechanics [115, 116, 117].

References

- [1] S. Suomela, R. Sampaio, and T. Ala-Nissila, *Phys. Rev. E* **94**, 032138 (2016).
- [2] E. Bäumer, M. Lostaglio, M. Perarnau-Llobet, and R. Sampaio (2018), *arXiv:1805.10096*.
- [3] M. K. E. L. Planck, *Verhandl. Dtsc. Phys. Ges.* **2**, 237 (1900).
- [4] M. K. E. L. Planck, *Verhandl. Dtsc. Phys. Ges.* **2**, 202 (1900).
- [5] A. Einstein, *Annalen der Physik* **322**, 132 (1905).
- [6] A. B. Arons and M. B. Peppard, *Am. J. Phys.* **33**, 367 (1965).
- [7] S. Vinjanampathy and J. Anders, *Contemporary Phys.* **57**, 545 (2016).
- [8] J. P. Pekola, *Nat. Phys.* **11**, 118 (2015).
- [9] G. Wendin, *Rep. Prog. Phys.* **80**, 106001 (2017).
- [10] M. Campisi, J. Pekola, and R. Fazio, *New J. Phys.* **17**, 035012 (2015).
- [11] R. Kosloff and A. Levy, *An. Rev. Phys. Chem.* **65**, 365 (2014).
- [12] W. Niedenzu, V. Mukherjee, A. Ghosh, A. G. Kofman, and G. Kurizki, *Nat. Comm.* **9**, 165 (2018).
- [13] A. Ronzani, B. Karimi, J. Senior, Y.-C. Chang, J. T. Peltonen, C. Chen, and J. P. Pekola, *Nat. Phys.* **14**, 991 (2018).
- [14] J. Klaers, S. Faelt, A. Imamoglu, and E. Togan, *Phys. Rev. X* **7**, 031044 (2017).
- [15] G. Manzano, F. Galve, R. Zambrini, and J. M. R. Parrondo, *Phys. Rev. E* **93**, 052120 (2016).
- [16] L. E. Reichl, *A modern course in statistical Phys.* (John Wiley & Sons, 2016).
- [17] G. Bochkov and Y. E. Kuzovlev, *Zh. Eksp. Teor. Fiz* **72**, 238 (1977).
- [18] C. Jarzynski, *Phys. Rev. Lett.* **78**, 2690 (1997).
- [19] U. Seifert, *Phys. Rev. Lett.* **95**, 040602 (2005).
- [20] G. E. Crooks, *Phys. Rev. E* **60**, 2721 (1999).
- [21] U. Seifert, *Rep. Prog. Phys.* **75**, 126001 (2012).
- [22] P. Talkner and P. Hänggi, *Phys. Rev. E* **94**, 022143 (2016).

- [23] H. M. Wiseman and G. J. Milburn, *Quantum measurement and control* (Cambridge university press, 2009).
- [24] F. Binder, L. A. Correa, C. Gogolin, J. Anders, and G. A. (Eds.), *Thermodynamics in the Quantum Regime: Fundamental Aspects and New Directions* (Springer International Publishing, 2018).
- [25] D. Bohm, Phys. Rev. **85**, 166 (1952).
- [26] D. Bohm, Phys. Rev. **85**, 180 (1952).
- [27] S. Teufel and D. Dürr, *Bohmian Mechanics* (Springer, 2009).
- [28] P. R. Holland, *The quantum theory of motion: an account of the de Broglie-Bohm causal interpretation of quantum mechanics* (Cambridge university press, 1995).
- [29] B. d'Espagnat, *Conceptual foundations of quantum mechanics* (CRC Press, 2018).
- [30] W. H. Zurek, Nature **341**, 119 (1989).
- [31] P. Kammerlander and J. Anders, Sci. Rep. **6**, 22174 (2016).
- [32] J. Goold, M. Huber, A. Riera, L. del Rio, and P. Skrzypczyk, J. Phys. A **49**, 143001 (2016).
- [33] S. Deffner, J. P. Paz, and W. H. Zurek, Phys. Rev. E **94**, 010103 (2016).
- [34] J. M. Parrondo, J. M. Horowitz, and T. Sagawa, Nat. Phys. **11**, 131 (2015).
- [35] T. Sagawa and M. Ueda, Phys. Rev. Lett. **102**, 250602 (2009).
- [36] H.-P. Breuer and F. Petruccione, *The theory of open quantum systems* (Oxford University Press on Demand, 2002).
- [37] D. Dürr, S. Goldstein, and N. Zanghi, J. Stat. Phys. **67**, 843 (1992).
- [38] Y. Aharonov, D. Z. Albert, and L. Vaidman, Phys. Rev. Lett. **60**, 1351 (1988).
- [39] T. Norsen and W. Struyve, A. Phys. **350**, 166 (2014).
- [40] G. Lindblad, Comm. Math. Phys. **48**, 119 (1976).
- [41] V. Gorini, A. Kossakowski, and E. C. G. Sudarshan, J. Math. Phys. **17**, 821 (1976).
- [42] K. Mølmer, Y. Castin, and J. Dalibard, J. Opt. Soc. Am. B **10**, 524 (1993).
- [43] R. Schmidt, M. F. Carusela, J. P. Pekola, S. Suomela, and J. Ankerhold, Phys. Rev. B **91**, 224303 (2015).
- [44] R. Schmidt, S. Maniscalco, and T. Ala-Nissila, Phys. Rev. A **94**, 010101 (2016).
- [45] G. Guarnieri, J. Nokkala, R. Schmidt, S. Maniscalco, and B. Vacchini, Phys. Rev. A **94**, 062101 (2016).
- [46] Sahar Alipour, *et al.*, in preparation (2018).
- [47] G. Lindblad, Comm. Math. Phys. **65**, 281 (1979).
- [48] B. Bylicka, D. Chruściński, and S. Maniscalco, Sci. Rep. **4**, 5720 (2014).
- [49] F. Verstraete, M. M. Wolf, and J. I. Cirac, Nat. Phys. **5**, 633 (2009).
- [50] J. F. Poyatos, J. I. Cirac, and P. Zoller, Phys. Rev. Lett. **77**, 4728 (1996).
- [51] D. Kienzler, H.-Y. Lo, B. Keitch, L. de Clercq, F. Leupold, F. Lindenefelser, M. Marinelli, V. Negnevitsky, and J. P. Home, Science **347**, 53 (2015).

- [52] S. G. Schirmer and X. Wang, Phys. Rev. A **81**, 062306 (2010).
- [53] S. Lorenzo, F. Plastina, and M. Paternostro, Phys. Rev. A **88**, 020102 (2013).
- [54] S. Luo, S. Fu, and H. Song, Phys. Rev. A **86**, 044101 (2012).
- [55] X.-M. Lu, X. Wang, and C. Sun, Phys. Rev. A **82**, 042103 (2010).
- [56] M. M. Wolf, J. Eisert, T. Cubitt, and J. I. Cirac, Phys. Rev. Lett. **101**, 150402 (2008).
- [57] R. Vasile, S. Maniscalco, M. G. Paris, H.-P. Breuer, and J. Piilo, Phys. Rev. A **84**, 052118 (2011).
- [58] H.-P. Breuer, E.-M. Laine, and J. Piilo, Phys. Rev. Lett. **103**, 210401 (2009).
- [59] A. Rivas, S. F. Huelga, and M. B. Plenio, Phys. Rev. Lett. **105**, 050403 (2010).
- [60] S. Suomela, A. Kutvonen, and T. Ala-Nissila, Phys. Rev. E **93**, 062106 (2016).
- [61] S. Wißmann, A. Karlsson, E.-M. Laine, J. Piilo, and H.-P. Breuer, Phys. Rev. A **86**, 062108 (2012).
- [62] H. Tasaki (2000), arXiv:cond-mat/0009244.
- [63] J. Kurchan (2000), arXiv:cond-mat/0007360.
- [64] P. Talkner, E. Lutz, and P. Hänggi, Phys. Rev. E **75**, 050102 (2007).
- [65] P. Talkner and P. Hänggi, J. Phys. A **40**, F569 (2007).
- [66] P. Talkner, P. S. Burada, and P. Hänggi, Phys. Rev. E **78** (2008).
- [67] M. Campisi, P. Hänggi, and P. Talkner, Rev. Mod. Phys. **83**, 771 (2011).
- [68] L. Fusco, S. Pigeon, T. J. G. Apollaro, A. Xuereb, L. Mazzola, M. Campisi, A. Ferraro, M. Paternostro, and G. De Chiara, Phys. Rev. X **4**, 031029 (2014).
- [69] S. Deffner and A. Saxena, Phys. Rev. E **92**, 032137 (2015).
- [70] M. Campisi, P. Talkner, and P. Hänggi, Phys. Rev. Lett. **102**, 210401 (2009).
- [71] S. Suomela, P. Solinas, J. P. Pekola, J. Ankerhold, and T. Ala-Nissila, Phys. Rev. B **90**, 094304 (2014).
- [72] S. Suomela, J. Salmilehto, I. G. Savenko, T. Ala-Nissila, and M. Möttönen, Phys. Rev. E **91**, 22126 (2015).
- [73] M. Perarnau-Llobet, K. V. Hovhannisyanyan, M. Huber, P. Skrzypczyk, N. Brunner, and A. Acín, Phys. Rev. X **5**, 041011 (2015).
- [74] K. Funo, Y. Watanabe, and M. Ueda, Phys. Rev. A **88**, 052319 (2013).
- [75] A. Misra, U. Singh, S. Bhattacharya, and A. K. Pati, Phys. Rev. A **93**, 052335 (2016).
- [76] K. Micadei, J. P. Peterson, A. M. Souza, R. S. Sarthour, I. S. Oliveira, G. T. Landi, T. B. Batalhão, R. M. Serra, and E. Lutz (2017), arXiv:1711.03323.
- [77] P. Solinas and S. Gasparinetti, Phys. Rev. E **92**, 042150 (2015).
- [78] C. Elouard, D. A. Herrera-Martí, M. Clusel, and A. Auffèves, NPJ Q. Info. **3**, 9 (2017).
- [79] G. Bochkov and Y. E. Kuzovlev, Zh. Eksp. Teor. Fiz **72**, 238 (1977).

- [80] G. Bochkov and Y. E. Kuzovlev, *Physica A* **106**, 443 (1981).
- [81] S. Yukawa, *J. Phys. Soc. Jpn.* **69**, 2367 (2000).
- [82] A. E. Allahverdyan and T. M. Nieuwenhuizen, *Phys. Rev. E* **71**, 066102 (2005).
- [83] P. Solinas and S. Gasparinetti, *Phys. Rev. A* **94**, 052103 (2016).
- [84] H. J. D. Miller and J. Anders, *New J. Phys.* **19**, 062001 (2017).
- [85] A. E. Allahverdyan, *Phys. Rev. E* **90**, 32137 (2014).
- [86] M. Lostaglio, *Phys. Rev. Lett.* **120**, 040602 (2018).
- [87] M. Perarnau-Llobet, E. Bäumer, K. V. Hovhannisyan, M. Huber, and A. Acin, *Phys. Rev. Lett.* **118**, 070601 (2017).
- [88] J. S. Bell, On the impossible pilot wave, in *Speakable and Unspeakable in Quantum Mechanics* (Cambridge University Press, 1987).
- [89] D. Dürr, S. Goldstein, and N. Zanghì, *Quantum Phys. without quantum philosophy* (Springer Science & Business Media, 2012).
- [90] De Broglie, Louis, *Ann. Phys.* **10**, 22 (1925).
- [91] de Broglie, Louis, *J. Phys. Radium* **8**, 225 (1927).
- [92] Benseny Albert, Albareda Guillermo, Sanz Ángel S., Mompart Jordi, and Oriols Xavier, *Europhys. J. D* **68**, 286 (2014).
- [93] A. B. Nassar and S. Miret-Artés, *Bohmian mechanics, open quantum systems and continuous measurements* (Springer, 2017).
- [94] W. Struyve, *Scientific Reports* **7**, 8161 (2017).
- [95] S. V. Antipov, S. Bhattacharyya, K. El Hage, Z.-H. Xu, M. Meuwly, U. Rothlisberger, and J. Vaníček, *Structural Dynamics* **4**, 061509 (2017).
- [96] Z. Zhan, E. Colomés, D. Pandey, S. Yuan, and X. Oriols (2018), arXiv:1805.04672.
- [97] D. Dürr, S. Goldstein, R. Tumulka, and N. Zanghì, *Found. Phys.* **35**, 449 (2005).
- [98] D. Dürr, S. Goldstein, and N. Zanghì, *J. Stat. Phys.* **116**, 959 (2004).
- [99] P. R. Holland, *The quantum theory of motion: an account of the de Broglie-Bohm causal interpretation of quantum mechanics* (Cambridge university press, 1995).
- [100] X. Oriols and A. Benseny, *New J. Phys.* **19**, 063031 (2017).
- [101] J. Piilo, S. Maniscalco, K. Härkönen, and K.-A. Suominen, *Phys. Rev. Lett.* **100**, 180402 (2008).
- [102] V. Semin, I. Semina, and F. Petruccione, *Phys. Rev. E* **96**, 063313 (2017).
- [103] W. T. Strunz, L. Diósi, and N. Gisin, *Phys. Rev. Lett.* **82**, 1801 (1999).
- [104] J. Gambetta and H. M. Wiseman, *Phys. Rev. A* **66**, 012108 (2002).
- [105] J. Gambetta and H. M. Wiseman, *Phys. Rev. A* **68**, 062104 (2003).
- [106] X. Oriols, *Phys. Rev. Lett.* **98**, 066803 (2007).
- [107] E. Colomés, Z. Zhan, D. Marian, and X. Oriols, *Phys. Rev. B* **96**, 075135 (2017).
- [108] W. Struyve and A. Valentini, *J. Phys. A* **42**, 035301 (2009).

- [109] Y. Xiao, Y. Kedem, J.-S. Xu, C.-F. Li, and G.-C. Guo, *Opt. Express* **25**, 14463 (2017).
- [110] A. Abedi and M. J. Sharifi, *J. Comput. Electron.* **17**, 68 (2018).
- [111] D. H. Mahler, L. Rozema, K. Fisher, L. Vermeyden, K. J. Resch, H. M. Wiseman, and A. Steinberg, *Sci. Adv.* **2**, e1501466 (2016).
- [112] S. Kocsis, B. Braverman, S. Ravets, M. J. Stevens, R. P. Mirin, L. K. Shalm, and A. M. Steinberg, *Science* **332**, 1170 (2011).
- [113] B. Hiley and P. Van Reeth, *Entropy* **20**, 353 (2018).
- [114] R. Flack and B. Hiley, *Entropy* **20**, 367 (2018).
- [115] N. Gisin, *Entropy* **20**, 105 (2018).
- [116] A. Cabello, M. Gu, O. Gühne, J.-A. Larsson, and K. Wiesner, *Phys. Rev. A* **94**, 052127 (2016).
- [117] C. G. T. Carina E. A. Prunkl, *S. Hist. Phil. Sci. B* **64**, 114 (2017).

ISBN 978-952-60-8414-5 (printed)
ISBN 978-952-60-8415-2 (pdf)
ISSN 1799-4934 (printed)
ISSN 1799-4942 (pdf)

Aalto University
School of Science
Department of Applied Physics
www.aalto.fi

**BUSINESS +
ECONOMY**

**ART +
DESIGN +
ARCHITECTURE**

**SCIENCE +
TECHNOLOGY**

CROSSOVER

**DOCTORAL
DISSERTATIONS**