The water we eat

Methods for estimating
water use of diets in
changing food systems

Mika Jalava

The water we eat

Methods for estimating
water use of diets in
changing food systems

Mika Jalava

A doctoral dissertation completed for the degree of Doctor of Science (Technology) to be defended, with the permission of the Aalto University School of Engineering, at a public examination held at the lecture hall TU1 of the school on 30 August 2019 at 12 noon.

**Aalto University
School of Engineering
Department of Built Environment
Water and Development Research Group**

Supervising professor

Professor Olli Varis

Thesis advisors

Dr. Joseph Guillaume, Australian National University, Australia

Assistant Professor Matti Kummu, Aalto University, Finland

Preliminary examiners

Olçay Ünver, Dr., Deputy Director, FAO Rome, Italy

Professor Jan Lundqvist, Stockholm International Water Institute, Sweden

Opponent

Professor Giuliano Di Baldassarre, Uppsala University, Sweden

Aalto University publication series

DOCTORAL DISSERTATIONS 145/2019

© 2019 Mika Jalava

ISBN 978-952-60-8672-9 (printed)

ISBN 978-952-60-8673-6 (pdf)

ISSN 1799-4934 (printed)

ISSN 1799-4942 (pdf)

<http://urn.fi/URN:ISBN:978-952-60-8673-6>

Images: Cover by Elina Lehtikainen (2019): Spherical Cow

Painosalama Oy

Turku 2019

Finland

Author

Mika Jalava

Name of the doctoral dissertation

The water we eat: Methods for estimating water use of diets in changing food systems

Publisher School of Engineering**Unit** Department of Built Environment**Series** Aalto University publication series DOCTORAL DISSERTATIONS 145/2019**Field of research** Water and Environmental Engineering**Manuscript submitted** 18 April 2019**Date of the defence** 30 August 2019**Permission for public defence granted (date)** 17 June 2019**Language** English☐ **Monograph**☒ **Article dissertation**☐ **Essay dissertation****Abstract**

Water scarcity already affects more than half of the human population. The largest water consuming human activity, food production, is at the centre of the quest for sustainable use of this natural capital. How much resources are required to satisfy the global food demand depends on more than just the number of us and how much we eat. It is largely determined by what the human diets consist of and where the food is produced, as well as how much of it is lost and wasted.

This dissertation focuses on water consumption in agriculture as a key determining factor in achieving food security. It does so by studying three overarching themes: the resource-saving effect of changes in human diets, modelling the supply chains within the complex food systems, and methods used to evaluate agricultural water use and its impacts. A substantial body of existing research aims to find relationships between consumer choices and resource use in agricultural production, but many aspects in the methodologies and in their results remain controversial. This dissertation points out challenges in using current methods to define alternative diets in global studies and presents a novel, optimisation-based method that adjusts diets according to chosen dietary guidelines and level of diet change. It combines the diet change analysis with food system models in a way that can capture interactions between different resource-saving measures and presents estimates of their global, regional and country-level potential to improve food security. A novel food system model, Aalto OptoFood was developed to allow analysing resource efficiency of large changes in food demand and in system dynamics, such as synergies between crop and livestock-based food production.

The dissertation found that replacing livestock-based foodstuffs in human diets with crop-based alternatives is an efficient way to reduce agricultural water consumption. Theoretically, the water thus saved could supply food for up to 1.8 billion people. Yet, the opportunity cost of livestock in terms of water may be lower in practice. Shifting water from the production of one foodstuff to another requires the availability of other resources, such as suitable cropland. Non-arable rangelands may support livestock production that positively contributes to food security. Nevertheless, many current production systems depend heavily on water and land resources that could be more efficiently used for crop production. Optimal spatial distribution of production changes would allow diet change to preferentially reduce livestock with highest opportunity cost in terms of water, and offer high initial returns with relatively small changes.

Finally, this dissertation uses the framework of the Cultural Theory to discuss how modelling outcomes are affected by decisions stemming from different worldviews. It argues that modellers should be conscious of such decisions and communicate them clearly to avoid real or perceived biases.

Keywords water scarcity, food system, diet change, food loss, cropland, livestock**ISBN (printed)** 978-952-60-8672-9**ISBN (pdf)** 978-952-60-8673-6**ISSN (printed)** 1799-4934**ISSN (pdf)** 1799-4942**Location of publisher** Helsinki**Location of printing** Turku**Year** 2019**Pages** 144**urn** <http://urn.fi/URN:ISBN:978-952-60-8673-6>

Tekijä

Mika Jalava

Väitöskirjan nimi

Syömämme vesi: ruokavalioiden vedenkulutuksen arvioiminen muuttuvissa ruokajärjestelmissä

Julkaisija Insinööritieteiden korkeakoulu**Yksikkö** Rakennetun ympäristön laitos**Sarja** Aalto University publication series DOCTORAL DISSERTATIONS 145/2019**Tutkimusala** Vesi- ja ympäristötekniikka**Käsikirjoituksen pvm** 18.04.2019**Väitöspäivä** 30.08.2019**Väittelyluvan myöntämispäivä** 17.06.2019**Kieli** Englanti☐ **Monografia**☒ **Artikkeliväitöskirja**☐ **Esseeväitöskirja****Tiivistelmä**

Jo yli puolet ihmiskunnasta kärsii veden puutteesta. Koska ruuan tuotantoon kuluu suurin osa ihmisen käyttämästä vedestä, se on keskeinen tekijä pyrittäessä vesivarojen kestäväään käyttöön. Globaalin ruuan kysynnän tyydyttämiseen tarvittavan veden määrä riippuu paitsi siitä, montako meitä on ja paljonko syömmme, myös ruokavaliostamme, ruokahävikistä ja ruuan tuotantoalueista.

Tämä väitöskirja keskittyy maatalouden vedenkulutukseen ruokaturvan avaintekijänä. Sen kolme pääteemaa ovat ihmisten ruokavalion muutoksien resurssveja säästävä vaikutus, tarjontaketjujen mallintaminen monimutkaisissa ruokajärjestelmissä ja maatalouden vedenkulutuksen arviointimenetelmät. Kuluttajien valintojen ja maatalouden resurssien käytön välisiä suhteita on tutkittu laajalti, mutta menetelmistä ja tuloksista on yhä ristiriitaa. Tämä väitöskirja osoittaa haasteita nykyisissä menetelmissä, joilla pyritään nykyistä resurssitehokkaampiin ruokavalioiden globaaleissa tutkimuksissa. Se esittää uuden, optimointiin perustuvan menetelmän, jolla ruokavaliota voidaan säätää suositusten ja haluttujen skenaarioiden mukaisiksi. Väitöskirja yhdistää ruokavaliomuutoksen analyysin ruokajärjestelmämalleihin mahdollistaen veden säästöön tähtäävien menetelmien yhteisvaikutuksien selvittämisen ja esittää arvioita ruokaturvan edistämisen mahdollisuuksista globaalisti, alueellisesti ja maakohtaisesti. Ruuan kysynnän laajojen muutoksien ja ruokajärjestelmän dynaamisten osien kuten kasvi- ja eläinperäisen ruuan tuottamiseen liittyvien synergioiden analysoimiseksi kehitettiin uusi ruokajärjestelmämalli, Aalto OptoFood.

Väitöskirja osoitti, että eläinperäisten ruoka-aineiden korvaaminen kasviperäisillä on tehokas tapa pienentää maanviljelyn vedenkulutusta. Teoreettisesti siten säästynyt vesi mahdollistaisi ruuan tarjonnan kasvattamisen 1,8 miljardin ihmisen tarpeita vastaavalla määrällä. Eläintuotannon vaihtoehtokustannus voi kuitenkin käytännössä olla pienempi. Tuotannon siirtyminen ruoka-aineesta toiseen edellyttää myös muita resurssveja, kuten viljelymaata. Muuhun viljelyyn sopimattomat laitumet voivat mahdollistaa ruokaturvaa parantavan eläintuotannon. Monet tuotantojärjestelmät kuitenkin tarvitsevat runsaasti vettä ja maata, joiden käyttö kasvintuotannossa olisi tehokkaampaa. Alueellisesti ihanteellinen tuotannon muutoksien sijoittelu mahdollistaisi veden suhteen vaihtoehtokustannuksiltaan kalleimman eläintuotannon ensisijaisen korvaamisen, jolloin suhteellisen pienillä muutoksilla saataisiin nopea vaikutus veden kulutukseen.

Lopuksi väitöskirja pohtii riskin kulttuuriteorian viitekehyksessä sitä, miten erilaisiin maailmankuviin tukeutuvat päätökset vaikuttavat mallinnuksen tuloksiin. Se esittää, että välttyäkseen tuomasta malleihin ennakoasenteen aiheuttamia todellisia vinoumia tai vaikutelmia niistä, mallintajien tulisi tehdä nämä päätökset tietoisesti ja kommunikoida ne selkeästi.

Avainsanat vesipula, ruokajärjestelmä, ruokavalion muutos, ruokahävikki, viljelymaa, eläintuotanto**ISBN (painettu)** 978-952-60-8672-9**ISBN (pdf)** 978-952-60-8673-6**ISSN (painettu)** 1799-4934**ISSN (pdf)** 1799-4942**Julkaisupaikka** Helsinki**Painopaikka** Turku**Vuosi** 2019**Sivumäärä** 144**urn** http://urn.fi/URN:ISBN:978-952-60-8673-6

Acknowledgements

Having travelled a winding road between careers in business and science, in many ways I have most enjoyed these years at Aalto University and its Water and Development Research Group. For this opportunity, for the guidance, and for all the support, as well as the long and interesting discussions about science, politics and so much else, I want to thank my supervisor, Professor Olli Varis. Together with Professor Varis, my sincere thanks for actually making my work and this dissertation possible belong to Assistant Professor Matti Kumm. Thank you for all the work to secure my funding, for all the instruction, advice and hands-on work with my papers!

I want to give my special thanks to Dr. Joseph Guillaume for his instruction with my papers as well as this dissertation. Joseph, you put more hours into coaching me through so many troubles than could ever be expected from an academic instructor. But it's not just the help to reach a goal – I am even more thankful for your persistence as a teacher. I do remember you saying “didn't we talk about this already?” quite a few times, but you always just went on to explain once again the piece of difficult math or whatever the question was.

I am thankful for the kind words, relevant comments and thoughtful advice of the two pre-examiners of my thesis, Professor Jan Lundqvist and Dr. Olcay Ünver. With their approval, I am now looking forward to my defence and meeting my opponent, Professor Giuliano Di Baldassarre. Thank you for accepting the invitation!

Few academic papers are individual efforts. Certainly, none of mine have been. In addition to those mentioned above, many people have helped substantially with the works included in this dissertation. Thank you, all my co-authors! Dr. Miina Porkka was my closest tutor in the early phases of my doctoral studies. Professor Stefan Siebert helped with both data and writing, but I especially appreciate the discussions with him about the complicated nature of international agricultural statistics. And I worked many months together with Dr. Vera Heck – she ran simulations over and over again, explained vegetation modelling, and spent long hours with my manuscripts.

In our group, postdocs play a very important role, and one in particular. I do not know how I could have finished my thesis without Dr. Maija Taka and her synthesis writing group, as well as all the other help she has provided!

All my other colleagues have put a considerable effort into making work easier and more enjoyable. I really miss the long discussions about politics, society and everything with Dr. Juho Haapala, a long time office roommate. I wish there were

such good people working in politics, not just talking about it! Doctors Suvi Sojamo, Aura Salmivaara, Mirja Kattelus and Timo Räsänen showed us “youngsters” what it takes to graduate. My current fellow Water and Development Research Group colleagues, doctoral students Lauri Ahopelto, Amy Fallon, Matias Heino, Anne Hyvärinen, Marko Kallio, Pekka Kinnunen, Elina Lehtikoinen, Hafsa Munia, and Venla Niva have not only participated in all the educated discussions but also put up with all my dog stories. MSc Vili Virkki will soon be in the doctoral students’ team. Thanks Vili for all the help with teaching, R, and GIS. Also, I want to thank Lic. Tech. Pirjo Rantanen, who kept me awake, or at least made my falling asleep less frequent, with her strong coffee and even stronger chili peppers! She was also a member of our synthesis group run by Maija, the others being Heidi Salo and Dr. Pia Väitalo – thank you all for all the insight and support in our peer survival group!

I also want to thank a few colleagues from way back when I started my master’s studies: Professors Harri Koivusalo and Riku Vahala were fellow students at that time, reminding me of the time that has passed and how another career took a major part of my life before I returned to the university. Though there is something very stable and almost permanent: the Senior Laboratory Technicians Aino Peltola and Antti Louhio. Aino, how many times have we reiterated our experiences with blown-up cables and other flashing memories? And Antti, you’d be rich if you charged even a penny for each invaluable piece of technical advice! All other current and former colleagues in Water and Environmental Engineering - thank you, it has been a pleasure working with you. And drinking coffee, of course!

The most important people in my life have taken my stress, absence and absent-mindedness so calmly, that it would be easy to forget to apologize. My wife Tarja has done so much more than her share to make it possible for me to concentrate on my thesis work. All too often I have not been able to spend time with my daughters, Hanna and Heli, either. And how many times have I told my best friend Essi that I miss her, but so rarely had the time to see her? All you loved ones, I hope I’ll be able to change this now that the thesis is no longer a reason or an excuse!

Finally, I must acknowledge two of the people mentioned above, Tarja and Essi, as well as my MSc supervisor Professor Emeritus Pertti Vakkilainen, as those who made it possible for me to engage in doctoral studies in the first place. I am not sure whether the ladies conspired together to convince me to finish my long neglected master’s studies. Anyway, at the end of 2008 and in early 2009, there was a sudden increase in subtle and not so subtle hints from both that I should do something about finally graduating. It worked – they usually manage to persuade me into, well, almost anything. While I don’t think either one of them ever contacted Pertti, he seemed to play in the same team. When I went to meet him without talking much in advance and asked whether he had any ideas about what to do about my thesis, he just opened a drawer in his desk, slowly looked down and said, “yes, I have this subject here waiting for you.” Mind you, this discussion occurred over 15 years later than Pertti expected! I never saw if he actually had anything on the paper in his drawer, but from there on, all went smoothly as if carefully planned and prepared. Thank you, what you did literally changed my life.

The works included in this dissertation have been funded by a multitude of institutions. My special thanks belong to Maa- ja vesitekniikan tuki ry. that have been involved with funding all parts of this research. Other important funding sources have been Aalto University core funds, Aalto School of Engineering doctoral programme, Academy of Finland, Finnish Cultural Foundation, Strategic Research Council (SRC), Emil Aaltonen Foundation, and Sven Hallin Foundation. I also want to thank CSC – IT Center for Science for computing resources, support, and training in high performance computing.

Espoo, August 2019

A handwritten signature in black ink, consisting of stylized, cursive letters that appear to be 'M. K.' followed by a horizontal line.

Contents

1	Introduction	11
1.1	Eating too much water?	11
1.2	Food for all, limited resources	14
1.3	Research gaps and questions	18
2	Data and methods	21
2.1	Diet change	23
2.2	Food system modelling: Supply chain losses	24
2.3	Food system modelling: Land use allocation with feed demand modelling	25
2.4	Food water use analysis	26
3	Findings	29
3.1	Diet change mechanism	29
3.2	Food supply chain modelling	30
3.3	Impacts of the measures – less water, more food	30
3.4	Linking demand and supply – where does consumption matter?	34
4	Discussion	37
4.1	Making best use of water in the food systems	37
4.2	Key dynamics influencing how much meat we should eat	40
4.3	Perspectives on water use from Cultural Theory	41
4.4	Cultural Theory Interpretation of Food System Modelling	43
5	Conclusions	51
	References	53

List of publications

This doctoral dissertation consists of a summary and of the following publications which are referred to in the text by their numerals

- I. **Jalava, M.**, Kummu, M., Porkka, M., Siebert, S., Varis, O., 2014. Diet change—a solution to reduce water use? *Environ. Res. Lett.* 9, 074016.
<https://doi.org/10.1088/1748-9326/9/7/074016> (Available by CC BY 3.0 license)
- II. **Jalava, M.**, Guillaume, J.H.A., Kummu, M., Porkka, M., Siebert, S., Varis, O., 2016. Diet change and food loss reduction: What is their combined impact on global water use and scarcity? *Earth's Future*.
<https://doi.org/10.1002/2015EF000327> (Available by CC BY-NC-ND 4.0 license)
- III. Kummu, M., Fader, M., Gerten, D., Guillaume, J.H., **Jalava, M.**, Jägermeyr, J., Pfister, S., Porkka, M., Siebert, S., Varis, O., 2017. Bringing it all together: linking measures to secure nations' food supply. *Current Opinion in Environmental Sustainability* 29, 98–117.
<https://doi.org/10.1016/j.cosust.2018.01.006> (© 2017 Elsevier B.V, reprint authorised by terms of use)
- IV. **Mika Jalava**, Joseph H.A. Guillaume, Matti Kummu, Vera Heck, Olli Varis, 2019. How much meat should we eat? Improved estimates accounting for food system dynamics influencing water use. Manuscript submitted on March 23rd 2019 and subsequently accepted for peer review.

Author's contribution

- I.** The author designed the diet optimisation model with supervision by Prof. Kummu. The author implemented the model and performed all analysis, and held primary responsibility for the article. Prof. Siebert provided improved water use data for irrigation. Dr. Porkka mentored with the use of food data and developed the code used in illustrations. Author wrote the article with contributions from all co-authors.
- II.** The author was responsible for the design and implementation of the diet model. The food loss model was adapted by the author from earlier work by Prof. Kummu. The author integrated it with the diet model with support from Dr. Guillaume. The author was responsible for performing the analysis and mainly responsible for writing the article, with contributions from all co-authors.
- III.** The author was responsible for the diet- and loss-related analysis and the diet-related text in the article, design of the analysis, and assisted in writing other parts of the article. Prof Matti Kummu and Dr Joseph Guillaume led the analyses and writing of the article, all co-authors participated. This analysis also relies on results from Publication II.
- IV.** The author designed the Aalto OptoFood model with help and guidance from Dr. Guillaume. The author implemented the model and performed all analysis. Dr. Heck performed the LPJmL runs to generate model inputs. Author wrote the article with contributions from all co-authors. Dr. Guillaume and Prof. Kummu assisted with illustration graphics.

In the end, “the world” is not fed, in aggregate,
and there is no collective “we” doing the feeding.

Prof. Timothy A. Wise

1 Introduction

1.1 Eating too much water?

Water scarcity already affects more than half of the human population, and the trend is increasing (Kummu et al., 2016). While water is a continuously renewable resource, the rate it becomes available is finite (Oki and Kanae, 2006) – the amount of water on our planet is practically constant, but the supply of clean, fresh water is limited at any point in time. As the largest water consuming sector of all human activities (FAO, 2012), food production is at the centre of the quest for using this natural capital in a sustainable way.

Water consumption in food production depends predominantly on two factors: what is produced, and where (Mekonnen and Hoekstra, 2010a, 2010b). These, in turn, are affected by what we eat. People eat food not just to stay alive or for enjoyment, but also to express cultural, religious, and other social traditions (Tiu Wright et al., 2001). The differences between diets and the widely different amounts of water to produce them was the key motivation for Publication I. Many studies (Baroni et al., 2006; Marlow et al., 2009; Renault and Wallender, 2000) compare the resource use of current diets with specific more resource efficient model diets. Resource efficiency of these diets is often improved by replacing animal products with plant-based alternatives while maintaining or improving nutritional values. However, I wanted to analyse the potential to save water through dietary change from a perspective that prioritised variation in the cultural value of diets – defining an acceptable diet change first, and only afterwards assessing its resource efficiency.

Assessing resource efficiency is itself a substantial task. Reducing water use in agriculture requires knowledge about the water consumption in current as well as projected scenarios. This is met by a number of challenges; much of the water used is precipitation directly on the crops, which is augmented by irrigation when needed (Hoff et al., 2010). This water is either consumed through evapotranspiration on site, or further transferred as runoff into streams or lakes, or infiltration into groundwater. Measuring these flows is far from trivial (Wang et al., 2014). Furthermore, different types of natural vegetation have their own characteristic evapotranspiration, not necessarily lower than that of cropland. This means that while converting untouched land to agricultural production may change ecosystems profoundly, water consumption may change more subtly. To complicate the matter further, an agricultural product, like a grain or a fruit, is typically only a fraction of the actual plant grown. And finally, a single crop may yield multiple

products, some of which are not necessarily human foodstuffs. Thus, to determine the water efficiency of food production, we need to understand all the products and services we actually get in return for the water consumed (ISO, 2014).

Food water footprints are a commonly used class of methods to estimate water use in agricultural production and allocate it to products. This in turn allows allocating water use to food demand, and to estimate water use effects of demand changes. Two main methodologies applied to food production estimate volumetric water use per production quantity and then assess human and environmental impacts of the water consumption. The main emphasis of the Water Footprint Network (WFN) methodology (Hoekstra et al., 2011) is on volumetric consumption, while Life Cycle Analysis (LCA) water footprinting (ISO, 2014) emphasises impact assessment. An important feature of the WFN water footprints is the accounting of precipitation used by plants, or green water, in addition to the blue water that is abstracted by humans from surface water or groundwater bodies (see **Figure 1**). Additionally, water spoiled by pollutants can be included as grey water.

Volumetric water footprint methodologies have been criticised for a multitude of problems. Without connection to arable land and other resources, they do not directly correspond to a useful opportunity cost in agricultural production, and production locations will be determined by other factors than water efficiency – water resources tend to be used in some way even when they are scarce (Wichelns, 2015). Other difficulties in volumetric water footprint applications include lack of impact analysis (Berger and Finkbeiner, 2013; Wichelns, 2015) and neglect of return flows (Berger and Finkbeiner, 2013). Nevertheless, while criticised as indicators, the WFN water footprints provide an essential measure of demand-side water use in a multitude of studies, both local and global in scope, including Publications **I–III** in this thesis.

On the other hand, the LCA water footprint analysis has proved a useful tool for comparing the impacts of alternative products and production systems within a limited spatial framework. Instead of attempting to produce a single indicator figure, it differs from the WFN water footprint method by communicating a wider range of measures, and by connecting with other types of environmental impact analysis (Pfister et al., 2017).

Publications **I** and **II** (and indirectly, **III**) used WFN water footprints to assess resource efficiency from the consumption perspective. Diet change, however, by definition means that demand for different foodstuffs will change, which in turn alters production, and hence potentially the foodstuffs' water footprints. The ex-

Figure 1: WFN water footprint colour codes

tensive analysis of food producing systems that forms the basis of both water footprinting methods has been criticised for being based on the status quo and not giving insight to how we got there (Yang et al., 2013). Resource efficiency of crop production depends on such variables as climate conditions, soil, crop cultivars etc. (Fischer et al., 2016). If changes in demand are large, it is possible that the calculated water footprints are no longer accurate. This was one of the motivations for the modelling approach used in Publication **IV**. Instead of water footprints that put emphasis on consumer's choices and the food demand side, it emphasises the food production side by exploring lower bounds for water use using optimised land use patterns that can satisfy the altered demand.

Despite the effort that has been put into estimating how much water is consumed to produce our food and how to reduce it, the concept of water consumption is in fact also open to debate. It is not even clear whether water consumption does indeed need to be reduced. After all, water is consumed by natural vegetation as well as by food and feed crops (Bosmans et al., 2017). Also, the importance of saving water is much higher where water is a limiting factor in agricultural productivity than in areas where waterlogged fields are a bigger problem than drought. There is no single answer to these questions, but taking different perspectives to resource availability and use (see Section 4.3) can explain why some experts emphasise human appropriation of water as the dividing line, while others look into the potential to increase food supply, and treat the environmental water flows as an external constraint.

An example of land use change illustrates the difficulty of valuing human appropriated water flows. Santos et al. (2017) found that in a Brazilian semi-arid region, evapotranspiration (ET) was highest in forested areas and dense savannah vegetation. Grassland and cropland had a considerably lower ET, and bare land lower still, especially in a wet year. If we start with an intact forest, and convert it to grassland or cropland, we are actually decreasing water consumption of the area. Grazing pasture and harvesting cropland decreases the leaf area of the vegetation, which again decreases evapotranspiration. In this example, human appropriation actually reduces water consumption. We can further reduce it by removing the vegetation altogether (Lin et al., 2015). It is obvious that even though evapotranspiration is the main mechanism for agricultural water consumption, high or low ET is not, per se, the deciding factor between “good” and “bad”. We must go much deeper to what purposes the resources serve and, possibly, for whom. My perspective is predominantly that of securing or increasing human food supply, while devoting special attention to finding constraints on food production, such that environmental impacts would be kept in check.

Obviously, while water is one of the most important limiting factors in food production (Mueller et al., 2012), it is only one of the resources required. Others include arable cropland, nutrients and suitable climate conditions, as well as a variety of other inputs in the form of work and materials. Many sectors of human societies compete for and depend on water resources, and water is abstracted from sources outside the immediate vicinity of most of its users. This gives it a unique significance that guided the focus of this thesis towards water, with connections to the other resources where necessary.

Below in Section 1.2, I will take a slightly broader look at the measures to reduce water consumption in food production that are studied in this thesis. In Section 1.3 I identify the research gaps in the field that motivated the thesis and the specific research questions answered by the included publications.

1.2 Food for all, limited resources

Feeding all humanity is an elusive goal. The global food production is currently more than is required to supply everyone with sufficient dietary energy (FAO, 2018a), but lack of economic as well as physical access to food still precludes achieving an acceptable level of food security for all. At the same time, there is a considerable risk that population is growing faster than agricultural production (FAO, 2017).

The term “feeding” can be understood to imply, that someone, or something, actively feeds the human population. This gives an arguably false feeling of control. Professor Timothy A. Wise titled his 2013 scoping paper “Can we feed the World in 2050”, and answered: “In the end, ‘the world’ is not fed, in aggregate, and there is no collective ‘we’ doing the feeding” (Wise, 2013). All measures aiming to improve food security must work in concert with the real-world food systems, ranging from smallholders’ farms providing livelihoods for a handful of people to large-scale industrial farming and food processing. The degree of external control on many of these entities is limited, and due to externalities and other market distortions, economics do not always lead to efficient outcomes from a food supply and resource perspective.

A multitude of measures to increase food supply or decrease resource use and environmental impacts have been studied and proposed. These can be divided into those affecting either the supply or consumption side of the food systems. Supply side measures, such as yield gap closure and spatial reallocation of cropland, have direct and relatively well understood effects on the resource use and supply response. Food loss reduction in the early phases of the food supply chain (FSC) can also be considered to belong to this category. Contrarily, demand side measures like diet change and food loss and waste reduction towards the end of the supply chain affect production and its impacts in an indirect way. Depending on trade structures, the effects on resources and environment may occur almost anywhere the food in question is produced.

Diet change

As noted above, what we eat can have a profound effect on water consumption for food production. In this dissertation, I use the term “food water use” to denote this production-side water consumption. A commonly suggested diet-related measure to improve food security and save resources is to reduce the amount of animal products consumed and replace them with plant-based alternatives with similar nutritional value (Aleksandrowicz et al., 2016). The reasoning is obvious: reduce the foodstuffs with the greatest water consumption relative to the role in nutrition, replace with something more water-efficient.

How much water can be saved, then, by going from one diet to another? At the simplest, the human diet can be approximated as a quantity of dietary energy.

This approach leaves out any concept of a healthy diet, but it allows combining estimates of water use in both plant and animal-based food production; livestock is seen as an inefficient source of energy that converts plant kilocalories into animal kilocalories (Gerten et al., 2011). The water use itself can be approximated by modelled or measured evapotranspiration from typical cropland area.

As the different food items serve quite different purposes, methods considering the actual diet composition rather than just kilocalories give a much better view of what actually changes in the food supply. The water footprint of the current diet can be compared with the water footprint of a specific model diet (Baroni et al., 2006; Blas et al., 2019; Marlow et al., 2009; Renault and Wallender, 2000), with the quantity of animal products reduced from the current average, or even adopting a completely vegan diet. The savings potential of water and other resources that could be achieved depends, in part, on how heavily the current diet relies on animal-based foodstuffs, especially meat (see **Figure 2**). It is easy to see that the potential is far smaller in Asia and Africa than in most of the other parts of the World due to the lower current livestock content in diets – although the consumption of animal products especially in Asia is growing fast (Allievi et al., 2015), likely increasing the relevance of diet change to reduce food water use there, too.

Figure 2: Animal based protein in the human diet. Reproduced from Publication I.

One global diet designed for overall resource efficiency would certainly not be met with unanimous enthusiasm around the world, as evidenced by the mixed reception of the sustainable diet suggested by the EAT-Lancet analysis (2019). Initial reactions in blog posts vary from concerns about nutrition (Harcombe, 2019) and disuse of resources (Clay, 2019) to accusations of a non-scientific approach (Teicholz, 2019). Concerns about the EAT-Lancet diet's effects on the global food

system have already been raised in peer-reviewed literature (Hemler and Hu, 2019).

Ways exist to implement diet change acknowledging local differences in the current food supply. A simple method is to keep the current diet otherwise intact, but replace animal products partly or completely with suitable protein-rich crops. This is the approach used in Publication **IV**. However, this is not sufficient if we want to simultaneously improve the nutritional quality of the diet.

Regionally or nationally focused studies (e.g. Vanham et al., 2018) can take a much more detailed approach by employing food based dietary guidelines (FBDGs), a number of which are listed by FAO (2018b). These divide food items into groups, from which the diet can be composed using accompanying recommendations regarding portion sizes and usage frequencies. With sufficient expert resources, multiple requirements ranging from availability and economics of different foodstuffs to health benefits and cultural desires can be applied to national FBDGs. Designing new FBDGs is laborious and requires extensive knowledge about local food supply and consumption practices. To overcome this, an optimisation-based approach parameterising the animal protein content in human diets was developed and used in Publications **I**, **II** and **III**. It uses optimisation to find a diet as close as possible to the current while following a dietary recommendation (see Section 2.1).

Food loss and waste reduction

An obvious way to reduce resource use and increase food supply is to utilise the food that is produced as completely as possible. Lost or wasted food has consumed resources in production but does not fulfil any nutritional needs. The term food loss represents the foodstuffs spoiled or otherwise lost across the early phases of the FSC (Parfitt et al., 2010): agricultural production, postharvest operations such as transport and storage as well as food processing. Food waste is the term used for the losses occurring in the last two phases: distribution and finally at the consumer's household. For brevity, where food loss and waste are discussed across the whole supply chain, the term loss is used alone to denote both.

Rates at which agricultural products and food are lost and wasted vary across the world. Typically, agricultural practices and methods in industrialised countries help avoid losses at the early stages of the FSC. In the developing world, fewer economic resources are available for investments, compounded with often challenging environmental conditions, leading to higher early losses. At the other end of the chain, the roles switch: economic disadvantage encourages efficient use of food supply at a household level, whereas abundance of money and food allow people to be more selective such that smaller defects lead to wastage. As Parfitt et al. (2010) put it, in traditional subsistence systems “a high proportion of [fresh fruits and vegetables] are consumed because every quality finds a ready consumer within the locality.” Fruits and vegetables are the food items most vulnerable to loss and waste (Gustavsson et al., 2011).

Addressing losses and waste at any food supply chain phase can reduce the amount of primary production required to match food demand, thus saving agricultural resources. In terms of water, the focus of this thesis, 24% of the resource is lost across the supply chain of crop products, according to Kummu et al. (2012).

Their analysis suggests that if best practices among world regions could be successfully implemented everywhere, up to 44% of these losses could be avoided. The methodology of this study was further developed in Publication **II** to cover livestock-based foods and to study the potential interaction between food loss reduction and diet change (see Section 2.2).

Spatial reallocation of cropland

Which crops are produced in different parts of the world depends on a variety of reasons. Obviously, not all crops can be grown everywhere; their efficient production is limited by zones defined by agro-climatic, soil and terrain conditions (Fischer et al., 2016). Nevertheless, resource efficiency is not the only factor leading to the current cropland allocation. It also depends on economic factors, such as the availability of workforce and level of mechanisation, and the local demand structure, which stems from customary diets and policy choices such as national food security, emergency supplies, employment etc. In fact, it has been shown that total production could be increased by 30% by allocating the available cropland more efficiently (Mauser et al., 2015). Another study by Davis et al. (2017) combines an increase in food supply and savings in food water use, feeding an additional 825 million people and reducing blue water use by 12% and green water consumption by 14%.

Spatial cropland reallocation is one of the important dynamic factors studied in Publication **IV**. Changes in the demand of crop and livestock-based foods must be backed by changes in agricultural production, and the eventual resource use depends on where the increases and decreases of each specific crop occur. Linear programming is used to find the best allocation determined by the production-consumption model that takes into account both direct human use of crops as well as the indirect use through the livestock sector (see Section 2.3).

Yield gap closure

Agricultural development, especially the “Green Revolution” (Evenson and Gollin, 2003), has greatly increased the yield of several important food crops. Still, a large percentage of cropland performs at a much lower than optimal level (van Ittersum et al., 2013) and improvements in nutrient and water management could increase yields by up to 70% (Mueller et al., 2012). This yield gap closure, or narrowing of the gap between actual and attainable yields, was studied in combination with diet change and food loss reduction in Publication **III**.

Other water-saving measures

Wada (2014) has presented a “wedge” approach to mitigate water stress. There, six largely independent strategies, or wedges, are defined that either increase water supply or limit its consumption. Two of the wedges, agricultural water productivity and irrigation efficiency improvements are covered above, and have been considered in the methodology of this thesis.

The remaining four wedges are not within the scope of this thesis, which concentrates on food water use. Two of them concern controlling water demand: improvements in domestic and industrial water use intensity, and limiting human

population growth. The last two wedges target water supply, increasing water storage in reservoirs and desalination of seawater.

Additional irrigation could increase crop production by up to 37% when measured in dietary energy (Rosa et al., 2018). Only using currently available sustainable water resources, the potential increase in production is still 24%. Water stress (high use of available water) requires allocation between water users and use sectors (Munia et al., 2016). Any measures relieving water stress may, if deemed necessary, make room for additional food production by allowing increased allocations to irrigation as long as other necessary resources are available.

1.3 Research gaps and questions

Based on the arguments above, evaluation of strategies to reduce water use and its impacts requires an integrated analysis consisting of three main steps: Defining a diet change mechanism, Food supply chain modelling and Evaluation of water use and impacts (**Figure 3**). Within each of these steps, this thesis has examined a number of key questions described here.

Figure 3: Main analysis steps of diet change effects on water use

Diet change mechanism

Predetermined model diets (Baroni et al., 2006; Marlow et al., 2009; Renault and Wallender, 2000) and food based dietary guidelines (FAO, 2018b) both have their strengths and weaknesses. Model diets are simple, and in a global context one average diet is all that is required. Using multiple model diets spatially quickly becomes difficult. FBDGs can accommodate specific requirements like those set by food availability, culture and economics. Their challenge is that the guidelines by different countries are created using different food item categorisations and many dietary recommendations are unique to specific nations, complicating parameterisation for different levels of diet change. Also, where an FBDG does not exist, putting one together requires extensive country-specific knowledge. A compromise solution was needed. Publication I aims to answer the Research Question 1 (**RQ1**): **how can we determine global diet change goals in a parameterised way?**

Food supply chain modelling

Diet change, moving away from animal products towards plant-based alternatives, aims to reduce resource use and increase food security. Alone, it is not suf-

ficient to enable the required increase of 50-70% in food supply (FAO, 2017; Tilman et al., 2011) that is needed to support the expected needs of the humanity in year 2050 (see Section 2.1, results of Publication I). Reducing food loss and waste serves the same goal by decreasing the amount of production required to fulfil food demand. Combining the strategies is complicated by the fact that changes in one element may affect others in non-trivial ways. An intentional oversimplification illustrates a very strong case of dependence: If we first reduce water use by, for example, 52% by removing the animal products from the diet (Renault and Wallender, 2000), then trying to reduce losses from animal products is pointless. More subtly, existing data showed differences in food loss and waste percentages between food groups (Gustavsson et al., 2011), with lower average losses for meat (17%) than for plant-based foods (with fruits and vegetables at 39%). We hypothesised that switching away from meat would increase losses, increasing the benefits of loss reduction. The overarching research question for Publication II is: **To what extent are water savings from diet change and food loss reduction interconnected (RQ2)?**

Evaluation of water use and its impacts

Global food systems involve a number of dynamics between crop production and human food consumption, making certain combinations of products synergistic. As seen above, water-saving strategies may target one or multiple elements in the food systems, and numerous studies have presented results concerning their effect on the total food water use. The combined effects of multiple strategies applied together are less well known. Publication III aims to characterise **what is the national-level combined potential of diet change, food loss reduction and yield gap closure in increasing human food supply (RQ3)?**

Publications I-III use the WFN water footprint analysis method to assess the consumption-based water use of different diets and the potential water savings from food loss reduction without having to model the whole production chains. The large production changes required to achieve meaningful savings prompted the question investigated by Publication IV: **should food water use be assessed through production modelling instead of static consumption-based footprints (RQ4)?**

Interpretations of food system modelling choices

Assessment of changes to the food system is fundamentally dependent on the methods used. In addition to the research questions answered by the Publications I-IV, in Section 4.4 of this thesis I therefore discuss how food system modelling depends on the modellers' decisions, often affected by their personal judgment and, potentially, background. I use the framework offered by the Cultural Theory, briefly introduced in Section 4.3, as a simple categorisation of worldviews to illustrate how different values affect modelling outcomes.

2 Data and methods

In this section I present the main methods used and developed in this thesis, together with the data employed. **Figure 4** illustrates the expanding scope of the publications, as additional aspects of food systems and methodologies to address their water use were introduced. While in terms of scope, Publication **IV** covers the diet change and loss reduction strategies explored by the earlier publications, the methods and data used are different. This is due to the methodological improvements specifically sought after (see above in Section 1.3), and the incommensurability of the data when going from country and region level analysis to spatial modelling. In **Table 1**, the methods specifically developed or extensively adapted for this thesis are presented with their main data sources. Publication **III** additionally employed methods and data not in the scope of this thesis, contributed by the other authors.

Figure 4: Scope of the publications included in this thesis

Table 1: Methods and main data sources overview

<div>Model element</div> <div>Ap- proach used in publications</div>	Diet change	Food system modelling	Food water use as- sessment
Publication I	Minimise change from current diet Diet data: FAO FBS ¹	Consumption-side analysis, differen- tiating local and imported produc- tion – change in production is not explicitly modelled	Water Footprint Analysis Data: Water Foot- prints (Mekonnen and Hoekstra, 2010a, 2010b)
Publication II		Sequential loss calculation adapted from Kummu et al. (2012) Loss data: Gus- tavsson et al. (2011)	
Publication III			
Publication IV	Replace combined livestock-based food with pulses and soy- beans Diet data: FAO FBS	Land use alloca- tion using spatial optimisation of cropland use to fulfil demand by humans and live- stock Cropland data: Portmann et al. (2010) Crop properties: LPJmL ² Loss data: Gus- tavsson et al. (2011)	Optimised land use Crop evapotranspira- tion (LPJmL) Irrigation water ab- straction (LPJmL)

¹ Food Balance Sheets (FBS) and Commodity Balance Sheets (CBS) in FAOSTAT are currently available at <http://www.fao.org/faostat/en/#data/> (FAO, 2019)

² Lund, Potsdam, Jena managed Land model (Bondeau et al., 2007)

2.1 Diet change

The novel method developed in Publication I for defining diet change scenarios (see **Figure 5**) uses optimisation to address three requirements by i) parameterising reduction of livestock products in diets at country level to allow comparison between different levels of animal protein, and ii) incorporating adjustments to national diets to follow a dietary recommendation (WHO, 2003), while iii) minimising the changes from the original diets.

For modelling purposes, national, regional and global average diets are better described by calculating average consumption of food picked from a number of food item groups rather than a fixed collection of specific food items. This allows changes in the diet by shifting consumption between the groups instead of replacing one food item at a time with another.

The diets are expressed in terms of 13 food item groups, each with protein, fat and dietary energy content derived from the FAO Food Balance Sheets (FBS). Optimisation is used to adjust the amount of each food item group in the diet.

The livestock reduction and dietary recommendation requirements are implemented together by first replacing the required percentage of livestock products with a mix of oilseeds and roots, and then building a set of optimisation constraints on nutritional content, amount of fruit and vegetables and restrictions on specific foodstuffs according to scenarios (for the actual constraints, see Publication I, Table 2). The changes from original diets are minimised by quadratic programming, using the sum of squares of the national differences from the current diet as the objective function, and optimising subject to these constraints.

Figure 5: Diet change optimisation method developed for Publication I, used also in Publications II and III

The same diet change optimisation method was used to analyse diet change effects in Publication **II** combined with food loss analysis. In Publication **III**, this combination was further combined with yield gap closure to assess the global food supply potential to meet the food demand projections for the year 2050.

An alternative, somewhat simpler diet change method was used in Publication **IV** to better detect the effects of livestock replacement on water use, without additional influences imposed by the dietary guidelines. The method was adapted from Vanham et al. (2013), who used it to determine reduced-livestock food based dietary guidelines. Livestock products are replaced by a combination of pulses and soy with similar dietary energy and protein content. An exact match in both parameters is not typically possible, and in this study, preserving protein was prioritised. As feed data was only available as an aggregate across all livestock species, the ratio between different livestock products is kept constant to avoid changes in feed composition and conversion efficiency.

2.2 Food system modelling: Supply chain losses

Publication **II** adapted the FSC loss and waste modelling approach from previous work by Kummu et al. (2012). It applies loss ratios of different FSC phases reported by Gustavsson et al. (2011) sequentially to the part of crop production (FAO FBS) directed towards human food consumption. The method was substantially improved by adding livestock-based food groups to the analysis and extending the water footprint computations to include green water (see **Figure 1**). Furthermore, two adjustments were made to the loss calculations. Processing losses³ are applied to processed foods (e.g. bread) instead of primary commodities (e.g. flour). This was necessary because not all processing outputs are food products, like the press cake from oilseeds which is used predominantly as feed. Postharvest loss⁴ modelling was also changed. The original choice of assigning these losses to producing countries was based on the assumption that much of the postharvest losses occur already at the farm level, which appeared to be well-justified (Afgnon et al., 2015). In the FAO FBS methodology, however, the waste element is considered a part of the consumption accounts covered by domestic food supply. Therefore, in the new model the postharvest losses were assigned to consuming countries instead of producing countries, consistent with a consumer-oriented footprint analysis. The FBS waste element is used interchangeably in the model with postharvest losses as the former is not available for all countries.

Publication **II** applies scenarios for reduced losses across the supply chain. The results are then also used in Publication **III**. Aalto OptoFood adjusts demand to take into account changes in supply chain losses using a similar process, but Publication **IV** does not apply loss reduction scenarios.

³ Losses due to spillage and degradation in either industrial or domestic food processing (Gustavsson et al., 2011)

⁴ Losses occurring during handling, storage and transportation between farm and distribution (Gustavsson et al., 2011)

2.3 Food system modelling: Land use allocation with feed demand modelling

In Publications **I–III**, a consumer perspective is used – the effect on production is not directly modelled. Instead, the analysis uses predetermined product water footprints (see next section, 2.4). This assumes that changes in demand merely lead to scaling up or down of the current mix of production systems, with a constant efficiency, productivity and impact per unit of production.

Tracking the actual source of all food consumed in a country is a complex task (Kastner et al., 2011), and changes in food consumption induce changes in trade patterns. The analyses in these publications do still, however, differentiate between domestic production and imports.

In contrast Publication **IV** develops a model of the food system, named *Aalto OptoFood*, to optimally allocate land and water resources to production globally in order to match human food demand. Changes in production resource efficiency can then be assessed spatially, as described below in food water use analysis (see Section 2.4).

Figure 6: Integrated food system modelling approach with Aalto OptoFood

The core of Aalto OptoFood is a linear programming model. For a given objective function, it optimises a set of decision variables: cropland use for each of 14 crop

functional types (CFTs) at 0.5-degree grid level, and regional production of six livestock-based food items.

The optimisation is subject to a number of constraints (see Table S4 in Supplementary Information of Publication **IV** for details). The main assumptions affecting the results set forth by the optimisation constraints are the following:

- i) *No expansion*: Cropland, irrigated area and irrigation water use are limited to their current amounts
- ii) *Landuse ‘stickiness’*: Production can only be moved for crops with changing demand. If production of a crop increases, all existing production must also be preserved along with the additional land, and if it decreases, land can only be removed and no new areas can be used for its production
- iii) *Livestock stickiness*: Livestock production is not allowed to increase in any region, as all scenarios call for decreasing total demand
- iv) *Demand constraints*: Demand must be met by production. Extrinsically set human food demand and intrinsically required livestock feed must be supplied for by crop and roughage production

2.4 Food water use analysis

Publications **I–III** use Water Footprint Network product water footprints (Meconnen and Hoekstra, 2010a, 2010b) to estimate water consumption as virtual water embedded in the human food supply. Here, water consumption is presented as the sum of blue and green water (see **Figure 1**). Grey water, being the water needed to dilute contaminants or salts to acceptable levels, is omitted. This is because the water may still be usable for other purposes, and secondly, other measures to control pollution exist, like buffer zones (Parkyn, 2004). Thus, while changes in freshwater quality are an important impact of food production, this can be addressed at least partly without changing which food items are produced.

The diet change and food system modelling differentiated demand by food group, country, and by whether it was produced domestically or imported. Water footprints for each country were therefore averaged across the current composition of the food groups used to represent the diet (see 2.1). Because the water footprints of food items vary between production areas, an identical diet in different parts of the world may have a different water footprint. The average water footprint of the exported food is calculated as the total blue and green water footprints of the products in the exporting countries, divided by total mass of the food.

Publications **I** and **II** also estimate changes in Green-Blue Water Scarcity. Green-Blue Water Scarcity measures water availability compared to the water needed to produce the energy in a reference diet (Gerten et al., 2011). Reduced water requirements are calculated by scaling down water use. Estimates of local impacts of production on water are not possible because changes in production are not explicitly modelled.

Publication **IV** differs from the preceding publications in that it determines food water use from grid-level crop evapotranspiration data and optimised land use. In LPJmL-generated input data from an “all crops” run (Bondeau et al., 2007),

evapotranspiration is available for all cells where each crop can grow, allowing water use estimates also when land use is altered so that some areas are converted to crops not grown there previously.

Explicit representation of production also allows broader consideration of impacts of water use. The Aalto OptoFood model allows any given objective to be used for land use allocation. Publication **IV** applies the following objectives: MinET (minimising the global sum of evapotranspiration), MinStress (minimise average stress among hydrological basins) and MinDiff (minimise sum of absolute differences from natural vegetation evapotranspiration). Additionally, when using MinET, part of grassland can be set aside as rangeland. This sets the evapotranspiration of grass in affected cells to zero for the optimisation runs. When used, this option implies that crop (or grass) evapotranspiration is to be considered a food production resource instead of a measure of human appropriation of green water. These objectives correspond to different perspectives on impacts of water use (see Section 4).

3 Findings

Consistent with the research questions in 1.3, this thesis provided findings on both methodological issues and impact of diet change on water use.

3.1 Diet change mechanism

RQ1: How can we determine global diet change goals in a parameterised way?

The novel diet change mechanism developed for Publication **I** was designed to allow parameterisation of the human diet in terms of livestock-based food content. It allows analysing the water-saving potential of country-level diet change scenarios without forcing one model diet globally. The five scenarios presented were characterised by a cap on the animal-based protein and, separately, meat protein in the human diets. Parameterisation of the model using optimisation constraints allows detailed, easily extensible control over the constituents of the resulting diets.

The diet change mechanism proved sufficient for a consumption side analysis, and a good basis for the combined modelling of diet change and food loss reduction in Publication **II**. A challenge, acknowledged in the discussion of Publication **I**, was the reduction of protein and fat in the global food supply and many of the national diets. Applying a single dietary recommendation (WHO, 2003) that accommodates person-to-person variation to nations may in some cases lead to changes that would be impractical, because the safe limits of population-level variation are smaller. This does suggest that improvements should be developed that allow novel ways to analyse changes in the diet than straightforward replacement of food groups with predetermined alternatives, such that tighter constraints on population level nutrition would be feasible without extreme shifts in food group ratios.

Publication **IV** takes a production side perspective to diet change. The emphasis is on the reorganisation of cropland use as a result to changing demand. This led to adoption of a simple animal protein replacement with pulses and soy, such that the water use and land allocation results are easily interpreted as a function of a single diet change level parameter acting on the demand side.

3.2 Food supply chain modelling

RQ2: To what extent are water savings from diet change and food loss reduction interconnected?

Publication II successfully demonstrated that there can be a further dependence between the food water use reduction potentials of diet change and food loss and waste reduction than the expected multiplicative effects of chained reductions. The practical effect of this specific dependency is relatively small with the method and data used, yielding results close to the simple multiplicative effects.

Figure 7: Summary of the findings of Publication II, comparing water use for multiple scenarios: original diet (OD), recommended diet with a cap of 25% livestock based protein (RD_A25), a food loss reduction scenario halving losses at all stages (HalfLoss), and a combined scenario (RD_A25_HalfLoss). Reproduced from Publication II.

Global blue water footprint reduction was 0.7% larger (green WFP reduction 0.6% larger) for combined diet change and food loss reduction scenarios than with the scenarios applied separately (see **Figure 7**). Regionally, the greatest difference in total water footprint (blue+green) reduction between combined and separate scenarios was up to 2.9 percentage points (pp) in North America and Oceania. At national level, the maximum difference was found to be -5 pp in green water and -8 pp in blue water (Guinea-Bissau). While small, the result is intuitive: when foodstuffs with low loss rates (e.g. meat, with global FSC losses of 17%) are replaced with ones with higher loss rates (plant-based foods, highest losses being with fruits and vegetables at 39%), reducing the losses generates greater savings.

3.3 Impacts of the measures – less water, more food

RQ3: What is the national-level combined potential of diet change, food loss reduction and yield gap closure in increasing human food supply?

In agreement with an extensive body of research using different methodologies, diet change was found to be an effective way to reduce the water footprint of human food consumption, according to the optimisation-based algorithm presented in Publication I. Converted to additional food supply, the water saved globally could theoretically be used to produce and supply food for 1.8 billion people.

Yet the new and more interesting result was the unevenness of the water footprint reduction due to differences in the current diet composition and sufficiency as well as the water footprints of the locally produced food. In **Figure 8**, increasing blue water footprints by country are presented as “bubbles” to the right of the vertical axis, and increasing green water footprints are above the horizontal axis. Of the 176 countries studied, 74 had a caloric deficiency, leading to increases in water footprint when switching to a diet following the WHO recommendation. Also, even in some countries where the algorithm led to reduction of dietary energy, at least some of the water footprints unexpectedly increased. In Finland, for example, the diet change towards plant-based foods led to an increase in the blue water footprint in all scenarios. This effect was tracked to the Finnish livestock production using predominantly rainfed feed supplies, but the replacing protein crops being partly sourced from the global import pool utilising irrigated cropland.

While lower than the maximum water saving potential of diet change, the effect of reducing food loss and waste is still substantial. The improved loss and waste model presented in Publication II that included livestock-based foodstuffs and green water footprint analysis showed that roughly 12% reduction in food water use can be attained by halving the loss and waste. This very closely matches the prior study by Kummu et al. (2012) that only analysed the blue water footprint of the crop-based part of the human diets. Furthermore, the water savings were very similar between a “best practices” MinLoss scenario, choosing the lowest loss factor for each World region and food group, and the HalfLoss scenario in which the loss at each phase was simply halved. This emphasises the intuitive robustness of the loss reduction strategy – differences in loss factors of different food groups and food supply chain phases even out. Reducing food losses by weight over the entire diet reduces food water use in roughly the same ratio. This is further supported by the synergy between diet change and loss reduction discussed above in Section 3.2.

While Publication III evaluated water savings to estimate the effects of diet change and loss reduction, the main results were presented in the form of potential to increase food supply by combining these strategies with yield gap closure. Although the methodology used to combine the effects of the different strategies was decidedly simple, the results build confidence that the human population in 2050 will be able to feed itself. At global scale, assuming trade could level out differences in food self-sufficiency, moderate to high level of implementation of these strategies were found to enable 111-223% increases in global food supply. The differences in food supply boosting potential between areas are highly variable, though, which means that the role of trade is crucial. At the country level, over 320% increases in food supply against projected population were observed on all

inhabited continents. But, alarmingly, with current trade levels even the high implementation levels of the supply-increasing strategies leaves most of Sub-Saharan Africa far below self-sufficiency in the year 2100. Moderate implementation will fall short of this goal already by 2050.

Figure 8: Effects of diet change on water footprints. Location of the bubbles represent water footprint changes when country level diets are adjusted from the Original Diet (OD) to the Recommended Diet (RD) scenarios following the WHO dietary recommendation. (B). Water footprint changes from RD to A25, setting a cap of 25% on livestock-based protein in the human diets. (C). Water footprint changes from RD to A0, where livestock-based protein is removed. Size of each bubble represents the population of the countries depicted (see legend). For further description of the scenarios, refer to Publication I, Table 2. Reproduced from Publication I

A number of key areas were identified to improve future analysis of combined scenarios, divided into three categories: core, supportive and linking developments (see **Figure 9**).

The core developments list five central elements needed to fill the knowledge gaps in quantitative modelling of the food system. CA1 (core action 1), fully integrated food system modelling, enables recognising feedbacks between measures applied to different parts of the system. CA2, climate change integration, is crucial

both due to the inherent connection between the global food production and greenhouse gas emissions as well as the inertia in changing the global food system; the time scale of changes may make the required increase in food supply a moving target. CA3, consistent scenarios, emphasises the importance of making the results of studies commensurate. CA4, inclusion of non-agricultural food sources, e.g. fish, aims to complete the picture of human diets. This is becoming even more crucial as aquaculture is becoming the dominant source of seafood and an increasing share of the inputs are derived from agricultural products (Gephart et al., 2017). Finally, CA5, filling in the value chain gaps, is required to complete understanding of the interactions between different elements of the food supply chain and agricultural sector. Concentrating on one element at a time may make material flows that are inputs to other important elements look like losses.

Figure 9: Methodological improvements suggested to obtain a more reliable understanding of the combined potential at country-scale to sustainably secure food supply, divided into three categories: core actions (CA1–CA5), supportive actions (SA1–SA4) and linking actions (LA1–LA4). The main linkages of these actions are shown as dashed lines. Reproduced from Publication III.

The supportive developments identify important context to understand the food supply potential and to model future scenarios. The linking developments, in

turn, identify issues requiring further knowledge and connections with other disciplines. These areas for improvement were, in part, a motivation for the Aalto OptoFood model presented in Publication IV.

3.4 Linking demand and supply – where does consumption matter?

RQ4: Should food water use be assessed through production modelling instead of static consumption-based footprints?

Figure 10: Change in land use from current diets to 20% reduced livestock using the objectives A) minimum evapotranspiration, and B) minimum difference from natural evapotranspiration. The land use change due to water stress minimisation is not shown, as the land use is not sensitive to rainfed agriculture, leaving the cropland use in low-stress areas random. Reproduced from the Supplementary Information of Publication IV.

Modelling food production using Aalto OptoFood and its three optimisation objectives allows a wider look into food water use and its impacts than demand-side water footprint analysis. Reallocating global cropland at increasing diet change levels shows a pronounced effect with all objectives (see **Figure 10**). In each case,

the effect is strongest in the initial phases of the shift away from livestock products. This suggests that while livestock production is consuming a large amount of water, causing considerable amount of water stress and changing the hydrological cycle with changes to evapotranspiration, relieving these issues is possible already with a moderate reduction of livestock-based food in our diets. Depending on the percentage of grassland that is non-arable rangeland, food security may benefit from some livestock production rather than leaving rangelands unused and replacing lost protein supply by crops. Assuming grassland on 75% of the grid cells is non-arable, up to 19% of the protein in the average global diet could be livestock-based products without compromising food security by inefficiently using water resources that could be better utilised by protein crops. Lower rangeland estimates push that efficient livestock protein percentage down. Still, assuming land use changes could be optimised, in all scenarios a majority of evapotranspiration reduction achievable by diet change can be obtained by cutting the livestock percentage in our diets from 34% down to 28%. Similarly, a comparable reduction in livestock percentage achieves most of the reduction in both water stress and deviation from natural ET – although with a very different land use allocation, illustrated by **Figure 10**.

Water footprints do take spatial variation into account but they are typically determined for current production only, instead of all potential food producing areas within the current agricultural land use extent or outside it. The amount of protein-rich crops to replace livestock products is large: 129% more pulses and 39% more soybeans need to be produced if all meat, milk and eggs are to be removed from human diets. This suggests that it is, indeed, better to model food water use through production modelling than using consumption-based water footprints.

4 Discussion

Food security and agricultural sustainability studies typically focus on reduction of meat and other animal products, sometimes abandoning them altogether, rather than on finding whether they provide a positive contribution to resource use or other measures of sustainability. Still, several relatively recent studies suggest that in some circumstances, some of the most inefficient food systems like grazing ruminant livestock can supply some food virtually for ‘free or using “leftover” resources’ (Davis and D’Odorico, 2015; Godfray et al., 2010; Rööß et al., 2016). Obviously, what “free” means depends on the definition of cost. In this thesis, I examine the cost mostly in terms of water consumption, and to some extent land use. The overarching theme is using water efficiently to match a predetermined food supply goal and avoid the unsustainable use of resources. In order to prioritise this objective, additional benefits that arise from the approaches explored are not made the subject of extensive enquiry. Even within this limited scope, water use efficiency and sustainability can still be seen from several perspectives.

The key findings presented in the previous section are well aligned with previous research, but extend it with important new knowledge. In this discussion, I first compare the new scientific findings with existing research in the field of the three overarching themes of the thesis: diet change, food system modelling and food water use analysis. I also identify the key dynamics in the global food systems that fundamentally affect the value of livestock. To illustrate why food production and consumption are controversial not only among consumers, policymakers and other stakeholders but also among researchers, I then take a look at how Cultural Theory (Douglas and Wildavsky, 1983) has been used to describe different views regarding water resources and discuss how food system modelling decisions can be interpreted in that context.

4.1 Making best use of water in the food systems

Diet change

Much emphasis has been put on the water consumption of food production, and water footprints (Hoekstra and Hung, 2002) quantifying the water volume consumed at all phases of producing a specific food item have started to appear in consumer marketing messages and stakeholder communication (EU Retail Forum, 2014). The findings of this thesis point out that not only is it important where the water footprints “land” and what the impacts are (Pfister et al., 2017), but also

that the effects of consumer actions, such as diet change, have considerable variation based on where they are applied (see **Figure 8**). This result from Publication **I** suggests that demand-side measures aiming at modifying the human diets may be more effective in reducing water consumption and scarcity if localised instead of relying on recommendations based on global estimates or generalising analysis based on data from limited areas (Baroni et al., 2006; Marlow et al., 2009; Renault and Wallender, 2000). This will require further study, but the diet change methodology developed in Publication **I** should provide a good starting point for such research. Although regional studies relying on food based dietary guidelines (Vanham et al., 2018, 2013) allow more freedom in defining diets according to multiple goals such as required in national dietary recommendations (FAO, 2018b), algorithmic diet scenarios like those presented in Publication **I** can easily be extended to multiple scenarios in a large number of study areas.

Food system modelling and issues with food water use analysis

While useful in assessing the food water use effects of changes occurring purely on the demand-side, the use of WFN water footprint analysis in Publications **I–III** brought up a number of problems with the indicator itself, and potentially with other water footprints as well. Some of this criticism has been presented by others (see Section 1.1).

First, pre-calculated product water footprints only represent the status of the production systems at the time of the analysis (Yang et al., 2013). In such a case, the food system model is at least partly embedded into the indicators themselves. For example, the WFN analysis (Hoekstra et al., 2011) allocates the water use over different product fractions (see Section 4.2 for discussion regarding food and feed) according to market values and yields of those fractions at a certain place and time. As many of the measures to improve food supply or reduce resource use specifically target some of these variables, for successful modelling of large changes in the systems they need to be disentangled from each other. In practice, this requires that water use is modelled as a part of food system modelling, not as a separate task aiming to make specific water use an intrinsic property of each food item.

Second, volumetric water footprints only attempt to represent consumptive water use (including evaporation and water incorporated into products). This is mainly useful if our attention is focused on total water consumption, such as if water is considered a universally useful natural capital (Hoekstra, 2009). A major drawback of this view, specifically targeted by impact-oriented water footprints (ISO, 2014; Ridoutt and Pfister, 2013), is that the value of water in human use is highly dependent on location and availability of other resources. However, if our concern is large-scale change from the natural state, such as feedback between evapotranspiration and precipitation, neither the water volumes nor local scarcity or degradation impacts alone can be used. Spatial distribution of cropland, natural vegetation and other land covers determines where and how much evapotranspiration occurs. While actual determination of how this in turn affects precipitation and thus water availability would require integration with climate analysis, the differences between results using alternative Aalto OptoFood optimisation objectives clearly show that modifying the food systems in order to reach different

goals can lead to very different outcomes (see **Figure 11** A, D and G, and **Figure 10**). None of these can be captured by product-based indicators that attribute changes only to the food demand, as the behaviour is strongly driven by changes in spatial distribution of production and across different elements of the food supply network.

Figure 11: Optimisation objective globally and spatially with alternative objective functions: (A-C) minimizing evapotranspiration, MinET, (D-F) minimizing average water stress, MinStress, and (G-I) deviation from natural evapotranspiration, MinDiff. Average water stress corresponds to the percentage of evaporative water consumption for irrigated agriculture relative to availability, averaged across all basins. Water stress is not sensitive to changes in rainfed agriculture, and ET and deviation from natural ET are therefore not shown for that objective. Reproduced from Publication IV.

4.2 Key dynamics influencing how much meat we should eat

Modelling diet changes in the global food system requires understanding some key dynamics between crop and livestock production sectors: the use of crops as feed and the land use competition between the sectors. Their combined effect is quantified in Publication **IV**. Here I give a qualitative summary to assist in understanding and interpreting food system modelling decisions in Section 4.3.

Fundamentally, the use of crops and land is a question of allocation. While allocations may be shifted by decisions stemming from economics, regulatory pressure or other reasons, they are at least partly determined by production and resource efficiency, as part of a complex system (Nesheim et al., 2015).

The value chains of various foodstuffs comprising the human diets are very different. Some food items like fruit and vegetables can be tracked across their supply chain relatively unchanged from field to consumer's plate. Others, like cereals or oilseeds, go through more involved processing steps before becoming food ready for consumption. Each step involves losses, and some of the crop or its components can be diverted to other uses. The livestock sector shares these properties with crop-based food production, but is interlinked with it through the inputs and, at least through manure, also through outputs.

Many crops are typically used both for human food and animal feed, with the most important exception of grasses as the main component in the diet of ruminant livestock. A common suggestion to improve food system efficiency and food security is to use the food crops directly instead of feeding them to animals (Clark and Tilman, 2017; Falkenmark and Lannerstad, 2010; Pimentel and Pimentel, 2003; and many others). After all, the animals require energy and nutrients for purposes that do not directly contribute to edible outputs. Thus, feed conversion efficiency, or the rate at which livestock can convert feed inputs into edible outputs, makes feeding food crops to animals and then eating the meat, milk and eggs inherently less efficient compared with consuming the crops directly (Mekonnen and Hoekstra, 2010, Vol 2.).

The situation is less straightforward if we take a more detailed look into the actual composition of food and feed flows. First, requirements for food-grade crops are more stringent than those for livestock feed (Bitzer and Riddell, 1984; Herrman and Kuhl, 1997). Livestock can utilise crops that would be considered of low or no value as human food, thus making otherwise potentially lost products again valuable elements in the human food supply chain. For example, between the years 2005 and 2017, the percentage of yearly wheat production in Finland that fulfilled the national food requirements varied from 14 in year 2017 to 77% in 2011 (LUKE, 2018). According to the same study, for production of oats intended as food, the quality requirements were fulfilled by 3 (2010) to 63% (2012) of the production. The variation can be assumed greater in Finland than in many other countries due to the climate with highly variable weather conditions during the growing period, but it serves to point out that not all crop production can be directed to human consumption without further action.

A second consideration is the intentional co-production of food and feed. According to the FAO Commodity Balance Sheets, less than half of rapeseed produc-

tion goes to human food, the rest being diverted to livestock feed. But these fractions are completely different – the food fraction is rapeseed oil, and the feed is the press cake remaining after the oil is extracted. Obviously, these products are not interchangeable. Economically feasible production of certain crops, oilseeds being a typical example, is largely dependent on market demand for multiple components – including those not used for human food.

The interface between food and feed production, however, is not fixed. While utilising substandard crops or oilseed press cakes directly as food may be infeasible or undesirable, those fractions may be useful in other ways than as livestock feed. Novel ways to utilise lower qualities or previously unused crop components in food processing industry could enable their utilisation directly in the human food supply chain, and other uses such as biofuel, bioplastics etc. can compete with feed use, changing the economics of the livestock sector. These alternative uses are intentionally left outside the scope of this thesis, given that it concentrates on actions improving rather than decreasing food security and resource efficiency of food production specifically, but their role is briefly discussed in Section 4.4.

4.3 Perspectives on water use from Cultural Theory

The system dynamics described above, along with other choices regarding model structure, data and constraints, can significantly affect modelling results and their interpretation. Unless these choices are clearly communicated and justified, the work may appear biased (Brunson, 1992). Many results and findings relevant to environmental policies are often challenged on the basis of real or perceived biases (Doremus, 2008). In fact, Huesemann (2002) argues that while completely unbiased environmental research is impossible, it is useful to recognise and reflect on one's own self-interest as well as institutional and socio-cultural background.

To illustrate how different perspectives to the same underlying issues affect research results, I am using Cultural Theory, or Cultural Theory of risk. It is a conceptual framework originating from the work by Mary Douglas (1970) and later formalised by her and Aaron Wildavsky (1983). It argues that social organisation strongly affects how people perceive and want to cope with societal dangers, and that these survival strategies further strengthen the societal structures. The theory assumes a particular classification of worldviews, discussed below, based on high and low positions over two dimensions labelled “group” and “grid”. Positioning on the group axis reflects the level of collective control by societal groups individuals associate themselves with. The grid axis describes the more fixed societal structures that establish status, authority or subordination (see **Figure 12**).

One approach to classify different perspectives to resource availability, use and environmental risk in terms of the Cultural Theory is presented by Hoekstra (2000). He describes a set of four perspectives to valuation of water by drawing on Cultural Theory, extending the theory's four worldviews.

Figure 12: The Grid-Group representation of Cultural Theory worldviews, adapted from Schwartz and Thompson (1990). The Grid axis represents external restrictions on individuals’ choices, where a prescribed behaviour has a high score and individually prescribing behaviour scores low. The Group axis represents the collective aspect of behaviour, differing from the Grid by self-assignment instead of prescription.

The individual worldviews are defined in considerable detail by literature. Hoekstra interprets the theory in terms of water resources management by giving the worldviews clear-cut stances on a number of field-specific aspects. These worldviews are only an example of how differences and incompatibilities in thinking about resources, needs and management options can be described in a systematic way. Cultural Theory has been criticised for confusing societal needs with individual’s perception of risk, and for stemming from a limited view to what culture is, at best describing different American political groups’ views to risk (van der Linden, 2016). Nevertheless, together with Hoekstra’s water resources specific interpretation, it does offer a framework for qualitative discussion regarding the importance of taking subjective choices into account in model development and use. The precise definition of these perspectives is open to debate, and my focus here is to provide some heuristics to help in interpreting approaches to food system modelling. The following characterisations are therefore only intended to give a superficial overview of the worldviews.

The low-grid, low-group *Individualist* worldview is presented by Cultural Theory as one that is willing to exploit the system’s abundant resources, fulfil individual’s needs and manage possible scarcity by market economy. Nature is seen as benign, resistant to change. Hoekstra’s interpretation of the Individualist’s perspective in the context of water resources emphasises the economic mind-set: allocation of resources is price-driven (group coordination is not essential), and problems are solvable by money and economic growth (there are few constraints on action).

The low-grid, high-group *Egalitarian* sees resources as scarce and depleting, unmanageable, and expects human needs to be managed socially to achieve sustainability. Nature is considered fragile; any perturbations could lead to sudden degradation. Water demand is a manageable desire, and water scarcity is a problem with excessive demand, to be solved through collective action.

The *Hierarchist* worldview characterised by high position on both axes takes resources as scarce, but manageable – if demand exceeds supply, they should both be managed with regulation. Nature is seen as tolerant to reasonable extent, but large perturbations should be avoided. Hoekstra suggests that the Hierarchist prefers technological approaches to increasing water supply rather than manages the need. The system is amenable to being controlled by a well-managed effort.

The high-grid, low-group *Fatalist* takes a passive view to life. Resources cannot be managed, their abundance is determined by luck. Nature is unpredictable, risks just have to be accepted. The Fatalist considers water scarcity a problem of individuals; water demand cannot be managed, and neither can availability. Emphasis is on reacting to events and stresses as they occur.

4.4 Cultural Theory Interpretation of Food System Modelling

In this section, I reflect on some important food system modelling choices using the Cultural Theory worldviews introduced above. Each model takes a certain perspective on the system or phenomenon it represents. Most questions cannot even be answered without accepting that at least some subjective judgements are being communicated or are embedded within results due to the effect of the judgements on model implementation. This is not limited to food system modelling. Even fundamental physics struggles between models and their interpretations, when all important questions cannot be answered by one concise framework. A well-known example of this is wave-particle duality, where two models of electromagnetic radiation both explain a different set of observable phenomena (e.g. Milonni, 1984). While in physics the different views typically stem from limited understanding of the natural phenomena, in fields that concern human societies, the imprecision of measuring what happens in the physically complex system is compounded with human preferences and impressions about what is desirable.

In food system modelling, non-trivial choices have to be made, because the systems are much more complex than can be captured in detail by any reasonable model or data available now or in the foreseeable future. Some of the choices are technical in nature, having practical implications in modelling accuracy, coverage, complexity of the analysis etc. While they may impact the results, the rationale of the choice is typically practical. Other choices reflect preferences and values, which should be openly communicated in a scientific context. Examples of such key questions related to food systems are: should we find ways to fulfil food or resource demand, or instead control that demand? Should we minimise water use, or its impacts? Or should we instead minimise the change from the state with natural vegetation? Whole resource-saving strategies, optimisation objectives and constraints represent different priorities which modellers and model users should

be sensitive about and communicate with their results. Below, I will discuss the choices taken to answer these questions using the Cultural Theory framework, emphasising how they relate to the models in this thesis. The choices are grouped into five categories, as summarised in Table 2.

Table 2: Choices in food systems modelling and possible interpretations in terms of Cultural Theory worldviews, as discussed in main text

Modelling choice	Used in Publications	Worldview (see Section 4.3)	Key rationale for chosen interpretation(s)
<i>Resource saving strategies</i>			
Diet change, decreasing livestock content in human food	I-IV	Egalitarian	Manage population's demand
Optimising system to maximise people fed, yield gap closure, cropland expansion, desalination	IV	Hierarchist	Controlled management using all resources but no more to meet any demand
Loss and waste reduction	II, III (possible in Aalto Opto-Food)	Hierarchist/ Egalitarian/ Individualist	Respectively where it involves managing supply, demand, and avoiding costs of losses
<i>Water use evaluation</i>			
Water footprint analysis	I-III	Egalitarian	Demand-based, sharing responsibility among consumers for human appropriation causing diversions from natural state
Agrobiological modelling of water use	IV	Hierarchist	Measure and manage system as a whole, including supply side
<i>Water use optimisation objectives</i>			
Minimise ET	IV	Hierarchist/Egalitarian	Respectively, optimising resource use, minimising footprints on vulnerable nature

Minimise difference from natural ET	IV	Egalitarian	Caution in face of vulnerable nature, but focussing on change in hydrological cycle rather than human appropriation
Minimise water stress	IV	Hierarchist/ Individualist	Manage demand, stress as cost
Economics-based models	<i>NA</i>	Individualist	Supply-demand balance, cost determines solution. Produce what is most profitable rather than to prioritise human needs
Trade-off analysis	<i>NA</i>	Hierarchist	Explicitly controlling both objectives to consider and who decides their value
<i>Optimisation constraints</i>			
Fixed water availability constraints	IV	Egalitarian	Manage demand, not resource
No-expansion land use scenarios	IV	Egalitarian	Manage demand, no increase in resource, resource usage is managed
Land use stickiness	IV	Egalitarian	Limiting disturbance to current system
Agent-based land use change modelling	<i>NA</i>	Individualist	Capture the emergent behaviour of individuals
Full reallocation of land use	<i>NA</i>	Hierarchist	Controlling and optimising land use
Imposing food-based dietary guidelines	I-III	Hierarchist	Setting acceptable limits on nutrition
<i>Modelling actions</i>			
Optimisation	I-III (diet), IV (land)	Hierarchist	Determining what should be done
Describing scenarios	I-IV	Egalitarian	Providing input for others to consider

Resource-saving strategies

Diet change as a resource-saving measure is a population-level (group) strategy requiring establishment of new norms. In general, it has a strong egalitarian character. It relies on managing food demand, and the management strategy is typically designed to restrict resource use. The egalitarian view is also present in how diet change implementation has been attempted so far by peer pressure and education, respecting individual decisions (Publication **I**), rather than enforcement of a collective decision (grid). Nevertheless, depending on how the objectives are framed, there can be some hierarchist connotations as well. If the preserved resources are to be used for feeding a larger population as Publication **I** and **III** do also emphasise, the question becomes that of allocating resources rather than preserving them.

While diet change as a public or personal health strategy is outside the scope of this dissertation except as an optimisation constraint in Publication **I**, it is often presented together with resource conservation and food security studies (Blas et al., 2019; Gordon et al., 2017; Willett et al., 2019). Coupling multiple justifications suggesting the same consumer action appeals to the egalitarian as a way of building peer pressure, more effectively targeting groups whose habits are already vulnerable to change (Verplanken and Wood, 2006).

Loss and waste reduction (Publications **II-IV**) has both supply and demand side implementations throughout the food supply chain. Thus, it appeals to both egalitarian and hierarchist views. The supply side action, improvements of efficiency from agricultural production to distribution, can be seen as hierarchist supply management, whereas consumer waste reduction is akin to diet change in cutting down the burden on resources by managing demand. The individualist, typically represented by a farmer, industrialist or a trade actor, might also want to reduce losses, but rather to increase profitability or to decrease cost than to save natural resources. This distinction is important and may play a fundamental role in choosing food system modelling techniques. Optimisation of resource efficiency has a relatively direct connection with the views of nature specified by the Cultural Theory. Maximisation of profit, as might be suggested by an economic model, does not directly correlate with minimisation of resource use or environmental impacts in most current societies. This is because some of the resources and impacts are treated as externalities, or their allocation is justified by other utilitarian arguments. Also, maximisation of profit may be an insufficient description of the goals of some actors in the supply side. For example, smallholder farmers may be willing to trade profit for security in their management choices.

Yield gap closure as a food supply side measure (Publication **III**) is an obvious choice for the hierarchist. Even when agriculture is seen as one of the consumers of allocated water in a management scheme, the water use is strictly controlled by the hierarchist yield-increasing practices and enabling technologies. The egalitarian would view yield gap closure as an unnecessary diversion from the natural state, but could be persuaded to accept the approach if it would allow reverting some of the cropland to natural ecosystems.

Other measures that are not covered by the publications in this thesis include for example cropland expansion and seawater desalination. These are both choices for the hierarchist. Both aim at increasing resource availability, the former

primarily by human appropriation, and the latter by application of new technology. The egalitarian would object to cropland expansion as a breach of natural ecosystems, and would consider the desalination approach unnecessary use of energy which could be avoided by reducing food demand. The individualist would carefully weigh the cost, and probably find the current price of desalinated water too high for any but the most valuable and water-efficient food products. Cropland expansion may be profitable in productive areas, especially where many of the environmental impacts are still left as externalities.

Water use evaluation

The method used to analyse water consumption may in itself reflect a specific worldview. Water footprint analysis (Publication **I-III**) imposes an egalitarian signature wherever it is used, as it assumes that all decisions regarding supply side measures have already been made, with the possible exception of the country of origin of the products. All management options available are related to consumption, and all effects are determined by it. This does not preclude interpreting the results through other worldviews, though, as noted above with diet change.

Modelling food water use with an agrobiological model, such as LPJmL (Publication **IV**), allows the analysis of alternative ways to fulfil food demand. Alternative land use models can be spatially more flexible than the area units used in existing water footprint assessments, and different agricultural management options may be explored. This allows a more direct analysis of supply side management, an inherent element of the hierarchist worldview. Again, the interpretation of the results can shift the emphasis towards other views.

It should be noted that there is a scale effect involved. Currently published water footprints tend to be used at relatively large spatial scales, and for consumers rather than producers. Water footprints can also be calculated at farm level for producers, in which case it is more amenable to a hierarchist use. Publications **I-III** used country level consumer footprints, putting them firmly in the egalitarian camp.

Water use optimisation objectives

All choices within one modelling framework are not necessarily made in support of the same worldview. For example, the Aalto OptoFood model provides three alternative objective functions: minimise ET, minimise difference between crop and natural ET, and minimise water stress. Each function represents a pursuit of a food system state that is optimal from some perspective, while none of them is objectively better than the others.

Reducing evapotranspiration is an approach analogous to, although not identical with reducing blue and green water footprints. The differences stem from WFP assessment including other types of water use that goes into producing food in addition to what is required to grow the crops and feed. The latter is by far the largest component, and the one that is most relevant in strategies aiming at sustainable use of agricultural resources.

Controlling evapotranspiration is a hierarchist objective, but the result can be seen as useful from several standpoints. If unused by humans, the saved water can be left to the needs of natural ecosystems – a central goal to the egalitarian. If

reduced evapotranspiration increases blue water availability to other sectors, the hierarchist could see opportunities in resource allocation – potentially in favour of an individualist willing to make profit.

Minimising the difference between the ET of crops and the ET of the area's natural vegetation is an egalitarian objective, although from a wider perspective. Instead of looking at the local water availability to natural ecosystems or local biodiversity effects of natural vs. agricultural land, the focus is on regional and potentially even global effects on the hydrological cycle (Ent et al., 2010). Publication I took a similarly egalitarian perspective in minimising change from current diets. Interestingly, it becomes a hierarchist objective if we establish limits on change, as in the planetary boundary perspective (Rockström et al., 2009). Rather than avoiding disturbing the system, the hierarchist tries to control the disturbance to keep the system stable. Suppose an egalitarian points out that concrete typically has a lower ET than natural vegetation (see Publication IV), so it is more important to share our water use than to minimise it. A hierarchist would respond that the aim is to minimise within constraints – there is more to go around if everybody uses water more efficiently, including the environment.

Water stress stems from the anthropogenic water demand exceeding easily available water resources. Assuming fixed water availability per hydrological basin and a fixed human population, water stress can be reduced by exploiting water resources outside the stressed basins by importing food (Porkka, 2016). Optimally, and as modelled by Aalto OptoFood, this is an allocation approach, preferred by the hierarchist worldview: simply move water-thirsty production out of the stressed basins to water-rich areas. In practice, there is no authority that could control food production in such a way. The solution has been trade, much to the joy and benefit of the individualist. A free market economist would argue that free trade without externalities and other market distortions would result in optimal distributions of water use and stress that reflect the value of water.

However, costs of water are not fully represented by water stress, and a more direct representation of the individualist worldview would have been an economic model. Models based on the biophysical inputs and socio-economic data, such as the global land use allocation model MAGPIE (Dietrich et al., 2018) allocate resources in order to supply food and other outputs, like bioenergy while minimising total cost of production. MAGPIE in particular takes a social planner approach, and might therefore be considered to have hierarchist tendencies, but similar results would be expected from a computable general equilibrium model capturing supply and demand responses of different sectors/individuals. In any case, using economics-based models does not preclude using constraints stemming from other perspectives.

An advantage of detaching the optimisation objective from the physical resources into a common abstraction like money is that multiple goals can be represented in common units, although this possibility does not guarantee that competing goals will be satisfactorily achieved. An example of such situation is bioeconomy, the use of crops as feedstocks in industrial processes for producing energy, biofuels, consumer products etc. Either the use of the crops themselves, or the resources used to produce them, may compete with food production, much

like livestock feed and human food supply chains are partly interlinked. Our results suggest that if enhancing food security is a direct goal, the livestock sector may to some extent increase food supply. However, without careful consideration of possibilities offered by circular economy and other ways of recycling resources, feedstocks to industrial bioeconomy are permanently removed from the FSC, thus representing a real competition with human food. The hierarchist would therefore prefer to analyse the problem as a trade-off (McCornick et al., 2008) and to treat the final choice as a political decision.

Optimisation constraints

Much like the optimisation objectives, definitions of many of the constraints are affected by the views and values of the modeller. A very obvious example, extensively discussed by Hoekstra (2000) and a determining factor between the worldviews is availability of water, or more generally, natural resources. The hierarchist carefully allocates water to optimally support different important uses and technology to appropriate a larger base of resources. The egalitarian agrees that the resource is scarce, but rather manages the demand than tries to increase supply. Aalto OptoFood assumes no expansion in agricultural land, water availability and irrigation equipment, adhering here to the egalitarian view. The stickiness constraints that keep unchanging production in place also appeal to the egalitarian, who wants to work in the small scale and minimise and avoid large technological solutions. Both the hierarchist and the individualist would choose a less restrictive scheme that would allow for more options to manage the supply and potentially to increase profitability by rearranging existing cropland. The individualist sees scarcity as a driver of price, which in itself controls the use and in turn also brings in investments to improving availability – and obviously, if given a choice, would use an economic model to describe the system.

In Publications **I–III** optimisation to create diets is constrained by nutrition guidelines. This is a hierarchist approach, based on population level information. An individualist might want a diet tailored to individual characteristics (e.g. with an agent-based model), and an egalitarian would take an exploratory approach, describing the impacts of diet for each individual to consider. A fatalist would stick to projecting current diet change without trying to change it.

Modelling actions

Regardless of what resource-saving strategies we consider, how we evaluate and optimise water use, and what constraints we place on change, ultimately the question is about action. The optimisation approach in Publication **IV** is at the face of it a hierarchist approach, in reallocating production globally. In focusing on an ideal case, it falls short, however, of providing a tool for policy makers to use.

Instead, it aims to tell a story – to present a scenario that helps people to make sense of the world around them and decide themselves whether they want to take an action. There is an egalitarian aspect in that the researchers recognise the need for humanity to face the problem collectively, but to make their own decisions. Everyone needs to be included regardless of their worldview. Individualists can use the scenarios for their own gain, hierarchists for collective management, and fatalists are free to ignore it or prepare to adapt to inevitable change.

Egalitarianism is in fact the dominant approach across all the papers in considering possible future actions. Publication **I** presents diet scenarios that respect cultural values. Publication **II** adds scenarios of food loss reduction, and Publication **III** scenarios of yield gap closure. Publication **IV** treats the optimisation objective and level of diet change as scenarios. Even though, along the way, other perspectives have crept in for practical reasons, my hope is that this work is in the end as inclusive as possible – that no matter their world view, everyone reading it will be able to find value and better understand the issues associated with changing food systems, and diet change in particular.

5 Conclusions

Food production is by far the largest sector of human-induced water consumption. It needs to have a central role in any comprehensive strategies aiming for sustainable water use. Consumption and production side measures such as diet change, food loss and waste reduction as well as efficient cropland allocation have all been shown to be important building blocks of such strategies. This thesis offers new insight to their effects and novel ways to examine them as part of future food system scenarios.

A novel method was developed to analyse replacement of livestock-based foods in human diet with plant-based alternatives (*New scientific finding 1*). It allows parameterisation of diet change scenarios and inclusion of a dietary recommendation to the replacement procedure. A previously developed, detailed food loss reduction model was considerably improved by addition of the handling of livestock-based foods as well as green water loss analysis. The diet and loss models were combined to investigate interaction between scenarios utilising these measures to reduce food water use. The effect of these measures was found to be approximately multiplicative, but with a small synergistic effect demonstrating the need to explicitly analyse dynamics in food systems potentially affecting resource-saving strategies (*New scientific finding 2*).

Combined with yield gap closure, diet change and food loss reduction were found to be sufficient measures to fulfil the global food demand in 2050, assuming the regional differences in production could be smoothed out by trade. Analysing more food supply chain elements together further emphasised the need for tighter integration in order to handle the system dynamics, especially between crop and livestock production (*New scientific finding 3*). This led to development of an integrated food system model that takes into account how effects of changes in food consumption or supply chains propagate down to agricultural production (*New scientific finding 4*).

Animal-based food, especially meat, consumes considerably more water to fulfil the same nutritional goals achievable by directly utilising plant-based proteins in human diets, when the production of each of the foodstuffs is individually analysed. Reducing consumption of animal-based products, however, does not always reduce total water consumption in the food system. The livestock sector may use feed products that cannot be directly consumed as food, and land that is not suitable for crop production the replacing protein crops may be sourced from inefficient production and supply chains. How these effects are modelled depends to a certain extent on value-based decisions made by the authors as well as users

of the modelling software. Clearly communicating the values affecting modelling choices is crucially important for the readers and researchers using results affected by such decisions as basis of further analysis to understand the embedded limitations and possible biases.

While the calculated results are inconclusive as to whether some meat consumption is better than none in order to minimise water use or its impacts, or to maximise food supply, they clearly suggest that most of the benefits from the diet change towards plant-based protein sources can be achieved even with a moderate level of implementation (*New scientific finding 5*). Hopefully, this can be seen as an encouragement towards resource-efficient policies regarding food production as well as sustainable food consumption habits. It is easy to dismiss strategies that require extensive changes to human behaviour to achieve measurable effects. Diet change may be different in this respect: the first steps towards sustainable eating habits may be the most important.

References

- Affognon, H., Mutungi, C., Sanginga, P., Borgemeister, C., 2015. Unpacking Postharvest Losses in Sub-Saharan Africa: A Meta-Analysis. *World Dev.* 66, 49–68. <https://doi.org/10.1016/j.worlddev.2014.08.002>
- Aleksandrowicz, L., Green, R., Joy, E.J.M., Smith, P., Haines, A., 2016. The Impacts of Dietary Change on Greenhouse Gas Emissions, Land Use, Water Use, and Health: A Systematic Review. *PLOS ONE* 11, e0165797. <https://doi.org/10.1371/journal.pone.0165797>
- Allievi, F., Vinnari, M., Luukkanen, J., 2015. Meat consumption and production – analysis of efficiency, sufficiency and consistency of global trends. *J. Clean. Prod.* 92, 142–151. <https://doi.org/10.1016/j.jclepro.2014.12.075>
- Baroni, L., Cenci, L., Tettamanti, M., Berati, M., 2006. Evaluating the environmental impact of various dietary patterns combined with different food production systems. *Eur. J. Clin. Nutr.* 61, 279–286. <https://doi.org/10.1038/sj.ejcn.1602522>
- Berger, M., Finkbeiner, M., 2013. Methodological Challenges in Volumetric and Impact-Oriented Water Footprints. *J. Ind. Ecol.* 17, 79–89. <https://doi.org/10.1111/j.1530-9290.2012.00495.x>
- Bitzer, M.J., Riddell, D.O., 1984. Corn Grades and Feed Value 3.
- Blas, A., Garrido, A., Unver, O., Willaarts, B., 2019. A comparison of the Mediterranean diet and current food consumption patterns in Spain from a nutritional and water perspective. *Sci. Total Environ.* 664, 1020–1029. <https://doi.org/10.1016/j.scitotenv.2019.02.111>
- Bondeau, A., Smith, P.C., Zaehle, S., Schaphoff, S., Lucht, W., Cramer, W., Gerten, D., Lotze-Campen, H., Müller, C., Reichstein, M., Smith, B., 2007. Modelling the role of agriculture for the 20th century global terrestrial carbon balance. *Glob. Change Biol.* 13, 679–706. <https://doi.org/10.1111/j.1365-2486.2006.01305.x>
- Bosmans, J.H.C., Beek, L.P.H. van, Sutanudjaja, E.H., Bierkens, M.F.P., 2017. Hydrological impacts of global land cover change and human water use. *Hydrol. Earth Syst. Sci.* 21, 5603–5626. <https://doi.org/10.5194/hess-21-5603-2017>
- Brunson, M., 1992. Professional Bias, Public Perspectives, and Communication Pitfalls for Natural Resource Managers. *Rangelands* 14, 292–295.
- Clark, M., Tilman, D., 2017. Comparative analysis of environmental impacts of agricultural production systems, agricultural input efficiency, and food choice. *Environ. Res. Lett.* 12, 064016. <https://doi.org/10.1088/1748-9326/aa6cd5>
- Clay, J., 2019. WWF Responds to EAT-Lancet Report on Global Diets. Trust Food. URL <https://www.trustinfood.com/wwf-responds-to-eat-lancet-report-on-global-diets/> (accessed 4.1.19).
- Davis, K.F., D’Odorico, P., 2015. Livestock intensification and the influence of dietary change: A calorie-based assessment of competition for crop production. *Sci. Total Environ.* 538, 817–823. <https://doi.org/10.1016/j.scitotenv.2015.08.126>

- Davis, K.F., Rulli, M.C., Seveso, A., D'Odorico, P., 2017. Increased food production and reduced water use through optimized crop distribution. *Nat. Geosci.* 10, 919. <https://doi.org/10.1038/s41561-017-0004-5>
- Dietrich, J., Humpenöder, F., bodirsky, Karstens, K., mishkos, Abhijeet Mishra, David Klein, Xiaoxi Wang, weindl, Geanderson Ambrósio, Ewerton Araujo, FelicitasBeier, David Chen, 2018. MAGPIE - An Open Source land-use modeling framework - Version 4.0. <https://doi.org/10.5281/zenodo.1446436>
- Doremus, H., 2008. Scientific and Political Integrity in Environmental Policy. *Tex. Law Rev. Austin* 86, 1601–1653.
- Douglas, M., 1970. *Natural symbols: explorations in cosmology.* Mary Douglas. Barrie and Rockliff, London.
- Douglas, M., Wildavsky, A., 1983. *Risk and Culture: An Essay on the Selection of Technological and Environmental Dangers.* University of California Press, Berkeley, Calif.
- Ent, R.J. van der, Savenije, H.H.G., Schaeffli, B., Steele-Dunne, S.C., 2010. Origin and fate of atmospheric moisture over continents. *Water Resour. Res.* 46. <https://doi.org/10.1029/2010WR009127>
- EU Retail Forum, 2014. Water Footprint Issue Paper.
- Evenson, R.E., Gollin, D., 2003. Assessing the Impact of the Green Revolution, 1960 to 2000. *Science* 300, 758–762. <https://doi.org/10.1126/science.1078710>
- Falkenmark, M., Lannerstad, M., 2010. Food security in water-short countries – Coping with carrying capacity overshoot, in: *Re-Thinking Water and Food Security: Fourth Botin Foundation Water Workshop.* CRC Press, pp. 3–22.
- FAO, 2019. FAOSTAT [WWW Document]. FAOSTAT. URL <http://www.fao.org/faostat/en/#data/> (accessed 1.5.19).
- FAO, 2018a. ESS : Food security indicators [WWW Document]. URL <http://www.fao.org/economic/ess/ess-fs/ess-fadata/en/#.XETkBFUzYdU> (accessed 1.20.19).
- FAO, 2018b. Food-based dietary guidelines [WWW Document]. Food Agric. Organ. U. N. URL <http://www.fao.org/nutrition/education/food-dietary-guidelines/home/en/> (accessed 12.29.18).
- FAO (Ed.), 2017. *The future of food and agriculture: trends and challenges.* Food and Agriculture Organization of the United Nations, Rome.
- FAO, 2012. *Coping with water scarcity. An action framework for agriculture and food security* (38). Rome, Italy.
- Fischer, G., Nachtergaele, F.O., Prieler, S., Teixeira, E., Toth, G., van Velthuizen, H.T., Verelst, L., Wiberg, D., 2016. *Global Agro-Ecological Zones (GAEZ v3.0).* IIASA.
- Gephart, J.A., Troell, M., Henriksson, P.J.G., Beveridge, M.C.M., Verdegem, M., Metian, M., Mateos, L.D., Deutsch, L., 2017. The 'seafood gap' in the food-water nexus literature—issues surrounding freshwater use in seafood production chains. *Adv. Water Resour.* 110, 505–514. <https://doi.org/10.1016/j.advwatres.2017.03.025>
- Gerten, D., Heinke, J., Hoff, H., Biemans, H., Fader, M., Waha, K., 2011. Global Water Availability and Requirements for Future Food Production. *J. Hydrometeorol.* 12, 885–899. <https://doi.org/10.1175/2011JHM1328.1>
- Godfray, H.C.J., Beddington, J.R., Crute, I.R., Haddad, L., Lawrence, D., Muir, J.F., Pretty, J., Robinson, S., Thomas, S.M., Toulmin, C., 2010. Food security: the challenge of feeding 9 billion people. *Science* 327, 812–818. <https://doi.org/10.1126/science.1185383>

- Gordon, L.J., Bignet, V., Crona, B., Henriksson, P.J.G., Holt, T.V., Malin Jonell, Lindahl, T., Troell, M., Barthel, S., Deutsch, L., Folke, C., Haider, L.J., Johan Rockström, Queiroz, C., 2017. Rewiring food systems to enhance human health and biosphere stewardship. *Environ. Res. Lett.* 12, 100201. <https://doi.org/10.1088/1748-9326/aa81dc>
- Gustavsson, J., Cederberg, C., Sonesson, U., van Otterdijk, R., Meybeck, A., 2011. Global food losses and food waste: Extent, causes and prevention. Food and agriculture organization of the United Nations (FAO) Rome.
- Harcombe, Z., 2019. The EAT Lancet diet is nutritionally deficient – Zoë Harcombe. URL <http://www.zoeharcombe.com/2019/01/the-eat-lancet-diet-is-nutritionally-deficient/> (accessed 4.1.19).
- Hemler, E.C., Hu, F.B., 2019. Plant-Based Diets for Cardiovascular Disease Prevention: All Plant Foods Are Not Created Equal. *Curr. Atheroscler. Rep.* 21, 18. <https://doi.org/10.1007/s11883-019-0779-5>
- Herrman, T., Kuhl, G., 1997. Grain Grading Standards in Feed Manufacturing (No. MF-2034). Manhattan, Kansas.
- Hoekstra, A.Y., 2009. Human appropriation of natural capital: A comparison of ecological footprint and water footprint analysis. *Ecol. Econ.* 68, 1963–1974. <https://doi.org/10.1016/j.ecolecon.2008.06.021>
- Hoekstra, A.Y., 2000. Appreciation of water: four perspectives. *Water Policy* 1, 605–622. [https://doi.org/10.1016/S1366-7017\(99\)00013-6](https://doi.org/10.1016/S1366-7017(99)00013-6)
- Hoekstra, A.Y., Chapagain, A.K., Aldaya, M.M., Mekonnen, M.M., 2011. The Water Footprint Assessment Manual: Setting the Global Standard. Earthscan, London.
- Hoekstra, A.Y., Hung, P.Q., 2002. Virtual water trade. *Quantif. Virtual Water Flows Nations Relat. Int. Crop Trade Value Water Res. Rep. Ser.* 11, 166.
- Hoff, H., Falkenmark, M., Gerten, D., Gordon, L., Karlberg, L., Rockström, J., 2010. Greening the global water system. *J. Hydrol., Green-Blue Water Initiative (GBI)* 384, 177–186. <https://doi.org/10.1016/j.jhydrol.2009.06.026>
- Huesemann, M.H., 2002. The inherent biases in environmental research and their effects on public policy. *Futures* 34, 621–633. [https://doi.org/10.1016/S0016-3287\(02\)00004-6](https://doi.org/10.1016/S0016-3287(02)00004-6)
- ISO, 2014. ISO 14046:2014 Environmental management -- Water footprint -- Principles, requirements and guidelines.
- Kastner, T., Kastner, M., Nonhebel, S., 2011. Tracing distant environmental impacts of agricultural products from a consumer perspective. *Ecol. Econ.* 70, 1032–1040. <https://doi.org/10.1016/j.ecolecon.2011.01.012>
- Kummu, M., de Moel, H., Porkka, M., Siebert, S., Varis, O., Ward, P.J., 2012. Lost food, wasted resources: global food supply chain losses and their impacts on freshwater, cropland, and fertiliser use. *Sci. Total Environ.* 438, 477–489. <https://doi.org/10.1016/j.scitotenv.2012.08.092>
- Kummu, M., Guillaume, J.H.A., Moel, H. de, Eisner, S., Flörke, M., Porkka, M., Siebert, S., Veldkamp, T.I.E., Ward, P.J., 2016. The world's road to water scarcity: shortage and stress in the 20th century and pathways towards sustainability. *Sci. Rep.* 6, 38495. <https://doi.org/10.1038/srep38495>
- Lin, B., Chen, X., Yao, H., Chen, Y., Liu, M., Gao, L., James, A., 2015. Analyses of landuse change impacts on catchment runoff using different time indicators based on SWAT model. *Ecol. Indic.* 58, 55–63. <https://doi.org/10.1016/j.ecoind.2015.05.031>

- LUKE, 2018. Luonnonvarakeskuksen tilastot [WWW Document]. URL <http://stat.luke.fi/satotilasto> (accessed 1.17.17).
- Marlow, H.J., Hayes, W.K., Soret, S., Carter, R.L., Schwab, E.R., Sabaté, J., 2009. Diet and the environment: does what you eat matter? [WWW Document]. URL <https://academic.oup.com/ajcn/article/89/5/1699S/4596964> (accessed 12.16.13).
- Mausser, W., Klepper, G., Zabel, F., Delzeit, R., Hank, T., Putzenlechner, B., Calzadilla, A., 2015. Global biomass production potentials exceed expected future demand without the need for cropland expansion. *Nat. Commun.* 6, 8946. <https://doi.org/10.1038/ncomms9946>
- McCornick, P.G., Awulachew, S.B., Abebe, M., 2008. Water–food–energy–environment synergies and tradeoffs: major issues and case studies. *Water Policy* 10, 23–36. <https://doi.org/10.2166/wp.2008.050>
- Mekonnen, M.M., Hoekstra, A.Y., 2010a. The green, blue and grey water footprint of crops and derived crop products (No. 47), Value of Water Research Report. Delft, the Netherlands.
- Mekonnen, M.M., Hoekstra, A.Y., 2010b. The green, blue and grey water footprint of farm animals and animal products (No. 48), Value of Water Research Report Series. Unesco-IHE Institute for Water Education, Delft, the Netherlands.
- Milonni, P.W., 1984. Wave-Particle Duality of Light: A Current Perspective, in: Diner, S., Fargue, D., Lochak, G., Selleri, F. (Eds.), *The Wave-Particle Dualism: A Tribute to Louis de Broglie on His 90th Birthday*, Fundamental Theories of Physics. Springer Netherlands, Dordrecht, pp. 27–67. https://doi.org/10.1007/978-94-009-6286-6_2
- Mueller, N.D., Gerber, J.S., Johnston, M., Ray, D.K., Ramankutty, N., Foley, J.A., 2012. Closing yield gaps through nutrient and water management. *Nature* 490, 254–257. <https://doi.org/10.1038/nature11420>
- Munia, H., Guillaume, J.H.A., Mirumachi, N., Porkka, M., Wada, Y., Kummu, M., 2016. Water stress in global transboundary river basins: significance of upstream water use on downstream stress. *Environ. Res. Lett.* 11, 014002. <https://doi.org/10.1088/1748-9326/11/1/014002>
- Nesheim, M.C., Oria, M., Yih, P.T., Committee on a Framework for Assessing the Health, E., Board, F. and N., Resources, B. on A. and N., Medicine, I. of, Council, N.R., 2015. *The U.S. Food and Agriculture System as a Complex Adaptive System*. National Academies Press (US).
- Oki, T., Kanae, S., 2006. Global Hydrological Cycles and World Water Resources. *Science* 313, 1068–1072. <https://doi.org/10.1126/science.1128845>
- Parfitt, J., Barthel, M., Macnaughton, S., 2010. Food waste within food supply chains: quantification and potential for change to 2050. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 365, 3065–3081. <https://doi.org/10.1098/rstb.2010.0126>
- Parkyn, S., 2004. Review of riparian buffer zone effectiveness. Ministry of Agriculture and Forestry.
- Pfister, S., Boulay, A.-M., Berger, M., Hadjikakou, M., Motoshita, M., Hess, T., Riddout, B., Weinzettel, J., Scherer, L., Döll, P., Manzardo, A., Núñez, M., Verones, F., Humbert, S., Buxmann, K., Harding, K., Benini, L., Oki, T., Finkbeiner, M., Henderson, A., 2017. Understanding the LCA and ISO water footprint: A response to Hoekstra (2016) “A critique on the water-scarcity weighted water footprint in LCA.” *Ecol. Indic.* 72, 352–359. <https://doi.org/10.1016/j.ecolind.2016.07.051>

- Pimentel, D., Pimentel, M., 2003. Sustainability of meat-based and plant-based diets and the environment. *Am. J. Clin. Nutr.* 78, 660S–663S.
- Porkka, M., 2016. Securing global food supplies with limited resources: Lessons from the past. <https://doi.org/10.13140/RG.2.2.12653.79841>
- Portmann, F.T., Siebert, S., Döll, P., 2010. MIRCA2000—Global monthly irrigated and rainfed crop areas around the year 2000: A new high-resolution data set for agricultural and hydrological modeling. *Glob. Biogeochem. Cycles* 24. <https://doi.org/10.1029/2008GB003435>
- Renault, D., Wallender, W.W., 2000. Nutritional water productivity and diets. *Agric. Water Manag.* 45, 275–296. [https://doi.org/10.1016/S0378-3774\(99\)00107-9](https://doi.org/10.1016/S0378-3774(99)00107-9)
- Ridoutt, B.G., Pfister, S., 2013. A new water footprint calculation method integrating consumptive and degradative water use into a single stand-alone weighted indicator. *Int. J. Life Cycle Assess.* 18, 204–207. <https://doi.org/10.1007/s11367-012-0458-z>
- Rockström, J., Steffen, W., Noone, K., Persson, Å., Chapin, F.S., Lambin, E.F., Lenton, T.M., Scheffer, M., Folke, C., Schellnhuber, H.J., Nykvist, B., de Wit, C.A., Hughes, T., van der Leeuw, S., Rodhe, H., Sörlin, S., Snyder, P.K., Costanza, R., Svedin, U., Falkenmark, M., Karlberg, L., Corell, R.W., Fabry, V.J., Hansen, J., Walker, B., Liverman, D., Richardson, K., Crutzen, P., Foley, J.A., 2009. A safe operating space for humanity. *Nature* 461, 472–475. <https://doi.org/10.1038/461472a>
- Röös, E., Patel, M., Spångberg, J., Carlsson, G., Rydhmer, L., 2016. Limiting livestock production to pasture and by-products in a search for sustainable diets. *Food Policy* 58, 1–13. <https://doi.org/10.1016/j.foodpol.2015.10.008>
- Rosa, L., Rulli, M.C., Davis, K.F., Chiarelli, D.D., Passera, C., Paolo D’Odorico, 2018. Closing the yield gap while ensuring water sustainability. *Environ. Res. Lett.* 13, 104002. <https://doi.org/10.1088/1748-9326/aadeef>
- Santos, C.A.G., Silva, R.M. da, Silva, A.M., Brasil Neto, R.M., 2017. Estimation of evapotranspiration for different land covers in a Brazilian semi-arid region: A case study of the Brígida River basin, Brazil. *J. South Am. Earth Sci.* 74, 54–66. <https://doi.org/10.1016/j.jsames.2017.01.002>
- Schwarz, M., Thompson, M., 1990. *Divided We Stand: Re-Defining Politics, Technology and Social Choice*. University of Pennsylvania Press.
- Teicholz, N., 2019. Majority of EAT-Lancet Authors (>80%) Favored Vegan/Vegetarian Diets. Nina Teicholz. URL <https://ninateicholz.com/majority-of-eat-authors-vegan-vegetarian/> (accessed 4.1.19).
- Tilman, D., Balzer, C., Hill, J., Befort, B.L., 2011. Global food demand and the sustainable intensification of agriculture. *Proc. Natl. Acad. Sci.* 108, 20260–20264. <https://doi.org/10.1073/pnas.1116437108>
- Tiu Wright, L., Kwok, P.M.H., Nancarrow, C., 2001. Food taste preferences and cultural influences on consumption. *Br. Food J.* 103, 348–357. <https://doi.org/10.1108/00070700110396321>
- van der Linden, S., 2016. *A Conceptual Critique of the Cultural Cognition Thesis*. *Sci. Commun.* 38, 128–138. <https://doi.org/10.1177/1075547015614970>
- van Ittersum, M.K., Cassman, K.G., Grassini, P., Wolf, J., Tittonell, P., Hochman, Z., 2013. Yield gap analysis with local to global relevance—A review. *Field Crops Res., Crop Yield Gap Analysis – Rationale, Methods and Applications* 143, 4–17. <https://doi.org/10.1016/j.fcr.2012.09.009>

- Vanham, D., Comero, S., Gawlik, B.M., Bidoglio, G., 2018. The water footprint of different diets within European sub-national geographical entities. *Nat. Sustain.* 1, 518. <https://doi.org/10.1038/s41893-018-0133-x>
- Vanham, D., Hoekstra, A.Y., Bidoglio, G., 2013. Potential water saving through changes in European diets. *Environ. Int.* 61, 45–56. <https://doi.org/10.1016/j.envint.2013.09.011>
- Verplanken, B., Wood, W., 2006. Interventions to Break and Create Consumer Habits. *J. Public Policy Mark.* 25, 90–103. <https://doi.org/10.1509/jppm.25.1.90>
- Wada, Y., Gleeson, T., Esnault, L., 2014. Wedge approach to water stress. *Nat. Geosci.* 7, 615–617. <https://doi.org/10.1038/ngeo2241>
- Wang, S., Huang, J., Li, J., Rivera, A., McKenney, D.W., Sheffield, J., 2014. Assessment of water budget for sixteen large drainage basins in Canada. *J. Hydrol.* 512, 1–15. <https://doi.org/10.1016/j.jhydrol.2014.02.058>
- WHO, 2003. Diet, nutrition and the prevention of chronic diseases [WWW Document]. WHO. URL <http://www.who.int/dietphysicalactivity/publications/trs916/en/> (accessed 7.23.13).
- Wichelns, D., 2015. Virtual water and water footprints do not provide helpful insight regarding international trade or water scarcity. *Ecol. Indic.* 52, 277–283. <https://doi.org/10.1016/j.ecolind.2014.12.013>
- Willett, W., Rockström, J., Loken, B., Springmann, M., Lang, T., Vermeulen, S., Garnett, T., Tilman, D., DeClerck, F., Wood, A., Jonell, M., Clark, M., Gordon, L.J., Fanzo, J., Hawkes, C., Zurayk, R., Rivera, J.A., De Vries, W., Majele Sibanda, L., Afshin, A., Chaudhary, A., Herrero, M., Agustina, R., Branca, F., Lartey, A., Fan, S., Crona, B., Fox, E., Bignet, V., Troell, M., Lindahl, T., Singh, S., Cornell, S.E., Srinath Reddy, K., Narain, S., Nishtar, S., Murray, C.J.L., 2019. Food in the Anthropocene: the EAT–Lancet Commission on healthy diets from sustainable food systems. *The Lancet* 393, 447–492. [https://doi.org/10.1016/S0140-6736\(18\)31788-4](https://doi.org/10.1016/S0140-6736(18)31788-4)
- Wise, T.A., 2013. Can We Feed the World in 2050? A Scoping Paper to Assess the Evidence (Working Paper No. 13– 04). GLOBAL DEVELOPMENT AND ENVIRONMENT INSTITUTE.
- Yang, H., Pfister, S., Bhaduri, A., 2013. Accounting for a scarce resource: virtual water and water footprint in the global water system. *Curr. Opin. Environ. Sustain., Aquatic and marine systems* 5, 599–606. <https://doi.org/10.1016/j.cosust.2013.10.003>

ISBN 978-952-60-8672-9 (printed)
ISBN 978-952-60-8673-6 (pdf)
ISSN 1799-4934 (printed)
ISSN 1799-4942 (pdf)

Aalto University
School of Engineering
Department of Built Environment
www.aalto.fi

**BUSINESS +
ECONOMY**

**ART +
DESIGN +
ARCHITECTURE**

**SCIENCE +
TECHNOLOGY**

CROSSOVER

**DOCTORAL
DISSERTATIONS**