Doubly nonlinear parabolic equations

Matias Vestberg

Doubly nonlinear parabolic equations

Matias Vestberg

The public defense on 15th May 2020 at 12:00 will be organized via remote technology.

Link: <https://aalto.zoom.us/j/66291986278>

Zoom Quick Guide: <https://www.aalto.fi/en/services/zoom-quick-guide>

A doctoral dissertation completed for the degree of Doctor of Science (Technology) to be defended, with the permission of the Aalto University School of Science, at an examination held at the lecture hall M1 of the school, and also via remote technology, on 15 May 2020 at 12.

**Aalto University
School of Science
Department of Mathematics and Systems Analysis
Nonlinear PDE research group**

Supervising professor

Professor Juha Kinnunen, Aalto University

Thesis advisor

Professor Juha Kinnunen, Aalto University

Preliminary examiners

Professor Simona Fornaro, University of Pavia, Italy

Professor Erik Lindgren, Uppsala University, Sweden

Opponent

Professor Ugo Gianazza, University of Pavia, Italy

Aalto University publication series

DOCTORAL DISSERTATIONS 68/2020

© 2020 Matias Vestberg

ISBN 978-952-60-3862-9 (printed)

ISBN 978-952-60-3863-6 (pdf)

ISSN 1799-4934 (printed)

ISSN 1799-4942 (pdf)

<http://urn.fi/URN:ISBN:978-952-60-3863-6>

Unigrafia Oy

Helsinki 2020

Finland

Author

Matias Vestberg

Name of the doctoral dissertation

Doubly nonlinear parabolic equations

Publisher School of Science**Unit** Department of Mathematics and Systems Analysis**Series** Aalto University publication series DOCTORAL DISSERTATIONS 68/2020**Field of research** Mathematics**Manuscript submitted** 5 February 2020**Date of the defence** 15 May 2020**Permission for public defence granted (date)** 14 April 2020**Language** English☐ **Monograph**☒ **Article dissertation**☐ **Essay dissertation****Abstract**

This thesis is devoted to the study of the existence and regularity of weak solutions to some doubly nonlinear parabolic equations. Such equations are nonlinear both with respect to the solution and its gradient, and are thus more difficult to handle than the parabolic p-Laplace equation and the porous medium equation, which are special cases of the equations considered here. Much of the work in this thesis concerns the so-called diffusive shallow medium equation, which presents an additional technical difficulty due to the given profile function appearing in the equation.

For the diffusive shallow medium equation we use a De Giorgi type iteration to prove that weak solutions are locally bounded. We show that bounded weak solutions are locally Hölder continuous using the method of intrinsic scaling introduced originally by DiBenedetto.

We prove the existence of a bounded solution to a Cauchy-Dirichlet problem associated with the diffusive shallow medium equation by introducing regularized problems for which the existence of solutions is easy to prove by means of Galerkin's method. In order to show that these solutions converge to a solution of the original problem, we have proved some new compactness results which might be useful also for other nonlinear equations.

There are various definitions for weak solutions appearing in the literature, and in this thesis we have devoted special attention to showing that the adopted definitions allow rigorous proofs of the existence and regularity results without any additional assumptions. For the diffusive shallow medium equation, a definition which is natural in this sense has not previously been used.

For doubly singular parabolic equations we have proved a range of regularity results such as local boundedness, an integral Harnack inequality, local Hölder continuity, expansion of positivity and a Harnack inequality in properly scaled space-time cylinders.

Keywords doubly nonlinear equation, regularity, Hölder continuity, local boundedness, existence, weak solution, Harnack estimate

ISBN (printed) 978-952-60-3862-9**ISBN (pdf)** 978-952-60-3863-6**ISSN (printed)** 1799-4934**ISSN (pdf)** 1799-4942**Location of publisher** Helsinki**Location of printing** Helsinki **Year** 2020**Pages** 181**urn** <http://urn.fi/URN:ISBN:978-952-60-3863-6>

Författare

Matias Vestberg

Doktorsavhandlingens titel

Dubbelt icke-linjära paraboliska ekvationer

Utgivare Högskolan för teknikvetenskaper**Enhet** Institutionen för matematik och systemanalys**Seriens namn** Aalto University publication series DOCTORAL DISSERTATIONS 68/2020**Forskningsområde** Matematik**Inlämningsdatum för manuskript** 05.02.2020**Datum för disputation** 15.05.2020**Beviljande av disputationstillstånd (datum)** 14.04.2020**Språk** Engelska☐ **Monografi**☒ **Artikelavhandling**☐ **Essäavhandling****Sammandrag**

Denna avhandling behandlar existens- och regularitetsfrågor för svaga lösningar till dubbelt icke-linjära paraboliska ekvationer. Dessa ekvationer är icke-linjära både med avseende på lösningen samt dess gradient, och är därmed svårare att hantera än den paraboliska p -Laplace ekvationen och ekvationen för porösa medier, som är specialfall av de ekvationer som betraktas här. En stor del av avhandlingen rör den s.k. diffusionsekvationen för grunda medier, som uppvisar en ytterligare teknisk svårighet p.g.a. den i ekvationen förekommande profilkfunktionen.

Vi använder De Giorgis iterationsmetod för att visa att svaga lösningar till diffusionsekvationen för grunda medier är lokalt begränsade. Vi bevisar att begränsade lösningar är lokalt Hölder-kontinuerliga m.h.a. den av DiBenedetto införda metoden som baserar sig på användningen av en intrinsisk skala.

Vi bevisar existensen av en lösning på ett Cauchy-Dirichlet-problem relaterat till diffusions-ekvationen för grunda medier genom att introducera ett regulariserat problem för vilket lösningar lätt kan hittas m.h.a. Galerkins metod. För att visa att dessa lösningar konvergerar mot en lösning på det ursprungliga problemet har vi bevisat några nya kompakthetsresultat som också kan vara användbara för andra icke-linjära ekvationer.

Det finns ett flertal definitioner på svaga lösningar i litteraturen, och i denna avhandling har vi fäst speciell uppmärksamhet vid att visa att vår definition möjliggör rigorösa bevis för existens- och regularitetsresultat utan extra antaganden. En i detta avseende naturlig definition av lösningar till diffusionsekvationen för grunda medier har inte använts tidigare.

För dubbelt singulära paraboliska ekvationer har vi visat ett antal regularitetsresultat såsom lokal begränsning, en integralform av Harnacks olikhet, lokal Hölderkontinuitet, expansion av positivitet samt en Harnack-olikhet för rum-tidscyldrar i lämplig skala.

Nyckelord dubbelt icke-linjär ekvation, regularitet, Hölderkontinuitet, lokal begränsning, existens, svag lösning, Harnack-estimat

ISBN (tryckt) 978-952-60-3862-9**ISBN (pdf)** 978-952-60-3863-6**ISSN (tryckt)** 1799-4934**ISSN (pdf)** 1799-4942**Utgivningsort** Helsingfors**Tryckort** Helsingfors**År** 2020**Sidantal** 181**urn** <http://urn.fi/URN:ISBN:978-952-60-3863-6>

Preface

The research for this doctoral thesis began in the Spring of 2016, when professor Juha Kinnunen brought my attention to a nonlinear model for shallow water dynamics, which had thus far received little attention in the mathematics community. This was to become one the main topics of the thesis, and I wish to thank professor Kinnunen both for the initial suggestion to study this equation, and for his later guidance as my advisor.

I want to thank my co-authors Verena Bögelein, Nicolas Dietrich, Thomas Singer and Vincenzo Vespri for the opportunity to work together, and for their significant contributions to the articles that make up this thesis. Moreover, I am grateful to Simona Fornaro and Erik Lindgren for carrying out the preliminary examination of the thesis and to Ugo Gianazza for agreeing to act as my opponent in the public examination.

For the financial support, I am indebted to the Academy of Finland and the Vilho, Yrjö and Kalle Väisälä Foundation. Furthermore, I would like to thank my colleagues at the department and especially the "coffee gang" for many rewarding and entertaining conversations.

Finally, I wish to thank my family and my friends.

Borgå, April 15, 2020,

Matias Vestberg

Contents

Preface	1
Contents	3
List of Publications	5
Author's Contribution	7
1. Introduction	9
2. The shallow medium equation	11
2.1 Weak solutions	12
2.2 Tools and notation	13
2.2.1 Estimates for Sobolev functions	14
2.2.2 Time mollification	16
3. Regularity properties	19
3.1 Continuity in time	19
3.2 Mollified weak formulation	20
3.3 Energy estimates	20
3.4 Local boundedness	22
3.5 Local Hölder continuity	23
3.5.1 Scaling and De Giorgi type lemma	24
3.5.2 The two alternatives	25
3.5.3 Iterating the reduction of the oscillation	27
4. Existence	29
4.1 Regularization of the problem	30
4.2 Properties of the approximating solutions	31
4.3 Compactness and convergence results	31
4.4 Existence	32

5. Doubly singular diffusion equations	33
5.1 Energy estimates	34
5.2 L^1 -Harnack inequality	34
5.3 Expansion of positivity	35
5.4 Local boundedness	35
5.5 Local Hölder continuity	36
5.6 Harnack Estimates	37
Bibliography	39
Publications	43

List of Publications

This thesis consists of an overview and of the following publications which are referred to in the text by their Roman numerals.

- I** Thomas Singer and Matias Vestberg. Local boundedness of weak solutions to the Diffusive Wave Approximation of the Shallow Water equations. *Journal of Differential Equations*, Volume 266, Issue 6, pages 3014-3033, March 2019.
- II** Thomas Singer and Matias Vestberg. Local Hölder continuity of weak solutions to a diffusive shallow medium equation. *Nonlinear Analysis*, Volume 185, pages 306-335, August 2019.
- III** Verena Bögelein, Nicholas Dietrich and Matias Vestberg. Existence of solutions to a diffusive shallow medium equation. *Submitted to a journal*, 35 pages, Available at: arXiv:2001.07942, January 2020.
- IV** Vincenzo Vespri and Matias Vestberg. An Extensive Study of the Regularity of Solutions to Doubly Singular equations. *Advances in Calculus of Variations*, 39 pages, Available at: <https://doi.org/10.1515/acv-2019-0102>, March 2020.

Author's Contribution

Publication I: “Local boundedness of weak solutions to the Diffusive Wave Approximation of the Shallow Water equations”

Both co-authors contributed equally to all parts of the paper.

Publication II: “Local Hölder continuity of weak solutions to a diffusive shallow medium equation”

Both co-authors contributed equally to all parts of the paper.

Publication III: “Existence of solutions to a diffusive shallow medium equation”

The author has made significant contributions to all parts of the paper. The proof of the uniform bound for the approximating solutions was developed independently by the author.

Publication IV: “An Extensive Study of the Regularity of Solutions to Doubly Singular equations”

The author has made significant contributions to all parts of the paper. The author developed the proof of local boundedness independently.

1. Introduction

This thesis is devoted to the study of doubly nonlinear parabolic differential equations. The main focus lies on diffusion equations of the form

$$\partial_t u - \nabla \cdot ((u - z)^\alpha |\nabla u|^{p-2} \nabla u) = f, \quad (1.1)$$

where u is the unknown quantity which is diffusing, $z(x)$ is a given profile function and $f(x, t)$ is a known source term. The parameters $\alpha > 0$ and $p > 1$ depend on the particular application. In the parameter range $p \in (1, 2)$, equation (1.1) can be used to model the diffusion of shallow water, whereas in the range $p > 2$, the equation has been used to describe the dynamics of glaciers.

The second type of equations of interest takes the form

$$\partial_t u - \nabla \cdot A(u, x, t, \nabla u^\beta) = f, \quad (1.2)$$

where $\beta > 1$ and A is a vector field satisfying the structure conditions

$$\begin{aligned} A(u, x, t, \xi) \cdot \xi &\geq C_0 |\xi|^p, \\ |A(u, x, t, \xi)| &\leq C_1 |\xi|^{p-1}, \end{aligned}$$

for some positive constants C_0 and C_1 . For certain parameter values we recover important special cases like the porous medium equation and the parabolic p -Laplace equation for which an extensive theory already has been developed. See for example the monographs [11, 51, 15] and the references therein.

There is a connection between (1.1) and (1.2) which becomes apparent in the case $z = 0$. Then, a formal application of the chain rule shows that equations of type (1.1) fall into the category of (1.2) with the choice $\beta = 1 + \frac{\alpha}{p-1}$. However, due to the general form of the vector field A , neither type of equations contains the other.

Prior to this thesis, very little was known about the theoretical aspects of (1.1) in the general case $z \neq 0$, although the equation had been studied numerically, see [44]. Even the natural definition of solutions was

unclear. Formulating a definition which allowed rigorous proofs of existence and regularity results is an important achievement of this thesis. The question of how to define solutions is relevant also for (1.2). Even for the porous medium equation various definitions have been used, and their relations are the subject of recent research, see for example [37].

One of the most important questions concerning any partial differential equation is the existence of solutions. Previously, existence of solutions to (1.1) was studied only for the flat case $z = 0$, see for example [1] which specifically treats the shallow water case, or [47] in which equations of type (1.2) are considered. In Chapter 4, which is a summary of Publication III, we show the existence of solutions to (1.1) satisfying suitable initial and boundary conditions.

Since the existence of solutions is typically established in a weak class, another relevant question is what kind of regularity properties the solutions possess. In this thesis we establish that weak solutions of (1.1) are locally bounded in the slow diffusion case. We follow the approach of De Giorgi, combining energy estimates with a Sobolev embedding. In addition, local Hölder continuity is obtained in the case $p > 2$ using the method of intrinsic scaling due to DiBenedetto [11]. These results can be found in Chapter 3, which is based on Publication I and Publication II.

It is likely that the method of DiBenedetto is applicable also in the more challenging case $p \in (1, 2)$, although the arguments would no doubt become more technical. In this thesis we have opted to lay out the first steps towards an alternative strategy of proving the local Hölder continuity in this range. Namely, we adapt a method involving integral Harnack estimates, which has previously been used to obtain a concise proof of the Hölder continuity of weak solutions to the singular parabolic p -Laplace equation. In Chapter 5, which summarises Publication IV, we extend this method to equation (1.2) in the so-called doubly singular parameter range. In the future, this strategy might be extended to cover also solutions to (1.2) in the singular-degenerate case and ultimately solutions to (1.1) with $p \in (1, 2)$.

Apart from the local Hölder continuity, in Publication IV we also prove other regularity results such as the integral Harnack estimate, local boundedness, expansion of positivity and Harnack estimates on cylinders.

2. The shallow medium equation

In this chapter we explain the general setting and present the precise definition of solutions to (1.1).

Let Ω be an open bounded subset of $\mathbb{R}^n$, where $n \geq 2$, and let $T > 0$. We use the shorthand notation Ω_T for the cylinder $\Omega \times (0, T)$. Physically, Ω represents a region in space and $(0, T)$ is thought of as a time interval. We will study the existence and regularity of weak solutions to the doubly nonlinear parabolic equation

$$\partial_t u - \nabla \cdot ((u - z)^\alpha |\nabla u|^{p-2} \nabla u) = f \quad \text{in } \Omega_T, \quad (2.1)$$

where $z: \Omega \rightarrow \mathbb{R}$ and $f: \Omega_T \rightarrow \mathbb{R}$ are sufficiently regular functions and $\alpha > 0$ and $p > 1$ are fixed parameters. Here ∂_t refers to the partial derivative with respect to the time variable, and the gradient ∇ and divergence $\nabla \cdot$ are taken only with respect to the spatial variables. For $p \in (1, 2)$, the equation has been used to model the diffusion of water in situations such as dam breaks, and floods in vegetated areas, see for example [53, 26, 18]. For physical motivations for this model, see [48, 22], or [1] for a different derivation. In papers focusing on shallow water dynamics it is customary to denote $p = \gamma + 1$. We have followed this convention in Publication I, but in the thesis we will always use p for consistency with the mathematical literature. In the range $p > 2$, the equation is used in models of glacier dynamics, see the classic work [39] or Chapter 2 of [42]. In such applications it is typical to take $p \approx 4$.

Regardless of the application, u represents the height of the water or ice with respect to some fixed ground level, and z represents the height of the land on top of which the water or ice is moving. Thus, we only look for solutions satisfying $u \geq z$. The physical interpretation of the source term f is precipitation, evaporation or infiltration and hence its sign may vary. Even though u only depends on two spatial coordinates in physical applications, such a restriction is not needed in the mathematical theory.

For this reason we allow the spatial dimension n to be arbitrarily large.

2.1 Weak solutions

In the forthcoming chapters we will study so-called weak solutions to (2.1). The concept of weak solutions to partial differential equations was introduced to facilitate existence proofs by dividing the problem into two steps. First, generalize the concept of solution into such a large class that existence is easy to prove. Then show that, in fact, the generalized solution also is a classical solution, or at least somewhat regular so that it might be considered as physically relevant.

For the classical heat equation this strategy works perfectly, as weak solutions turn out to be smooth and hence also solutions in the classical sense, see [17]. A typical feature of nonlinear equations such as the parabolic p -Laplace equation or the porous medium equation is that weak solutions might be less regular, and the relevant question becomes how much regularity one can obtain.

The function space in which existence is initially established depends on the equation. For the parabolic p -Laplace equation

$$\partial_t u - \nabla \cdot (|\nabla u|^{p-2} \nabla u) = 0,$$

it turns out to be a Sobolev space of the form $L^p(0, T; W^{1,p}(\Omega))$, which means that solutions are p -integrable functions with a p -integrable spatial gradient. Note that no regularity with respect to time is assumed.

Comparing the doubly nonlinear equation (1.2) with the parabolic p -Laplace equation we see that in the elliptic part, the quantity u^β replaces u . Therefore, a natural requirement for weak solutions in this case is that $u^\beta \in L^p(0, T; W^{1,p}(\Omega))$.

In order to motivate the natural notion of weak solution in the case of (1.1), we re-write the equation in a form which resembles (1.2). Note first that the vector field appearing in the elliptic part of (2.1) can be re-written as

$$(u - z)^\alpha |\nabla u|^{p-2} \nabla u = |(u - z)^{\frac{\alpha}{p-1}} \nabla u|^{p-2} (u - z)^{\frac{\alpha}{p-1}} \nabla u.$$

Adding and removing the gradient of z and formally applying the chain rule we obtain

$$(u - z)^{\frac{\alpha}{p-1}} \nabla u = (u - z)^{\frac{\alpha}{p-1}} \nabla(u - z) + (u - z)^{\frac{\alpha}{p-1}} \nabla z = \beta^{-1} \nabla v^\beta + v^{\beta-1} \nabla z,$$

where $v = u - z$ and $\beta = 1 + \frac{\alpha}{p-1}$. Thus, (1.1) takes the form

$$\partial_t v - \nabla \cdot [\beta^{1-p} |\nabla v^\beta + \beta v^{\beta-1} \nabla z|^{p-2} (\nabla v^\beta + \beta v^{\beta-1} \nabla z)] = f. \quad (2.2)$$

Note that if z is constant, the equation satisfied by v is exactly of type (1.2). With this formulation, it is natural to assume that v^β has a gradient, rather than v or u . A comparison with the function space of solutions to (1.2), suggests that one should require $v^\beta \in L^p(0, T; W^{1,p}(\Omega))$. To simplify notation, we denote

$$A(v, \nabla v^\beta) = \beta^{1-p} |\nabla v^\beta + \beta v^{\beta-1} \nabla z|^{p-2} (\nabla v^\beta + \beta v^{\beta-1} \nabla z). \quad (2.3)$$

To arrive at the definition of weak solutions we follow the approach which is common to all diffusion equations. Namely, we multiply (2.2) by a smooth compactly supported test function and integrate formally by parts. Thus, a function $u: \Omega_T \rightarrow \mathbb{R}$ is a weak solution to (2.1) if and only if $v = u - z$ is non-negative, $v^\beta \in L^p(0, T; W^{1,p}(\Omega))$ and

$$\iint_{\Omega_T} A(v, \nabla v^\beta) \cdot \nabla \varphi - v \partial_t \varphi \, dx \, dt = \iint_{\Omega_T} f \varphi \, dx \, dt, \quad (2.4)$$

for all $\varphi \in C_0^\infty(\Omega_T)$. Moreover, in this case we say that v is a solution to (2.2).

In order for the right-hand side of (2.4) to be defined, we must at least require that f is locally integrable. Later, when investigating existence and regularity properties we will need to require higher integrability of f . By assumption, ∇v^β is p -integrable and we require ∇z to be sufficiently integrable so that also the term $v^{\beta-1} \nabla z$ is in L^p . A simple calculation using Hölder's inequality shows that it is sufficient to require that ∇z belongs to $L^{\beta p}(\Omega)$.

In principle, somewhat weaker integrability properties of ∇u^β and ∇z would already guarantee that the left-hand side of (2.4) is defined. However, in the regularity theory we will need to use (2.4) with certain nonsmooth test functions, and the higher integrability ensures that this is possible. In general, one can justify a nonsmooth test function if it can be approximated by smooth compactly supported test functions so that also the corresponding integrals in the weak formulation are approximated.

2.2 Tools and notation

In this section we present some tools and well-known mathematical results that have been used extensively in the thesis. First we explain some notation.

By $B_\rho(x)$ we denote the open ball of radius ρ centered at x . We will frequently use the notation $z_o = (x_o, t_o)$ for a point in Ω_T . For such a point we define the space-time cylinder

$$Q_{\rho,\theta}(z_o) := B_\rho(x_o) \times (t_o - \theta, t_o).$$

For $w, v \geq 0$ we define

$$\begin{aligned} \mathfrak{b}[v, w] &:= \frac{1}{\beta+1}(v^{\beta+1} - w^{\beta+1}) - w^\beta(v - w) \\ &= \frac{\beta}{\beta+1}(w^{\beta+1} - v^{\beta+1}) - v(w^\beta - v^\beta). \end{aligned} \quad (2.5)$$

This quantity appears naturally in the energy estimates satisfied by weak solutions to doubly nonlinear equations. Proofs for the properties of $\mathfrak{b}$ which have been used in the publications of this thesis can be found in [3]. As we shall not make explicit use of these properties in the introduction, we do not repeat them here.

Let U be an open subset of $\mathbb{R}^n$ and let $(a, b) \subset \mathbb{R}$ be an interval. By the parabolic boundary of $U \times (a, b)$ we intend the set $(U \times \{a\}) \cup (\partial U \times [a, b])$.

2.2.1 Estimates for Sobolev functions

Here we present some inequalities satisfied by Sobolev functions.

Lemma 1. *Let v be a function in $L^p(0, T; W_0^{1,p}(\Omega))$ and let $p, \kappa \geq 1$. Then the following inequality holds:*

$$\iint_{\Omega_T} |v|^{\kappa p} dx dt \leq c \left[\operatorname{ess\,sup}_{t \in [0, T]} \int_{\Omega} |v(x, t)|^{(\kappa-1)n} dx \right]^{\frac{p}{n}} \iint_{\Omega_T} |\nabla v|^p dx dt, \quad (2.6)$$

where the constant depends only on p, κ, n .

This result looks similar to Proposition 3.1 in Section I.3 of [11], except for the naming of parameters. However, we remark that the exponent $(\kappa - 1)n$ in our case can be any number greater than or equal to zero, whereas in [11] the range of this exponent is restricted to $[1, \infty)$. The proof in [11] is divided into two different cases. We provide an alternative proof which is valid for the whole parameter range.

Proof of Lemma 1. Let $u \in C_0^\infty(\Omega)$. Then, by Hölder's inequality

$$\begin{aligned} \int_{\Omega} |u|^{\kappa p} dx &= \int_{\Omega} |u|^{1+\kappa(p-1)} |u|^{\kappa-1} dx \\ &\leq \left(\int_{\Omega} |u|^{(1+\kappa(p-1))\frac{n}{n-1}} dx \right)^{\frac{n-1}{n}} \left(\int_{\Omega} |u|^{(\kappa-1)n} dx \right)^{\frac{1}{n}}. \end{aligned} \quad (2.7)$$

We want to apply the Gagliardo-Nirenberg inequality for functions in $W_0^{1,1}(\Omega)$ to the first integral on the last line. If $p = 1$ then $1 + \kappa(p -$

1) = 1 and we apply the inequality to u . If $p > 1$ then we observe that $t \mapsto |t|^{1+\kappa(p-1)}$ is continuously differentiable and hence the composition $|u|^{1+\kappa(p-1)}$ is in $C_0^1(\Omega)$ and we can again apply the Gagliardo-Nirenberg inequality. Thus, in both cases we have

$$\begin{aligned} \left(\int_{\Omega} |u|^{(1+\kappa(p-1))\frac{n}{n-1}} dx \right)^{\frac{n-1}{n}} &\leq c \int_{\Omega} |\nabla(|u|^{1+\kappa(p-1)})| dx \\ &= c \int_{\Omega} |u|^{\kappa(p-1)} |\nabla u| dx \\ &\leq c \left(\int_{\Omega} |u|^{\kappa p} dx \right)^{\frac{p-1}{p}} \left(\int_{\Omega} |\nabla u|^p dx \right)^{\frac{1}{p}}, \end{aligned}$$

where the last step follows from Hölder's inequality. Combining the last estimate with (2.7) and denoting $q = (\kappa - 1)n$ we conclude that

$$\int_{\Omega} |u|^{\kappa p} dx \leq c \left(\int_{\Omega} |u|^q dx \right)^{\frac{p}{n}} \int_{\Omega} |\nabla u|^p dx, \quad (2.8)$$

for all $u \in C_0^\infty(\Omega)$. The next step is to extend this inequality to all u in $W_0^{1,p}(\Omega)$. It is sufficient to consider $u \in W_0^{1,p}(\Omega) \cap L^q(\Omega)$ since otherwise the right-hand side is infinite and the inequality holds trivially. Moreover, by dividing the function into its positive and negative parts, we can assume $u \geq 0$. For such a function u , pick $\varphi_j \in C_0^\infty(\Omega)$ so that $\varphi_j \geq 0$ and $\varphi_j \rightarrow u$ in $W^{1,p}(\Omega)$. Define

$$u_j = \min\{\varphi_j, u\}, \quad u_j^\varepsilon = u_j \star \eta_\varepsilon,$$

where η_ε is a standard mollifier. Clearly $u_j \in W^{1,p}(\Omega)$ and $u_j \rightarrow u$ in $W^{1,p}(\Omega)$. Since u_j has compact support contained in Ω we see that u_j^ε belongs to $C_0^\infty(\Omega)$ for small ε , and hence

$$\int_{\Omega} |u_j^\varepsilon|^{\kappa p} dx \leq c \left(\int_{\Omega} |u_j^\varepsilon|^q dx \right)^{\frac{p}{n}} \int_{\Omega} |\nabla u_j^\varepsilon|^p dx. \quad (2.9)$$

We want to pass to the limit $\varepsilon \rightarrow 0$. The second integral on the right-hand side poses no problem since $\nabla u_j^\varepsilon \rightarrow \nabla u_j$ in L^p as $\varepsilon \rightarrow 0$. To treat the other integrals we note that for a suitable sequence (ε_k) converging to zero we may assume that $u_j^{\varepsilon_k}$ converges pointwise a.e. to u_j as $k \rightarrow \infty$. Furthermore, $|u_j^\varepsilon| \leq \|\varphi_j\|_\infty < \infty$. Therefore, the Dominated Convergence Theorem guarantees that the integrals with ε_k in place of ε converge as $k \rightarrow \infty$ and we obtain

$$\begin{aligned} \int_{\Omega} |u_j|^{\kappa p} dx &\leq c \left(\int_{\Omega} |u_j|^q dx \right)^{\frac{p}{n}} \int_{\Omega} |\nabla u_j|^p dx \\ &\leq c \left(\int_{\Omega} |u|^q dx \right)^{\frac{p}{n}} \int_{\Omega} |\nabla u_j|^p dx. \end{aligned}$$

The last estimate follows from the fact that $0 \leq u_j \leq u$. By passing to a subsequence still denoted (u_j) we may assume that u_j converges pointwise a.e. to u . Thus, Fatou's lemma shows that

$$\begin{aligned} \int_{\Omega} |u|^{\kappa p} dx &\leq \liminf_{j \rightarrow \infty} \int_{\Omega} |u_j|^{\kappa p} dx \leq \liminf_{j \rightarrow \infty} \left[c \left(\int_{\Omega} |u|^q dx \right)^{\frac{p}{n}} \int_{\Omega} |\nabla u_j|^p dx \right] \\ &= c \left(\int_{\Omega} |u|^q dx \right)^{\frac{p}{n}} \int_{\Omega} |\nabla u|^p dx. \end{aligned}$$

Thus, we have verified (2.8) for all $u \in W_0^{1,p}(\Omega)$. In order to prove (2.6) we apply (2.8) to $v(\cdot, t)$ at a.e. t to conclude

$$\begin{aligned} \iint_{\Omega_T} |v|^{\kappa p} dx dt &\leq c \int_0^T \left(\int_{\Omega} |v(x, t)|^q dx \right)^{\frac{p}{n}} \int_{\Omega} |\nabla v(x, t)|^p dx dt \\ &\leq c \left[\operatorname{ess\,sup}_{t \in [0, T]} \int_{\Omega} |v(x, t)|^q dx \right]^{\frac{p}{n}} \iint_{\Omega_T} |\nabla v|^p dx dt. \end{aligned}$$

□

In several of the publications we have needed the following isoperimetric inequality which was proved by De Giorgi in his classical work [9].

Lemma 2. *Let $v \in W^{1,1}(B_{\rho}(x_o))$ for some $\rho > 0$ and $x_o \in \mathbb{R}^n$. Let k and l be real numbers such that $k < l$. Then there exists a constant c depending only on n (and thus independent of k, l, v, x_o and ρ) such that for any representative of v , we have*

$$\begin{aligned} (l - k) |\{x \in B_{\rho}(x_o) : v(x) > l\}| |\{x \in B_{\rho}(x_o) : v(x) < k\}| & \quad (2.10) \\ &\leq c \rho^{n+1} \int_{\{k < v < l\} \cap B_{\rho}(x_o)} |\nabla v| dx. \end{aligned}$$

2.2.2 Time mollification

One challenge with the definition of weak solutions is the a priori absence of any regularity in time. However, we will often have the reason to use arguments in which some time regularity is actually needed of the solution. The key to overcoming this problem is to show that in many cases one can replace the solution by a version of it which has been mollified in time. There are several different time-mollifications that are used in the theory of nonlinear PDEs, such as the Steklov average, and the standard mollification taken only w.r.t. the time variable. The exponential time mollification, also known as the Naumann mollification, used in [34, 41] has turned out to be especially useful in the doubly nonlinear setting. For a function $w \in L^1(\Omega_T)$ the exponential time mollification is defined as

$$w_h(x, t) := \frac{1}{h} \int_0^t e^{\frac{s-t}{h}} w(x, s) ds, \quad (x, t) \in \Omega_T. \quad (2.11)$$

Moreover, we define its reversed analogue by

$$w_{\bar{h}}(x, t) := \frac{1}{h} \int_t^T e^{\frac{t-s}{h}} w(x, s) ds.$$

The time-mollification has many useful properties. For a function w in $L^p(\Omega_T)$ the mollification w_h is also p -integrable with a norm bounded by the norm of w . Moreover $w_h \rightarrow w$ in $L^p(\Omega_T)$ as $h \rightarrow 0$. Corresponding statements hold if the space $L^p(\Omega_T)$ is replaced by $L^p(0, T; W^{1,p}(\Omega))$. One of the most important properties of the exponential time mollification is that w_h has a weak time derivative on Ω_T given by

$$\partial_t w_h = \frac{1}{h} (w - w_h). \quad (2.12)$$

Correspondingly, for $w_{\bar{h}}$ we have

$$\partial_t w_{\bar{h}} = \frac{1}{h} (w_{\bar{h}} - w).$$

Since the time derivatives are linear combinations of the mollified function and the original function, it follows from the previously mentioned properties that $\partial_t w_h$ and $\partial_t w_{\bar{h}}$ both belong to $L^p(\Omega_T)$. Finally, we note that if $w \in L^p(0, T; L^p(\Omega))$ then $w_h \in C([0, T]; L^p(\Omega))$. The proofs of the properties presented above can be found in [34, Lemma 2.9], [4, Appendix B] and [5, Lemma 2.2].

3. Regularity properties

In this chapter we present the strategies of Publication I and Publication II, which allow us to conclude the local boundedness and local Hölder continuity of weak solutions to (1.1). Due to the nature of the equation, the arguments are somewhat technical and we seek to highlight the main ideas, explaining also why the adopted definition ensures rigorous proofs. All regularity results will be obtained in the *slow diffusion* case, i.e.

$$\alpha + p > 2.$$

This is also the range which is relevant for applications in glacier and shallow water dynamics.

3.1 Continuity in time

The first regularity property which we can prove directly from the definition of weak solutions is a certain continuity w.r.t. time. Namely, regarded as a map $[0, T] \rightarrow L_{\text{loc}}^{\beta+1}(\Omega)$ every weak solution has a continuous representative. This result allows us to interpret the value of v at a fixed time unambiguously, a fact which is needed for the mollified weak formulation presented below.

Using (2.4) with a suitable test function, we are able to show that the for any compact $K \subset \Omega$ we can approximate v essentially uniformly on the time interval by functions of the form $([v^\beta]_h)^{\frac{1}{\beta}}$ in the $L^{\beta+1}(K)$ -norm as $h \rightarrow 0$. Since the approximating functions are continuous on the time interval into $L^{\beta+1}(K)$, this guarantees that also v has the desired time continuity.

3.2 Mollified weak formulation

In order to prove that weak solutions satisfy energy estimates we need to use (2.4) with a test function which is proportional to the solution or some of its powers. However, since we are considering weak solutions, we cannot guarantee that such a test function is smooth, which is required by the definition. Especially the time derivative $\partial_t \varphi$ in (2.4) is problematic, since we did not assume that the solution has any regularity in time.

To overcome this problem we use a version of (2.4) in which the time derivative has been moved from the test function to v . This is not directly possible since v does not necessarily have a weak time derivative. Instead we need to use a mollified version of the weak formulation, in which v has been replaced by the time mollified function v_h . The time mollified formulation states that

$$\iint_{\Omega_T} [A(v, \nabla v^\beta)]_h \cdot \nabla \phi + \partial_t v_h \phi \, dx \, dt - \int_{\Omega} v(0) \phi_h(0) \, dx = \iint_{\Omega_T} f_h \phi \, dx \, dt, \quad (3.1)$$

for all $\phi \in C_0^\infty(\Omega_T)$, and can be proved by using (2.4) with the test function $\varphi = \phi_h \eta_\varepsilon$, where

$$\eta_\varepsilon(t) := \begin{cases} \frac{t}{\varepsilon} & \text{for } t \in [0, \varepsilon], \\ 1 & \text{for } t \in (\varepsilon, T]. \end{cases}$$

The nonsmooth test function is admissible due to the remarks at the end of Section 2.1.

Note that the time derivative in (3.1) is operating on v_h rather than ϕ . The term containing $v(0)$ on the left-hand side of (3.1) has a unique interpretation due to the previously established time continuity.

3.3 Energy estimates

One of the main tools in regularity theory are the energy estimates or Caccioppoli estimates that weak solutions satisfy. These provide a bound of the L^p -norm of the gradient in a super- or sublevel set in terms of the corresponding norm of the solution itself. For the shallow medium equation, the energy estimates state that for any $k \geq 0$ and $\varphi \in C^\infty(\Omega_T; \mathbb{R}_{\geq 0})$

vanishing in a neighborhood of the parabolic boundary, we have

$$\begin{aligned}
& \iint_{\Omega_T} |\nabla(v^\beta - k^\beta)_\pm|^p \varphi^p \, dx \, dt + \operatorname{ess\,sup}_{\tau \in (0, T)} \int_{\Omega} (v^{\frac{\beta+1}{2}} - k^{\frac{\beta+1}{2}})_\pm^2 \varphi^p(x, \tau) \, dx \quad (3.2) \\
& \leq c \iint_{\Omega_T} |\nabla \varphi|^p (v^\beta - k^\beta)_\pm^p \, dx \, dt + c \iint_{\Omega_T} (v^{\frac{\beta+1}{2}} - k^{\frac{\beta+1}{2}})_\pm^2 \varphi^{p-1} |\partial_t \varphi| \, dx \, dt \\
& \quad + c \iint_{\Omega_T} |\nabla z|^p v^{(\beta-1)p} \varphi^p \chi_{\{(v-k)_\pm > 0\}} \, dx \, dt + c \iint_{\Omega_T} (v^\beta - k^\beta)_\pm \varphi^p |f| \, dx \, dt,
\end{aligned}$$

for a constant c depending only on p and β . Comparing with the energy estimates for previously studied nonlinear equations, we see that the presence of z shows up as an extra term on the right-hand side. Also the source term gives rise to an extra term, as expected. To formulate the definition of weak solutions we only needed f to be locally integrable, but to prove (3.2) this is not sufficient. In Publication I we require that f is p' -integrable, where p' is the Hölder exponent of p . This also guarantees that the last term on the right-hand side is finite. A more careful analysis shows that one can replace $|f|$ by the positive part f_+ and require only f_+ to be p' -integrable.

To prove (3.2) we use the mollified weak formulation (3.1) with the test function $\phi = (v^\beta - k^\beta)_\pm \varphi^p \xi_\tau^\varepsilon$, where $\tau \in (0, T)$ and

$$\xi_\tau^\varepsilon(t) = \begin{cases} 1, & t \in [0, \tau], \\ 1 - \varepsilon^{-1}(t - \tau), & t \in (\tau, \tau + \varepsilon], \\ 0, & t \in (\tau + \varepsilon, T]. \end{cases}$$

Since there is no time derivative acting on ϕ in (3.1), using the non-smooth test function can be justified by approximation with smooth functions. The treatment of the term involving the time derivative merits some comments. In this term, we would like to apply the chain rule for Sobolev functions so we add and remove:

$$\partial_t v_h (v^\beta - k^\beta)_+ = \partial_t v_h ((v_h)^\beta - k^\beta)_+ + \partial_t v_h [(v^\beta - k^\beta)_+ - ((v_h)^\beta - k^\beta)_+].$$

Here it is important that in the quantity $(v_h)^\beta$ the mollification is performed on v before raising to the exponent β and not vice versa. The main reason for using the exponential time mollification becomes apparent from the second term. Due to the form of the time derivative (2.12), we see that this term is nonnegative, and we can thus omit this term replacing the equation by an inequality. Note that taking a test function which directly contains $(v_h)^\beta$ would create problems with the elliptic part of the equation since we do not know whether this function has a gradient. If instead we considered a test function containing $[v^\beta]_h$, the test function would have

a gradient and passing to the limit $h \rightarrow 0$ in the elliptic term would be no problem. However, this would not allow us to use the chain rule in the diffusion term, since the operations of taking time mollification and exponent do not commute.

We also remark that the vector field $A(v, \nabla v)$ belongs to $L^{\frac{p}{p-1}}$ whereas the gradient of the test function is in L^p . Since we have matching Hölder exponents, passing to the limit $h \rightarrow 0$ in this term works without problems. At this point we can also explain why the original form of the equation is problematic. With a vector field of the form $(u - z)^\alpha |\nabla u|^{p-2} \nabla u$, we would need to take a test function proportional to u , but assuming that u is in $L^p(0, T; W^{1,p}(\Omega))$ does not guarantee that the quantity $(u - z)^\alpha |\nabla u|^{p-2} \nabla u$ is in $L^{\frac{p}{p-1}}$, so we cannot ensure that the exponents match. All these considerations together show that the reformulated version of the equation is the natural formulation and that the exponential time mollification is the correct tool to obtain rigorous proofs of the energy estimates.

3.4 Local boundedness

In order to prove the local boundedness of solutions, we use a De Giorgi type iteration, combining the energy estimate (3.2) with a Sobolev embedding. For this strategy to work, we need to assume more regularity for the source term and the profile function than was needed to obtain the energy estimates. More precisely, we assume that there exists some $\sigma > \frac{p+n}{p}$ such that

$$f \in L^{p'\sigma}(\Omega_T) \quad \text{and} \quad z \in W^{1,\beta p\sigma}(\Omega). \quad (3.3)$$

Observe that the Sobolev exponent for z is so high that z is actually Hölder continuous. From a practical point of view this is not a big restriction, since in applications z is normally a Lipschitz function.

The De Giorgi iteration is based on the fact that the gradient of the solution appears only on the left-hand side in the energy estimates, and on the right-hand side in the parabolic Sobolev embedding (2.6), allowing us to eliminate the gradient and obtain an iterative estimate of the form

$$Y_{j+1} \leq C b^j Y_j^{1+\delta}, \quad (3.4)$$

where $C, b > 1, \delta > 0$ due to the lower bound on σ , and Y_j is of the form

$$Y_j = \iint_{Q_j} \left(v^{\frac{\beta+1}{2}} - k_j^{\frac{\beta+1}{2}} \right)_+^{\frac{2\beta p}{\beta+1}} dx dt.$$

Here Q_j is a sequence of shrinking cylinders converging to a limit cylinder Q and k_j is an increasing sequence converging to a positive number k . Moreover, $k_0 = 0$. The lemma of fast geometric convergence states that a sequence satisfying (3.4) will converge to zero provided that

$$Y_0 \leq C^{-\frac{1}{\delta}} b^{-\frac{1}{\delta^2}}, \quad (3.5)$$

see for example [24, Lemma 7.1]. In our case, the calculations show that C is proportional to a negative power of k , and Y_0 is a fixed number, so (3.5) is valid if we choose k sufficiently large. Observing furthermore that

$$\iint_Q \left(v^{\frac{\beta+1}{2}} - k^{\frac{\beta+1}{2}} \right)_+^{\frac{2\beta p}{\beta+1}} dx dt \leq Y_j,$$

we see that the integral on the left-hand side must be zero, i.e. $v \leq k$ a.e. in Q when k is chosen sufficiently large. Thus we have verified the local boundedness of v . Moreover, condition (3.5) and the k -dependence of C allow us to calculate a sufficiently large value for k in terms of Y_0 . In this way we obtain the following quantitative estimate for cylinders of the form $Q_\rho(z_o) := B_\rho(x_o) \times (t_o - \rho^{\frac{\beta+1}{\beta}}, t_o + \rho^{\frac{\beta+1}{\beta}})$:

$$\operatorname{ess\,sup}_{Q_{\frac{\rho}{2}}(z_o)} v \leq c \rho^{-\frac{n+p}{\beta+1}} \left[1 + \iint_{Q_\rho(z_o)} v^{\beta p} dx dt \right]^{\frac{1}{\beta+1}},$$

where $0 < \rho \leq 1$ and $c \geq 1$ is a constant that depends only on $n, \beta, p, \|f\|_{L^\sigma(\Omega_T)}, \|z\|_{W^{1,\beta p\sigma}(\Omega)}$ and σ . Note that since z is continuous, the local boundedness of v is actually equivalent to the local boundedness of the original function u .

3.5 Local Hölder continuity

In Publication II, we prove the local Hölder continuity of bounded weak solutions to (1.1) in the case $p > 2$. Since in this work boundedness of the weak solution is an assumption, the integrability conditions (3.3) for f and z from the previous section are not needed. In fact, (3.3) can be replaced by the weaker assumptions

$$f \in L^\sigma(\Omega_T) \quad \text{and} \quad z \in W^{1,p\sigma}(\Omega), \quad (3.6)$$

where $\sigma > \frac{p+n}{p}$ as before. Note that since $p\sigma > n$, z is Hölder continuous also in this case.

We use the method of intrinsic scaling which was originally introduced by DiBenedetto in [10] to prove local Hölder continuity for solutions to

the degenerate parabolic p -Laplace equation. Later, the method was extended to the singular case by Chen and DiBenedetto, [8]. See also the monograph [11]. A particularly clear exposition of the method in the degenerate case can be found in [50]. Systems of parabolic p -Laplace type have been considered in [12, 13]. For doubly nonlinear equations, Hölder regularity of weak solutions was verified in [52] and [43] under the extra assumption that the vector field in the elliptic part has sufficiently high integrability. This assumption is needed since the authors use a definition in which u (rather than a certain power of u) has a gradient. Solutions which are bounded away from zero were previously considered in [27]. The special case called Trudinger's equation, which is homogeneous, was investigated in [35] and [36]. In [25], doubly nonlinear equations with a definition consistent with ours are considered in a metric measure space setting for $p > 2$. Our result is a partial generalization of this work, since in the case $f = 0 = z$ the shallow medium equation is covered by their result.

The method of intrinsic scaling relies on showing the reduction of the oscillation in cylinders whose scale depends on the solution itself. In this way one can resolve problems stemming from the fact that unlike the classical heat equation, nonlinear equations are typically not homogeneous. The oscillation of the solution v on a subset $E \subset \mathbb{R}^{n+1}$ is defined as

$$\operatorname{ess\,osc}_E v := \operatorname{ess\,sup}_E v - \operatorname{ess\,inf}_E v.$$

We show that the oscillation decreases when passing from a parabolic cylinder to a smaller cylinder. Iterating this result leads to the local Hölder continuity. There are several reasons why obtaining such a result is nontrivial in the current setting. As mentioned earlier, we need to use an intrinsic scaling for the cylinders, i.e. the scaling depends on the solution itself. Moreover, as in the previous works [7, 25, 35, 36], the particular form of the scaling will depend on the relation between given upper and lower bounds of the solution, as we will now explain.

3.5.1 Scaling and De Giorgi type lemma

Let $0 \leq \mu_- < \mu_+ < \infty$ and set $\omega^\beta := \mu_+^\beta - \mu_-^\beta$. The numbers μ_- and μ_+ will represent lower and upper bounds for the solution in a suitable cylinder. We now introduce the scaling relevant in this setting. The two regimes

and scalings that we need to consider are:

$$\text{Degenerate:} \quad \mu_- \leq \frac{1}{2}\mu_+ \text{ and } \theta = \omega^{1-(p-1)\beta},$$

$$\text{Non-degenerate: } \mu_- > \frac{1}{2}\mu_+ \text{ and } (2\mu_+)^{1-\beta}\omega^{\beta(2-p)} \leq \theta \leq (\frac{1}{2}\mu_+)^{1-\beta}\omega^{\beta(2-p)}.$$

With these definitions we have the following result, which is known as a De Giorgi type lemma. Suppose $Q_{\rho,\theta\rho^p}(z_o) \subset \Omega_T$ is a parabolic cylinder satisfying

$$\operatorname{ess\,inf}_{Q_{\rho,\theta\rho^p}(z_o)} v \geq \mu_- \quad \text{and} \quad \operatorname{ess\,sup}_{Q_{\rho,\theta\rho^p}(z_o)} v \leq \mu_+,$$

and that

$$\xi := \rho^{p-\frac{n+p}{\sigma}}(\theta^{-p-\frac{1}{\sigma}}\omega^{\beta p(1-p)} + \theta^{-\frac{1}{\sigma}}\omega^{\beta(1-p)}) \max\{\|\nabla z\|_{L^\sigma}^p, \|f\|_{L^\sigma}\} \leq 1. \quad (3.7)$$

Then, there exists a number $\nu_o = \nu_o(n, p, \beta, \sigma) \in (0, 1)$ such that if

$$|Q_{\rho,\theta\rho^p}(z_o) \cap \{v^\beta \leq \mu_-^\beta + \frac{1}{2}\omega^\beta\}| \leq \nu_o |Q_{\rho,\theta\rho^p}(z_o)| \quad (3.8)$$

then

$$v^\beta \geq \mu_-^\beta + \frac{1}{4}\omega^\beta$$

a.e. in $Q_{\rho/2,\theta(\rho/2)^p}(z_o)$.

In practice, the lemma states that if v is close to the lower bound μ_- in only a small portion of the full cylinder $Q_{\rho,\theta\rho^p}(z_o)$ then we can find a better lower bound for v by passing to a subcylinder. In this way we have converted information about the measure into information about the pointwise behaviour of v .

The proof of the De Giorgi type lemma is based on a De Giorgi iteration combining energy estimates of a form similar to (3.2) with a Sobolev embedding. The presence of the source term and the term involving the gradient of z on the right-hand side of the energy estimate leads to the condition on ξ .

3.5.2 The two alternatives

When proving the reduction of the oscillation we initially consider space-time cylinders of the form $Q_{\rho,\theta_+\rho^p}(x_o, t_o)$ where $(x_o, t_o) \in \Omega_T$ and ρ is chosen so small that the cylinder is contained in Ω_T . Here, the number θ_+ is proportional to θ and can be written as $\theta_+ := 2^{\Lambda(p-2)}\theta$, where initially the parameter $\Lambda \in \mathbb{N}$ is not determined. In the end it turns out that one can choose a sufficiently large number Λ which only depends on the data.

One of the following alternatives must hold, since the second alternative is simply the negation of the first:

Alternative 1: There exists a time $t^* \in (t_o - \theta_+ \rho^p, t_o)$ such that the time interval $(t^* - \theta \rho^p, t^*)$ is contained in $(t_o - \theta_+ \rho^p, t_o)$ and on the corresponding cylinder $Q_{\rho, \theta \rho^p}(x_o, t^*)$ the measure estimate

$$|Q_{\rho, \theta \rho^p}(x_o, t^*) \cap \{v^\beta \leq \mu_-^\beta + \frac{1}{2}\omega^\beta\}| \leq \nu_o |Q_{\rho, \theta \rho^p}(x_o, t^*)| \quad (3.9)$$

holds true.

Alternative 2: The measure estimate

$$|Q_{\rho, \theta \rho^p}(x_o, t^*) \cap \{v^\beta \leq \mu_-^\beta + \frac{1}{2}\omega^\beta\}| > \nu_o |Q_{\rho, \theta \rho^p}(x_o, t^*)|$$

holds for all times $t^* \in (t_o - \theta_+ \rho^p, t_o)$ such that the time interval $(t_o - \theta \rho^p, t_o)$ is contained in $(t_o - \theta_+ \rho^p, t_o)$.

Here, ν_o denotes the constant from the De Giorgi type lemma. If the first alternative holds, then the De Giorgi type lemma implies the reduction of the oscillation in a subcylinder. However, in order to apply our method we need to replace the endpoint t^* of the time interval with t_o . This can be done by means of so-called logarithmic estimates which are obtained in a similar fashion as the energy estimates, but with a different test-function. Such estimates are used also in the regularity theory for the parabolic p -Laplacian, but in our case the presence of z shows up as an extra term on the right-hand side. Despite this additional complication, the logarithmic estimates allow us to control the measure on time-slices of the set where v is close to the lower bound all the way up to t_o . Aided by this result, another De Giorgi type iteration shows the reduction of the oscillation from below in a cylinder with terminal time t_o .

If instead the second alternative holds, we obtain a lower bound of the portion of $Q_{\rho, \theta \rho^p}(x_o, t^*)$ where $v^\beta > \mu_-^\beta + \frac{1}{2}\omega^\beta$. This can be used to conclude a corresponding lower bound on some time-slice $B_\rho(x_o) \times \{t^o\}$. We do not know the exact value of t^o , but logarithmic estimates can again be utilized to obtain measure estimates on time slices up to t_o . Finally, another De Giorgi type iteration shows, that if the scale Λ is chosen large enough, the oscillation is reduced from above. In this argument we also use the estimate (2.10).

The arguments in both the first and second alternative require that the quantity ξ is sufficiently small, and the bound only depends on the data. This can be achieved by choosing ρ sufficiently small. The conclusion of taking into account both alternatives is that there is a number $\tau_o \in (0, 1)$

depending only on the data such that

$$\operatorname{ess\,osc}_{Q_{\rho/8, \theta(\rho/8)^p}(z_o)} v^\beta \leq \tau_o \omega^\beta.$$

Thus, we are able to decrease the upper bound for the oscillation by passing to a smaller cylinder.

3.5.3 Iterating the reduction of the oscillation

We now explain how the reduction of oscillation can be iterated to demonstrate the local Hölder continuity. For all $j \in \mathbb{N}$, set

$$\omega_j^\beta := \tau_o^j \omega^\beta.$$

Consider first the case $\mu_- \leq \frac{1}{2}\mu_+$. We define

$$\theta_j = \omega_j^{1-(p-1)\beta}, \quad \theta_+^j = 2^{\Lambda(p-2)} \theta_j, \quad Q_j = Q_{\frac{\rho}{C^j}, \theta_+^j(\frac{\rho}{C^j})^p}(z_o),$$

where C is a sufficiently large number depending only on the data. The number has been chosen so that ξ on every level of the iteration will remain sufficiently small. Furthermore, we set

$$\mu_-^j = \operatorname{ess\,inf}_{Q_j} v, \quad (\mu_+^j)^\beta = (\mu_-^j)^\beta + \omega_j^\beta. \quad (3.10)$$

Suppose that $\mu_-^j \leq \frac{1}{2}\mu_+^j$ for all j smaller than some given integer j_0 . Then iterating the reduction of the oscillation shows that

$$\operatorname{ess\,osc}_{Q_j} v^\beta \leq \omega_j^\beta, \quad j \in \{1, \dots, j_0\}.$$

If $\mu_-^{j_0} > \frac{1}{2}\mu_+^{j_0}$, we need to redefine the cylinders starting from this index. For $j \geq j_0$ we set

$$\theta_j = (\mu_+^{j_0})^{1-\beta} \omega_j^{\beta(2-p)}, \quad \theta_+^j = 2^{\Lambda(p-2)} \theta_j, \quad Q_j = Q_{\frac{\rho}{C^j}, \theta_+^j(\frac{\rho}{C^j})^p},$$

where $\mu_+^{j_0}, \mu_+^{j_0}$ and ω_j are defined as before. For $j > j_0$ we use (3.10) to define μ_-^j and μ_+^j . One can now show that we remain in the nondegenerate regime on all of the following cylinders, i.e. $\mu_-^j > \frac{1}{2}\mu_+^j$ for $j \geq j_0$. Due to the modified scaling of the cylinders this allows us to again use the reduction of the oscillation to conclude that

$$\operatorname{ess\,osc}_{Q_j} v^\beta \leq \omega_j^\beta, \quad j \geq j_0.$$

In the case $\mu_- > \frac{1}{2}\mu_+$ the argument is easier, since then we can use the nondegenerate scaling from the start. Thus, we have found a shrinking sequence of cylinders with a fixed base point z_o so that the oscillation of

v^β shrinks on every level. In fact, we can verify uniform estimates for the oscillation when the base point z is allowed to vary in a neighborhood of z_o . Then the iterated reduction of oscillation implies estimates of the form

$$Q_{r, c\mu_+^{1-\beta}, \omega^{\beta(2-p)}(z)}^{\text{ess osc}} v^\beta \leq C\omega^\beta \left(\frac{r}{\rho}\right)^\alpha,$$

for all $r \in (0, \rho)$. Here $c, C > 0$ and $\alpha \in (0, 1)$ only depend on the data. Again the base point of the cylinder is allowed to vary in a neighborhood of z_o . The local Hölder continuity is a direct consequence of this estimate.

4. Existence

In Publication III we show the existence of solutions to a Cauchy-Dirichlet problem. Existence of solutions to certain doubly nonlinear equations have previously been established in [29, 30, 31]. See also [47] where different assumptions for the source term are considered. Existence of variational solutions to doubly nonlinear equations have been obtained in [6].

In Publication III we show the existence of solutions to the Cauchy-Dirichlet problem

$$\begin{cases} \partial_t u - \nabla \cdot ((u - z)^\alpha |\nabla u|^{p-2} \nabla u) = f & \text{in } \Omega_T, \\ u = z & \text{on } \partial\Omega \times (0, T), \\ u(\cdot, 0) = \psi & \text{in } \overline{\Omega}, \end{cases} \quad (4.1)$$

in the parameter range $\alpha > 0$, $p > 1$. Contrary to the regularity theory, we do not restrict ourselves to the slow diffusion case in this chapter. For the source term f and the land elevation z we use the conditions (3.3) which occurred earlier in the proof of local boundedness. Moreover, we need to assume that f is nonnegative.

From the viewpoint of applications the required integrability is reasonable, since typically f is bounded and z is Lipschitz in realistic models. The restriction to nonnegative source terms rules out infiltration and evaporation, but is necessary for our method to work.

For the initial values ψ we require only that

$$\Psi := \psi - z \in L^\infty(\Omega; \mathbb{R}_{\geq 0}).$$

The requirement that Ψ is nonnegative corresponds to the physical fact that the medium at time zero resides above the ground. Also the boundedness of Ψ is evidently physically motivated.

To formulate the exact definition of solutions we again make use of the function $v := u - z$. Stated in terms of v , the Cauchy-Dirichlet problem

takes the form:

$$\begin{cases} \partial_t v - \nabla \cdot A(v, \nabla v^\beta) = f & \text{in } \Omega_T, \\ v = 0 & \text{on } \partial\Omega \times (0, T), \\ v(\cdot, 0) = \Psi & \text{in } \overline{\Omega}, \end{cases} \quad (4.2)$$

where A is given by (2.3). We say that $u \geq z$ is a weak solution to the Cauchy-Dirichlet problem (4.1) if v solves the equation weakly (i. e. (2.4) is true), $v^\beta \in L^p(0, T; W_0^{1,p}(\Omega))$, $v \in C([0, T]; L^{\beta+1}(\Omega))$ and $v(0) = \Psi$. Note that the continuity in time means that $v(0)$ has a meaning. Moreover, in this case we say that v is a solution to (4.2).

In order to prove the existence of solutions to (4.2) we first consider a sequence of regularized Cauchy-Dirichlet problems for which it is easier to obtain solutions. A compactness result which has been proved in Publication III is then used to conclude the existence of a subsequence which converges to a function that solves the original problem.

4.1 Regularization of the problem

Consider the regularized Cauchy-Dirichlet problems

$$\begin{cases} \partial_t v_k - \nabla \cdot A_k(v_k, \nabla v_k) = f - k^{-\alpha} \nabla \cdot (|\nabla z|^{p-2} \nabla z) & \text{in } \Omega_T, \\ v_k = \frac{1}{k} & \text{on } \partial\Omega \times (0, T), \\ v_k(\cdot, 0) = \frac{1}{k} + \Psi & \text{in } \overline{\Omega}, \end{cases}$$

where $k > 1$ is an integer and the regularized vector field A_k is defined as

$$A_k(v, \xi) := \begin{cases} A(\frac{1}{k}, \beta \frac{1}{k^{\beta-1}} \xi), & v < \frac{1}{k} \\ A(v, \beta v^{\beta-1} \xi), & \frac{1}{k} \leq v \leq k \\ A(k, \beta k^{\beta-1} \xi), & v > k. \end{cases} \quad (4.3)$$

Since A_k can be written as $A_k(v, \xi) = T_k(v)^\alpha |\xi + \nabla z|^{p-2} (\xi + \nabla z)$, where T_k denotes the truncation $T_k(s) := \min\{k, \max\{s, \frac{1}{k}\}\}$, the regularized equation is essentially of p -Laplace type.

The existence of a solution v_k to the regularized problem is proved using Galerkin's method, i. e. by approximation with finite-dimensional problems. This technique has been applied to nonlinear parabolic equations also in the past, see for example Section III.4 in [45]. The existence of solutions to the finite-dimensional problems is guaranteed by an extended version of the Cauchy-Peano theorem, which we have also proved in Publication III.

4.2 Properties of the approximating solutions

The last term on the right-hand side of the regularized equation disappears in the limit $k \rightarrow \infty$, and ensures that the constant function $\frac{1}{k}$ is a solution to the approximating equation in the case $f = 0$. This allows us to use a comparison principle argument as in [30] to conclude that $v_k \geq \frac{1}{k}$. In this argument we also need the nonnegativity of f .

There is also a uniform upper bound for the functions v_k . Establishing this fact is one of the more delicate points of our argument. In fact, we must initially work with the truncated functions $\tilde{v}_k := \min\{v_k, k\}$. The first step is to establish an energy estimate for $\tilde{v}_k$ in which the gradient of $\tilde{v}_k^\beta$ appears on the left-hand side. Combining this result with the Poincaré inequality, we obtain a uniform bound for the $L^{\beta p}$ -norm of $\tilde{v}_k$. Finally, a De Giorgi type iteration shows that $\tilde{v}_k$ is bounded by $L^{\beta p}$ -norm. From this we conclude that $\tilde{v}_k$ and hence also v_k has an upper bound which is independent of k . The necessity of the assumption $\sigma > \frac{n+p}{p}$ for the parameter occurring in the conditions (3.3) for the source and profile function shows up in the De Giorgi type iteration.

One consequence of the lower and upper bounds for v_k is that for large k we have $\frac{1}{k} \leq v_k \leq k$ which means that A_k can be replaced by A in the regularized equation. The equation together with the uniform upper bound for v_k then imply that also the L^p -norm of ∇v_k^β is bounded independently of k .

4.3 Compactness and convergence results

The uniform bound for the functions v_k and the L^p -bound for the gradients ∇v_k^β show that (v_k^β) is a bounded sequence in the reflexive space $L^p(0, T; W^{1,p}(\Omega))$. Therefore, there is a subsequence which converges weakly to an element of $L^p(0, T; W^{1,p}(\Omega))$.

In Publication III we have proved some compactness results which allow us to conclude a stronger form of convergence. They are consequences of the results of Simon [46]. Some similar results for uniformly bounded functions that are piecewise constant in time have previously been obtained, see [32] and its references. Even though we prove several theorems in the hope of wider applicability, only Corollary 4.7 of Section 4 is used in the existence proof. This result provides sufficient conditions for the precompactness of a subset of $L^p(0, T; L^q(\Omega))$ where $p, q \geq 1$ are suitable

parameters. Apart from the previously mentioned bounds for v_k , in order to apply the corollary, we also need the fact that

$$\lim_{h \rightarrow 0+} \|\tau_h v_k - v_k\|_{L^{\beta p}(0, T-h; (W_0^{1,p}(\Omega))')} = 0, \quad \text{uniformly in } k.$$

We have verified this condition using the equation with a suitable test function. The conclusion is that there is a subsequence of (v_k^β) which converges strongly in $L^p(\Omega_T)$. Due to the uniform bound on the functions v_k we can even conclude that the corresponding subsequence of v_k converges strongly in $L^q(\Omega_T)$ for every $q \in [1, \infty)$. By passing to another subsequence we may also assume pointwise convergence almost everywhere.

Using the approximating equation and the monotonicity property of the vector field A we have also verified that the subsequence of (∇v_k^β) converges strongly in $L^p(\Omega \times I)$ for every closed interval $I \subset (0, T)$.

4.4 Existence

Aided by the convergence results from the previous section we can now pass to the limit in the approximating equation and conclude that

$$\iint_{\Omega_T} [A(v, \nabla v^\beta) \cdot \nabla \varphi - v \partial_t \varphi] dx dt = \iint_{\Omega_T} f \varphi dx dt + \int_{\Omega} \Psi \varphi(0) dx, \quad (4.4)$$

for every $\varphi \in C^\infty(\bar{\Omega} \times [0, T])$ with support contained in $K \times [0, \tau]$ where $K \subset \Omega$ is compact and $\tau \in (0, T)$. This immediately implies that v satisfies the original equation since the integral involving Ψ disappears if φ is compactly supported in Ω_T . Moreover, using (4.4) with the test function $\varphi(x)\zeta_\varepsilon(t)$ where $\varphi \in C_0^\infty(\Omega)$ and

$$\zeta_\varepsilon(t) = \begin{cases} \frac{1}{\varepsilon}(\varepsilon - t), & t \in [0, \varepsilon], \\ 0, & t \geq \varepsilon, \end{cases}$$

one sees that

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} \int_0^\varepsilon \int_{\Omega} v(x, t) \varphi(x) dx dt = \int_{\Omega} \Psi \varphi dx.$$

On the other hand, as in Chapter 3 we can verify that the weak solution v belongs to $C([0, T]; L^{\beta+1}(\Omega))$. This implies that also

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} \int_0^\varepsilon \int_{\Omega} v(x, t) \varphi(x) dx dt = \int_{\Omega} v(0) \varphi dx$$

holds. Since $\varphi \in C_0^\infty(\Omega)$ is arbitrary, it follows that $v(0) = \Psi$, i. e. v satisfies the correct initial value. Finally, since $v_k^\beta - k^{-\beta}$ belongs to the closed subspace $L^p(0, T; W_0^{1,p}(\Omega))$ of $L^p(0, T; W^{1,p}(\Omega))$, the weak convergence shows that also $v^\beta \in L^p(0, T; W_0^{1,p}(\Omega))$, so that the correct boundary values are satisfied. Thus, v is a solution to the Cauchy-Dirichlet problem (4.2).

5. Doubly singular diffusion equations

In Publication IV we prove a wide range of regularity results for doubly singular equations of the form

$$\partial_t u - \nabla \cdot A(x, t, u, \nabla u^\beta) = 0, \quad (5.1)$$

where $A(x, t, u, \xi)$ is a vector field satisfying

$$|A(x, t, u, \xi)| \leq C_1 |\xi|^{p-1} \quad (5.2)$$

$$A(x, t, u, \xi) \cdot \xi \geq C_0 |\xi|^p. \quad (5.3)$$

Here, $\beta = 1 + (m - 1)/(p - 1)$, and the parameters m and p satisfy

$$p \in (1, 2), \quad m > 1, \quad \text{and} \quad 2 < m + p < 3. \quad (5.4)$$

Applying the chain rule formally as in Section 2.1 one can see that the model case equation

$$\partial_t u - \nabla \cdot (u^{m-1} |\nabla u|^{p-2} \nabla u) = 0,$$

falls into this category.

There were several motivations for studying this type of equations. The first objective was to give a simple proof of the local Hölder continuity of weak solutions, aided by the L^1 -Harnack inequality and the expansion of positivity which were previously obtained in [19, 20]. However, the authors make use of the gradient of u , whereas the definition of solutions requires only u^β to have a gradient. Even though their approach might be justified, since in the fast diffusion case one can reason as in Section 5 of [28] to conclude that also u itself should have a locally p -integrable gradient, we thought it interesting to show that the original strategies of [19, 20] can be modified so that only the gradient of u^β appears in the calculations. Thus, we have also proved the L^1 -Harnack inequality and the expansion of positivity directly from the definition.

To be specific, we say that a function $u: \Omega_T \rightarrow \mathbb{R}$ is a weak solution to (5.1) if and only if $u \geq 0$, $u^\beta \in L^p(0, T; W^{1,p}(\Omega))$, $u \in L^{\beta+1}(\Omega_T)$ and

$$\iint_{\Omega_T} A(x, t, u, \nabla u^\beta) \cdot \nabla \varphi - u \partial_t \varphi \, dx \, dt = 0, \quad (5.5)$$

for all $\varphi \in C_0^\infty(\Omega_T)$.

The extra $L^{\beta+1}$ -integrability is necessary to justify the test function in the proof of the energy estimates, since $\beta p < \beta + 1$ in the fast diffusion case. In the slow diffusion case the inequality is reversed so that the assumption can be avoided.

5.1 Energy estimates

The method presented in Section 3.1, can be used to show that also weak solutions to (5.1) are time continuous on the closed interval $[0, T]$ into $L_{loc}^{\beta+1}(\Omega)$. This allows us to prove a mollified weak formulation, which then enables us to conclude the following energy estimate:

$$\begin{aligned} & \iint_{\Omega_T} |\nabla(u^\beta - k^\beta)_\pm|^p \varphi^p \, dx \, dt + \operatorname{ess\,sup}_{\tau \in [0, T]} \int_{\Omega} \mathfrak{b}[u, k] \chi_{\{(u-k)_\pm > 0\}} \varphi^p(x, \tau) \, dx \\ & \leq C \iint_{\Omega_T} (u^\beta - k^\beta)_\pm^p |\nabla \varphi|^p \, dx \, dt + \iint_{\Omega_T} \mathfrak{b}[u, k] \chi_{\{(u-k)_\pm > 0\}} |\partial_t \varphi^p| \, dx \, dt, \end{aligned}$$

for all smooth $\varphi \geq 0$ defined on $\bar{\Omega}_T$, vanishing in a neighborhood of the parabolic boundary of Ω_T . The constant C only depends on the data.

5.2 L^1 -Harnack inequality

Following the strategy of [19], we have shown the L^1 -Harnack inequality for weak solutions. Contrary to [19], we do not need the gradient of u in our argument. The precise statement is as follows. Let u be a weak solution to (5.1) where the vector field $A(x, t, u, \xi)$ satisfies the structure conditions (5.2) and (5.3), and the parameters satisfy the conditions (5.4). Then there exists a positive constant γ depending only on m, n, p, C_0, C_1 such that for all cylinders $\bar{B}_{2\rho}(y) \times [s, t] \subset \Omega \times [0, T]$,

$$\operatorname{ess\,sup}_{\tau \in [s, t]} \int_{B_\rho(y)} u(x, \tau) \, dx \leq \gamma \operatorname{ess\,inf}_{\tau \in [s, t]} \int_{B_{2\rho}(y)} u(x, \tau) \, dx + \gamma \left(\frac{t-s}{\rho^\lambda} \right)^{\frac{1}{3-m-p}}, \quad (5.6)$$

where $\lambda = n(p + m - 3) + p$.

5.3 Expansion of positivity

Using the strategy of [20], we have verified that weak solutions exhibit expansion of positivity. The main difference in our approach is that we do not need the gradient of u . Instead, we rely only on the existence of ∇u^β . The expansion of positivity result states the following. Suppose that $(x_o, s) \in \Omega_T$ and u is a weak solution satisfying

$$|B_\rho(x_o) \cap \{u(\cdot, s) \geq M\}| \geq \alpha |B_\rho(x_o)|,$$

for some $M > 0$ and $\alpha \in (0, 1)$. Then there exist $\varepsilon, \delta, \eta \in (0, 1)$ depending only on $m, p, n, C_0, C_1, \alpha$ such that if $B_{16\rho}(x_o) \times (s, s + \delta M^{3-m-p}\rho^p) \subset \Omega_T$ then

$$u \geq \eta M \text{ in } B_{2\rho}(x_o) \times (s + (1 - \varepsilon)\delta M^{3-m-p}\rho^p, s + \delta M^{3-m-p}\rho^p).$$

The proof makes use of a change in variables and a transformed function which satisfies a doubly nonlinear equation with a source term which depends on the solution itself. Combining energy estimates with the isoperimetric inequality (2.10) leads to measure estimates for sublevel sets of this function in a certain cylinder. After this, one segments the cylinder to get the proper intrinsic scaling so that a De Giorgi type lemma can be applied. This converts the measure information to pointwise information, namely a lower bound for the function. Finally, we return to the original coordinates and obtain the expansion of positivity.

5.4 Local boundedness

Using a De Giorgi type iteration we have showed the local boundedness of weak solutions in the range

$$m + p > 3 - \frac{p}{n - (\frac{n-p}{p})}. \quad (5.7)$$

This condition is necessary for our method to work. In the case $m = 1$, i.e. equations of singular parabolic p -Laplace type, the condition reduces to $p > 2n/(n + 2)$ and the extra integrability assumption becomes $u \in L^2$. It is well-known that these are sufficient conditions for local boundedness for the singular parabolic p -Laplace equation, see for example Chapter V of [11] where these conditions correspond to the parameter choice $r = 2$. The author even claims that such conditions are sharp.

The De Giorgi iteration provides a bound for the essential supremum in terms of the $L^{\beta+1}$ -norm of the solution on a cylinder which is compactly

contained in Ω . This result can be somewhat improved by another iterative procedure. First, note that (5.7) can be phrased as

$$(\beta + 1)p + n(m + p - 3) > 0.$$

Thus, there exist $r \in (0, \beta + 1)$ such that

$$\lambda_r := rp + n(m + p - 3) > 0. \quad (5.8)$$

It turns out that we can obtain a bound also in terms of the L^r -norm of u . Combining these two boundedness results we can state that for any $r \in (0, \beta + 1]$ for which (5.8) is valid, and any cylinder of the form $Q_{\rho, 2\tau}(z_o)$ contained in Ω_T and any weak solution u , we have

$$\operatorname{ess\,sup}_{Q_{\rho, \tau}(z_o)} u \leq c \left[\tau^{-\frac{n+p}{p}} \iint_{Q_{2\rho, 2\tau}(z_o)} u^r \, dx \, dt \right]^{\frac{p}{\lambda_r}} + c \left(\frac{\tau}{\rho^p} \right)^{\frac{1}{3-m-p}} \quad (5.9)$$

where the constant c depends only on r, m, n, p and the constants C_0 and C_1 appearing in the structure conditions.

5.5 Local Hölder continuity

We have proved the local Hölder continuity of weak solutions in the supercritical range

$$m + p > 3 - \frac{p}{n}. \quad (5.10)$$

It is well-known that the expansion of positivity combined with the integral Harnack inequality lead to a concise proof of the local Hölder continuity of bounded weak solutions to the singular parabolic p -Laplace equation, although references for this specific approach, which also we have adopted, do not seem to be available in the literature. For a related method relying solely on the expansion of positivity, we refer to Section 5 of [16]. The degenerate parabolic p -Laplace equation was considered in [23]. Recently, a unified approach which covers signed solutions of both the parabolic p -Laplace and the porous medium equation has been presented in [38]. These various approaches provide shorter proofs than the classical argument of DiBenedetto.

In the case of the singular parabolic p -Laplace equation, the two alternatives of DiBenedetto's approach can be replaced by a single argument since for a solution $0 \leq u \leq 1$ also $1 - u$ is a solution. In the doubly nonlinear setting this is no longer true, so our proof is somewhat more complicated. We first prove a De Giorgi type lemma which provides sufficient

conditions for obtaining the reduction of the oscillation from above. If the conditions of the lemma fail, we can use the integral Harnack inequality to obtain the reduction of the oscillation from below.

The condition (5.10) is used when proving the reduction of the oscillation from below, and is needed for controlling the size of the second term on the right-hand side of the integral Harnack inequality (5.6). As usual, also an intrinsic scaling is required for these arguments.

5.6 Harnack Estimates

Apart from the previously obtained integral Harnack inequality we have also verified that in the supercritical range (5.10), weak solutions satisfy Harnack estimates on properly scaled cylinders. More precisely, suppose that u is a weak solution in a domain containing $B_{34R}(\bar{0}) \times [-T, T]$. Suppose that $u(0, 0) > 0$ and

$$4 R^p \sup_{B_{2R}(\bar{0})} u(\cdot, 0)^{3-m-p} \leq T. \quad (5.11)$$

Then there exist constants $C \geq 1$, $\theta > 0$ depending only on m, n, p, C_0, C_1 such that

$$\begin{aligned} C^{-1} \sup_{B_{R/4}(\bar{0})} u(\cdot, s) &\leq u(0, 0) \leq C \inf_{B_R(\bar{0})} u(\cdot, t), \\ \text{for } -\theta u(0, 0)^{3-m-p} R^p &\leq s, t \leq \theta u(0, 0)^{3-m-p} R^p. \end{aligned} \quad (5.12)$$

Here we are considering the continuous representative of u . The result has been formulated for a cylinder centered at the origin, but obviously the result is translation invariant. Since an infimum can only increase when passing to a smaller set, we could replace the ball in the right inequality of (5.12) by $B_{R/4}(\bar{0})$ so that the supremum and infimum are taken over the same ball.

Note that the admissible times s and t in (5.12) belong to the same interval. This type of Harnack estimates are exclusive to the fast diffusion case, and have been established previously for solutions to the singular parabolic p -Laplace equation in the supercritical range $p \in (\frac{2n}{n+1}, 2)$, see [11], Chapter 6 of [15] or [14]. In these sources it is also demonstrated that the subcritical range is sharp.

By contrast, in the slow diffusion case, there is a time gap between s and t . See for example Chapter 5 of [15] which treats the degenerate parabolic p -Laplace equation. The necessity of the time gap can be seen from the

finite speed of propagation exhibited for example by the self-similar solutions found by Barenblatt [2]. A time gap occurs also in the case of linear diffusion, see [40], and for Trudinger's equation, which corresponds to the border case $m + p = 3$, see [49, 33].

Harnack estimates of the form (5.12) have been obtained for doubly nonlinear equations in the supercritical range in [21] under the assumption that u has a gradient. We avoid this extra assumption. Our method follows roughly the pattern of [16]. A key observation is that in the supercritical range, we can take $r = 1$ in (5.9). Combining this estimate with the integral Harnack inequality on a suitable subcylinder allows us to use the reduction of the oscillation to get a pointwise lower bound in a smaller cylinder. Using the expansion of positivity we then obtain a lower bound on the whole cylinder, thus proving the right inequality in (5.12).

In order to prove the left inequality we apply similar arguments to a suitably rescaled and translated version of the solution to obtain a bound for the supremum of u on the time-slice $B_R(\bar{0}) \times \{0\}$. After this, an upper bound for u on the whole space-time cylinder can be obtained by combining (5.9) and the integral Harnack inequality as before.

Bibliography

- [1] R. Alonso, M. Santillana, and C. Dawson. On the diffusive wave approximation of the shallow water equations. *European J. Appl. Math.*, 19(5):575–606, 2008.
- [2] G.I. Barenblatt. On self-similar motions of compressible fluids in porous media (in Russian). *Prikl. Mat. Meth.*, 16:679–698, 1952.
- [3] V. Bögelein, F. Duzaar, R. Korte, and C. Scheven. The higher integrability of weak solutions of porous medium systems. *Advances in Nonlinear Analysis*, 8(1):1004–1034, 1 2019.
- [4] V. Bögelein, F. Duzaar, and P. Marcellini. Parabolic systems with p, q -growth: A variational approach. *Archive for Rational Mechanics and Analysis*, 210(1):219–267, 2013.
- [5] V. Bögelein, F. Duzaar, and P. Marcellini. Existence of evolutionary variational solutions via the calculus of variations. *Journal of Differential Equations*, 256(12):3912–3942, 2014.
- [6] V. Bögelein, F. Duzaar, P. Marcellini, and C. Scheven. Doubly nonlinear equations of porous medium type. *Archive for Rational Mechanics and Analysis*, 229(2):503–545, Aug 2018.
- [7] V. Bögelein, T. Lukkari, and C. Scheven. Hölder regularity for degenerate parabolic obstacle problems. *Ark. Mat.*, 55(1):1–39, 09 2017.
- [8] Y.-Z. Chen and E. DiBenedetto. Hölder estimates of solutions of singular parabolic equations with measurable coefficients. *Archive for Rational Mechanics and Analysis*, 118(3):257–271, Sep 1992.
- [9] E. De Giorgi. Sulla differenziabilità e l’analiticità delle estremali degli integrali multipli regolari. *Mem. Accad. Sci. Torino. Cl. Sci. Fis. Mat. Nat.*, III(3):25–43, 1957.
- [10] E. DiBenedetto. On the local behaviour of solutions of degenerate parabolic equations with measurable coefficients. *Annali della Scuola Normale Superiore di Pisa - Classe di Scienze*, 13(3):487–535, 1986.
- [11] E. DiBenedetto. *Degenerate Parabolic Equations*. Springer Verlag, 1993.
- [12] E. DiBenedetto and A. Friedman. Regularity of solutions of nonlinear degenerate parabolic systems. *Journal für die reine und angewandte Mathematik*, 349:83–128, 1984.

- [13] E. DiBenedetto and A. Friedman. Hölder estimates for nonlinear degenerate parabolic systems. *Journal für die reine und angewandte Mathematik*, 357:1–22, 1985.
- [14] E. DiBenedetto, U. Gianazza, and V. Vespri. Forward, backward and elliptic harnack inequalities for non-negative solutions to certain singular parabolic partial differential equations. *Annali della Scuola Normale Superiore di Pisa*, 9(2):385–422, 2010.
- [15] E. DiBenedetto, U. Gianazza, and V. Vespri. *Harnack's Inequality for Degenerate and Singular Parabolic Equations*. Springer Monographs in Mathematics, 2012.
- [16] F. G. Düzgün, S. Mosconi, and V. Vespri. Harnack and pointwise estimates for degenerate or singular parabolic equations. In Serena Dipierro, editor, *Contemporary Research in Elliptic PDEs and Related Topics*, pages 301–368. Springer INdAM Series, 07 2019.
- [17] L. C. Evans. *Partial Differential Equations*. American Mathematical Society, 2 edition, 2010.
- [18] K. Feng and F. J. Molz. A 2-d diffusion based, wetland flow model. *J. Hydrol.*, 196(1):230–250, 1997.
- [19] S. Fornaro, M. Sosio, and V. Vespri. Energy estimates and integral harnack inequality for some doubly nonlinear singular parabolic equations. *Contemporary Mathematics*, 594:179–199, 01 2013.
- [20] S. Fornaro, M. Sosio, and V. Vespri. $L^r_{\text{loc}} - L^\infty_{\text{loc}}$ estimates and expansion of positivity for a class of doubly non linear singular parabolic equations. *Discrete and Continuous Dynamical Systems*, 7(4):737–760, 2013.
- [21] S. Fornaro, M. Sosio, and V. Vespri. Harnack type inequalities for some doubly nonlinear singular parabolic equations. *Discrete and Continuous Dynamical Systems*, 35(12):5909–5926, 2015.
- [22] P. Di Giammarco, E. Todini, and P. Lamberti. A conservative finite elements approach to overland flow: the control volume finite element formulation. *Journal of Hydrology*, 175(1):267 – 291, 1996.
- [23] U. Gianazza, M. Surnachev, and V. Vespri. A new proof of the hölder continuity of solutions to p -laplace type parabolic equations. *Advances in Calculus of Variations*, 3:263–278, 07 2010.
- [24] E. Giusti. *Direct Methods in the Calculus of Variations*. World Scientific, 2003.
- [25] E. Henriques and R. Laleoglu. Local and global boundedness for some nonlinear parabolic equations exhibiting a time singularity. *Differential Integral Equations*, 29(11/12):1029–1048, 2016.
- [26] T. V Hromadka, C. E. Berenbrock, J. R Freckleton, and G. L. Guymon. A twodimensional dam-break flood plain model. *Adv. Water Resour.*, 8(1):7–14, 1985.
- [27] A. V. Ivanov. Hölder estimates for quasilinear doubly degenerate parabolic equations. *Journal of Soviet Mathematics*, 56(2):2320–2347, Aug 1991.

- [28] A. V. Ivanov. On the maximum principle for doubly nonlinear parabolic equations. *Journal of Mathematical Sciences*, 80:2236–2254, 08 1996.
- [29] A. V. Ivanov and P. Z. Mkrtchyan. Existence of Hölder continuous generalized solutions of the first boundary value problem for quasilinear doubly degenerate parabolic equations. *Journal of Mathematical Sciences*, 62:2725–2740, 1992.
- [30] A. V. Ivanov, P. Z. Mkrtchyan, and W. Jäger. Existence and uniqueness of a regular solution of the Cauchy-Dirichlet problem for a class of doubly nonlinear parabolic equations. *Journal of Mathematical Sciences*, 84(1):411–435, 1997.
- [31] A. V. Ivanov, P. Z. Mkrtchyan, and W. Jäger. Erratum to: “Existence and uniqueness of a regular solution of the Cauchy-Dirichlet problem for a class of doubly nonlinear parabolic equations”. *J. Math. Sci., New York*, 184(6):786–787, 2012.
- [32] A. Jüngel. Boundedness of weak solutions to cross-diffusion population systems with Laplacian structure. *arXiv e-prints*, page arXiv:1812.08719v1, Dec 2018.
- [33] J. Kinnunen and T. Kuusi. Local behaviour of solutions to doubly nonlinear parabolic equations. *Mathematische Annalen*, 337(3):705–728, Mar 2007.
- [34] J. Kinnunen and P. Lindqvist. Pointwise behaviour of semicontinuous supersolutions to a quasilinear parabolic equation. *Annali di Matematica Pura ed Applicata*, 185:411–435, 08 2006.
- [35] T. Kuusi, R. Laleoglu, J. Siljander, and J-M. Urbano. Hölder continuity for Trudinger’s equation in measure spaces. *Calculus of Variations and Partial Differential Equations*, 45(1):193–229, 2012.
- [36] T. Kuusi, J. Siljander, and J-M. Urbano. Local Hölder continuity for doubly nonlinear parabolic equations. *Indiana University Mathematics Journal*, 61(1):399–430, 2012.
- [37] P. Lehtelä. *Generalized solutions to the porous medium equation*. PhD thesis, Aalto University, 2018.
- [38] N. Liao. A unified approach to the Hölder regularity of solutions to degenerate and singular parabolic equations. *Journal of Differential Equations*, 2019.
- [39] M. W. Mahaffy. A three-dimensional numerical model of ice sheets: Tests on the Barnes ice cap, northwest territories. *J. Geophys. Res.*, 81(6):1059–1066, 1976.
- [40] J. Moser. A Harnack inequality for parabolic differential equations. *Communications on Pure and Applied Mathematics*, 17(1):101–134, 1964.
- [41] J. Naumann. *Einführung in die Theorie parabolischer Variationsungleichungen*. Teubner-Texte zur Mathematik 64. Leipzig: B.G. Teubner, 1984.
- [42] D. Partridge. *Numerical Modelling of Glaciers: Moving Meshes and Data*. PhD thesis, University of Reading, 2013.

- [43] M.M. Porzio and V. Vespri. Holder estimates for local solutions of some doubly nonlinear degenerate parabolic equations. *Journal of Differential Equations*, 103(1):146 – 178, 1993.
- [44] M. Santillana and C. Dawson. A numerical approach to study the properties of solutions of the diffusive wave approximation of the shallow water equations. *Computational Geosciences*, 14(1):31–53, 2010.
- [45] R. E. Showalter. *Monotone Operators in Banach Space and Nonlinear Partial Differential Equations*. American Mathematical Society, 1997.
- [46] J. Simon. Compact sets in the space $L^p(0, T; B)$. *Annali di Matematica Pura ed Applicata*, 146:65–96, 01 1986.
- [47] S. Sturm. Existence of weak solutions of doubly nonlinear parabolic equations. *J. Math. Anal. Appl.*, 455(1):842–863, 2017.
- [48] E. Todini and M. Venutelli. Overland flow: A two-dimensional modeling approach. In David S. Bowles and P. Enda O’Connell, editors, *Recent Advances in the Modeling of Hydrologic Systems*, pages 153–166. Springer Netherlands, Dordrecht, 1991.
- [49] N. Trudinger. Pointwise estimates and quasilinear parabolic equations. *Communications on Pure and Applied Mathematics*, 21(3):205–226, 1968.
- [50] J. M. Urbano. *The Method of Intrinsic Scaling*. Springer-Verlag Berlin Heidelberg, 2008.
- [51] J.L. Vazquez. *The Porous Medium Equation: Mathematical Theory*. Oxford Mathematical Monographs. Clarendon Press, 2006.
- [52] Vincenzo Vespri. On the local behaviour of solutions of a certain class of doubly nonlinear parabolic equations. *manuscripta mathematica*, 75(1):65–80, Dec 1992.
- [53] Th. Xanthopoulos and Ch. Koutitas. Numerical simulation of a two dimensional flood wave propagation due to dam failure. *J. Hydraulic Res.*, 14(4):321–331, 1976.

ISBN 978-952-60-3862-9 (printed)

ISBN 978-952-60-3863-6 (pdf)

ISSN 1799-4934 (printed)

ISSN 1799-4942 (pdf)

Aalto University

School of Science

Department of Mathematics and Systems Analysis

www.aalto.fi

**BUSINESS +
ECONOMY**

**ART +
DESIGN +
ARCHITECTURE**

**SCIENCE +
TECHNOLOGY**

CROSSOVER

**DOCTORAL
DISSERTATIONS**