Design of selective Sodium Calcium Exchanger inhibitors for cardiac indications:

Chronic treatments and acute on/off use.

Leena Otsomaa

Design of selective Sodium Calcium Exchanger inhibitors for cardiac indications:

Chronic treatments and acute on/off use.

Leena Otsomaa

A doctoral dissertation completed for the degree of Doctor of Science (Technology) to be defended, with the permission of the Aalto University School of Technology, at a public examination held at the lecture hall Ke2 (Komppa) of the school on the 30th of August 2019 at 12:00.

Aalto University
School of Chemical Engineering
CHEM, CMAT

Supervising professor

Professor Ari Koskinen, Aalto University, Finland

Thesis advisor

Senior Vice President R&D, M.D., Ph.D., Christer Nordstedt, R&D, Orion Corporation, Finland

Preliminary examiners

Professor, Dr Christa Müller, University of Bonn, Germany

Professor Jari Yli-Kauhaluoma, University of Helsinki, Finland

Opponent

Professor Nick Turner, University of Manchester, UK

Aalto University publication series

DOCTORAL DISSERTATIONS 139/2019

© 2019 Leena Otsomaa

ISBN 978-952-60-8660-6 (printed)

ISBN 978-952-60-8661-3 (pdf)

ISSN 1799-4934 (printed)

ISSN 1799-4942 (pdf)

<http://urn.fi/URN:ISBN:978-952-60-8661-3>

Unigrafia Oy

Helsinki 2019

Finland

Publication orders (printed book):

leena.otsomaa@orionpharma.com

Author

Leena Otsomaa

Name of the doctoral dissertation

Design of selective Sodium Calcium Exchanger inhibitors for cardiac indications:

Publisher School of Chemical Engineering

Unit CHEM, CMAT

Series Aalto University publication series DOCTORAL DISSERTATIONS 139/2019

Field of research Medicinal Chemistry

Manuscript submitted 8 July 2019

Date of the defence 30 August 2019

Permission for public defence granted (date) 19 June 2019

Language English

☐ **Monograph**

☒ **Article dissertation**

☐ **Essay dissertation**

Abstract

The $\text{Na}^+/\text{Ca}^{2+}$ exchanger (NCX) is a transmembrane protein which plays an important role in regulating cardiomyocyte Ca^{2+} homeostasis. NCX operates in two modes: Forward (exit mode) and Reverse (entry mode). In the forward mode, NCX extrudes Ca^{2+} ions from the intracellular matrix that is proposed to have effects on relaxation in conditions where the intracellular Ca^{2+} concentration is high. In the reverse mode, NCX intrudes Ca^{2+} ions from the extracellular matrix in conditions where the intracellular Na^+ concentration is high to enable depolarization of the membrane potential. Selective inhibition of NCX1 has been proposed to create cardio protection in certain conditions, hence various research groups in academia and industry have been studying the role, function and therapeutic value of NCX1.1 and have had activities to develop novel NCX 1.1 inhibitors that could be used as cardioprotective therapy.

Based on current public data, six pharma companies have worked in the field with NCX inhibitors including Orion Corporation. The objective of this thesis was to develop potent selective and novel NCX1 inhibitors for chronic treatments and acute on-off use for cardiac indications with desired PK properties. While for the chronic indications the long-term and steady exposure to the inhibitor molecule is the ideal, for the acute indications an inhibitor with relatively fast PD effect followed by fast clearance would be preferred.

The presented work describes the development of two novel potent and selective NCX1.1 inhibitors with different chemical structures, which enable scientific communities to study further functions and therapeutic value of NCX inhibition.

The first part of the thesis focuses on the background and the rational for developing NCX 1.1 inhibitors for acute and chronic indications. The second and the major part of the thesis describes the discovery of two distinctively different and novel chemical series of NCX inhibitors and optimization strategy towards desired DMPK properties. Optimization was guided by the results from in vitro test battery of the project workflow. The work resulted in three patent applications filed by the Orion Corporation and the most selective NCX 1.1 inhibitors (ORM-10962 and ORM-11372) reported in the literature.

The third part of the thesis focuses on the studies made in Orion and in the collaboration with academic groups to further reveal the pharmacological properties of the two lead molecules ORM-10962 and ORM-11372 from the different chemical series. At the same time these most selective tool molecules (ORM-10962 and ORM-11372) has given the opportunity for the academic groups to study further the role, function and therapeutic value of NCX1.1.

Final part of the thesis describes the status of NCX 1.1 inhibitor research in Orion Corporation and pilot toxicology result of the ORM-11372.

Keywords Design, Optimisation, NXC, Medicinal Chemistry

ISBN (printed) 978-952-60-8660-6

ISBN (pdf) 978-952-60-8661-3

ISSN (printed) 1799-4934

ISSN (pdf) 1799-4942

Location of publisher Helsinki

Location of printing Helsinki

Year 2019

Pages 300

urn <http://urn.fi/URN:ISBN:978-952-60-8661-3>

Tekijä

Leena Otsomaa

Väitöskirjan nimi

Selektiivisten NCX inhibiittoreiden suunnittelu sydänsairauksien krooniseen ja akuuttiin hoitoon.

Julkaisija Kemian tekniikan korkeakoulu**Yksikkö** CHEM, CMAT**Sarja** Aalto University publication series DOCTORAL DISSERTATIONS 139/2019**Tutkimusala** Lääkeainekemia**Käsikirjoituksen pvm** 08.07.2019**Väitöspäivä** 30.08.2019**Väittelyluvan myöntämispäivä** 19.06.2019**Kieli** Englanti☐ **Monografia**☒ **Artikkeliväitöskirja**☐ **Esseeväitöskirja****Tiivistelmä**

Transmembraaniproteiini Na⁺/Ca²⁺-vaihtajalla (NCX) on tärkeä rooli kardiomyosyytien kalsium (Ca²⁺) tasapainon säätelyssä. NCX toimii kaksisuuntaisesti: poistaa Ca²⁺:a solusta (suunta ulospäin) tai tuo Ca²⁺:a solun sisään (suunta sisäänpäin). Nykykäsityksen mukaan NCX vaikuttaa relaksaatioon solunsisäisen Ca²⁺-ionikonsentraation ollessa suuri vaihtamalla solunsisäisestä matriisista Ca²⁺-ioneja ulos. Vastaavasti solunsisäisen Na⁺-ionikonsentraation ollessa suuri NCX siirtää Ca²⁺ ioneja solunulkoisesta matriisista solun sisään membraanipotentiaalin neutraloimiseksi. Selektiivisellä NCX1-inhibitiolla on esitetty olevan sydäntä suojaavia vaikutuksia tietyissä sydänsairauksissa. Tämän vuoksi monet tutkimusryhmät tiedemaailmassa mukaan lukien Orion Oyj ovat keskittyneet tutkimaan NCX:n toimintaa, merkitystä ja mahdollista terapeutista arvoa. Lisäksi nykyisen tiedon valossa kuudella lääkeyrityksellä on ollut NCX1.1-inhibiittoritutkimusprojekteja tarkoituksena kehittää uusia tehokkaita NCX1-inhibiittoreita krooniseen sekä akuuttiin "katkaisija"-tyyppiseen hoitoon ideaalisella farmakokineettisellä profiililla. Kroonisessa indikaatiossa ideaalilla molekyylillä saavutetaan pitkäaikainen ja tasainen altistus oraalisesti. Akuutissa indikaatiossa toivottu "katkaisija"-tyyppinen farmakokineettinen profiili saavutetaan suhteellisen nopealla farmakodynaamisella vaikutuksella, jota seuraa nopea lääkeaineen poistuminen elimistöstä (suuri puhdistuma).

Väitöskirjatyön tavoitteena oli kehittää uusia tehokkaita NCX1-inhibiittoreita krooniseen sekä akuuttiin "katkaisija"-tyyppiseen hoitoon ideaalisella farmakokineettisellä profiililla. Kroonisessa indikaatiossa ideaalilla molekyylillä saavutetaan pitkäaikainen ja tasainen altistus oraalisesti. Akuutissa indikaatiossa toivottu "katkaisija"-tyyppinen farmakokineettinen profiili saavutetaan suhteellisen nopealla farmakodynaamisella vaikutuksella, jota seuraa nopea lääkeaineen poistuminen elimistöstä (suuri puhdistuma). Väitöskirjassa kuvataan kahden uuden, kemiallisesti hyvin erilaisen, tehokkaan ja selektiivisen NCX1.1-inhibiittorin keksiminen ja kehittäminen, jotka mahdollistavat luotettavamman NCX-inhibition toiminnan, merkityksen sekä terapeutin potentiaalin tutkimisen tiedeyhteisössä. Työn ensimmäisessä osassa keskitytään taustaan sekä perusteluun, miksi NCX inhibitiomekanismilla on esitetty olevan sydäntä suojaavia ominaisuuksia akuuteissa ja kroonisissa indikaatioissa. Väitöskirjatyön toinen ja myös pääasiallinen osa kuvaa kahden uuden ja toisistaan erilaisen kemiallisen sarjan keksimisen sekä kemiallisen rakenteen optimointistrategian kohti toivottua farmakokineettistä profiilia. Työn tuloksena syntyi kolme Orionin jättämää patenttihakemusta sekä kaksi selektiivisintä NCX-inhibiittoria, joita kirjallisuudessa on koskaan raportoitu (ORM-10962 ja ORM-11372).

Työn kolmannessa osassa kuvataan farmakologiset *in vivo* ja *ex vivo*-mallit, joilla molekyylit ORM-10962 ja ORM-1137 karakterisoitu tarkemmin ja niistä saadut tulokset. Tutkimukset ovat tehty joko Orionilla tai yhdessä akateemisten ryhmien kanssa. Samalla nämä kaksi tunnetuinta selektiivistä ja tehokasta työkalumolekyyliä (ORM-10962 ja ORM-11372) ovat mahdollistaneet akateemisten ryhmien tutkimukset NCX1.1:n merkityksestä, toiminnasta ja terapeuttisesta arvosta.

Väitöskirjatyön viimeisessä osassa päivitetään NCX1.1:n tutkimuksen tilanne Orionissa sekä ORM-1137:lla tehdyn toksikologisen pilottitutkimuksen tulokset.

Avainsanat Suunnittelu, optimointi, NCX, Lääkeainekemia**ISBN (painettu)** 978-952-60-8660-6**ISBN (pdf)** 978-952-60-8661-3**ISSN (painettu)** 1799-4934**ISSN (pdf)** 1799-4942**Julkaisupaikka** Helsinki**Painopaikka** Helsinki**Vuosi** 2019**Sivumäärä** 300**urn** <http://urn.fi/URN:ISBN:978-952-60-8661-3>

Acknowledgements

I am thankful and feel emotional, but at the same time extremely tired, after pushing through the writing process in parallel with my daily duties. It is definitely not a plain and easy to do on top of the every-day work life. However, I found it absolutely great to see the whole story, I've done among the NCX inhibitors during my early days in Orion. Unfortunately, the work did not deliver new medication for the cardiac patients who have unmet needs, even though it was always my primary motivation. However, it at least gave me the opportunity to increase knowledge in a science community and publish my results in the PhD thesis. I am truly grateful to my employer Orion Corporation for giving me the opportunity to publish the work and for being very supportive for my efforts.

I definitely want to give my sincere thanks to my supervising Professor Ari Koskinen. You have encouraged me through the process and difficult times when I had my doubts. I also feel pleased that I was able to pursue for doctoral degree under your supervision in Aalto University – this all started long time ago in the University of Oulu when I was making my licentiate thesis.

I am thankful for my advisor, Senior Vice President R&D, Christer Nordstedt for his support. You truly gave me the motivation to write the thesis, but you also supported my efforts and worried about my wellbeing. Many thanks for that!

I am also very grateful to all my co-authors which are many in this kind of cross disciplinary topic. Especially, I want to gratefully acknowledge Jouko Levijoki, Norbert Jost, Kaisa Syrjänen and Hugh Chapman for their co-operation and scientific input. Jouko's support in the field of pharmacology has been extremely important during the experimental part, but also during the writing of the Paper I. We indeed had couple of good laugh! Didn't we?

I also like to give my big thanks to my colleague Timo Lotta for keeping up my spirit, giving valuable comments and his support. Indeed, you are extremely positive and selfless supportive person. Great to have such a colleague!

In addition, there are many other current and former Orion colleagues, I like to thank for all their support and help. Just to mention few teams e.g. Global Medicine Design supervisory team and R&D Management team. I feel privileged to work with all of you.

My most grateful thanks go to my family and friends for supporting me through this journey and being always understanding and empathic. Last but not least, my warmest thanks are for my husband Jussi. Your everlasting support and tender care to maintain balance in life, has been the most critical for me during this journey.

List of publications

This thesis consists of an overview of the following four publications, which from here onwards are referred to by their Roman numerals in the text:

- I** **Otsomaa L.**, Levijoki J., Wohlfahrt G., Chapman H., Koivisto A.P., Syrjänen K., Koskelainen T., Peltokorpi S.E., Finckenberg P., Heikkilä A., Abi-Gerges N., Ghetti A., Page G., Miller P.E., Mervaala E., Kohajda Z., Jost N., Virág L., Varró A., Papp J.Gy. Discovery and characterization of unique positively inotropic sodium-calcium exchanger inhibitor ORM-11372. Submitted to *Br J Pharmacol*. Manuscript ID 2019-BJP-0720-RP.
- II** Jost N., Nagy N., Corici C., Kohajda Z., Horvath A., Acsai K., Biliczki P., Levijoki J., Pollesello P., Koskelainen T., **Otsomaa L.**, Toth A., Papp J.Gy., Varró A., Virag L. (2013) ORM-10103, a novel specific inhibitor of the $\text{Na}^+/\text{Ca}^{2+}$ exchanger, decreases early and delayed afterdepolarizations in the canine heart. *Br J Pharmacol* **170**, 768-778. DOI: 10.1111/bph.12228
- III** Kohajda Z., Farkas-Morvay N., Jost N., Nagy N., Geramipour A., Horvath A., Varga R.S., Hornyik T., Corici C., Acsai K., Horvath B., Prorok J., Ordog B., Deri S., Toth D., Levijoki J., Pollesello P., Koskelainen T., **Otsomaa L.**, Toth A., Bacsko I., Lepran I., Nanasi P.P., Papp J.Gy., Varró A., Virag L. (2016) The effect of a novel highly selective inhibitor of the sodium/calcium exchanger (NCX) on cardiac arrhythmias in *in vitro* and *in vivo* experiments. *PLoS One*. **11**, e0166041/1-e0166041/28. DOI: 10.1371/journal.pone.0166041
- IV** Oravec K., Kormos A, Gruber A., Márton Z., Kohajda Z., Mirzaei L., Jost N., Levijoki J., Pollesello P., Koskelainen T., **Otsomaa L.**, Tóth A., Papp J.Gy., Nánási P.P., Antoons G., Varró A., Acsai K., Nagy N. Inotropic effect of NCX inhibition depends on the relative activity of the reverse NCX assessed by a novel inhibitor ORM-10962 on canine ventricular myocytes. (2018) *European Journal of Pharmacology* **818**,278–286. DOI: 10.1016/j.ejphar.2017.10.039

Author's contribution

I Paper I is based on the invention described in appendix 03, where the author is the main inventor. The author was responsible for devising the conceptual approach for the optimization program, defined the design strategy, and led the optimization based on the pharmacological results. She analysed data and drew conclusions for the structure optimization. Furthermore, the author also synthesized many of the derivatives during the optimization program. J. Levijoki, N. Jost, L. Virág, A. Varró and J. Papp were responsible for the pharmacological conceptualization and *in vivo* and *ex vivo* characterisation, as well as data visualization and analyses. G. Wohlfahrt performed the virtual screening to find original hit for the chemical series. K. Syrjänen synthesised ORM-11372 for the first time based on defined optimization program. She developed the synthesis method for ORM-11372. The author and J. Levijoki were responsible for the selection the compound for the studies based on the property profile gained during the SAR optimization of the scaffold, which is presented in appendix 03. T. Koskelainen looked after the scaling up of the synthesis for the further studies. Others were responsible for biological methodologies and testing, including bioanalysis as follows: A-P Koivisto and H. Chapman contributed by the electrophysiological investigation, methodology, and data analysis including iPS; S.E. Peltokorpi contributed to the pharmacological investigation; P. Finckenberg contributed to the histopathological methodology, formal and data analysis; A. Heikkilä contributed to the bioanalytical methodology and data analysis; E. Mervaala participated in the pharmacological conceptualization; Z. Kohajda contributed to pharmacological investigation. N. Abi-Gerges, A. Ghetti, G. Page and P.E. Miller were responsible for human *ex-vivo* tissue study as follows: N. Abi-Gerges was responsible for study design, data analysis, data interpretation, and contributed to the writing. G. Page and P.E. Miller contributed to study design and data interpretation. A. Ghetti contributed to study design, performed experiments and data analysis. The author wrote the original draft of the manuscript together with another main co-author J. Levijoki. They also share the senior authorship of the manuscript. K. Syrjänen participated to the writing of the supplementary material under supervision of the author; H. Chapman contributed to the *in vitro* methodology part in the manuscript; E. Mervaala and N. Jost critically reviewed original draft of the manuscript and contributed to the writing; G. Wohlfahrt generated the pharmacophore and critically reviewed the manuscript.

II Paper II is based on the appendix 01 invention, where the author is the one of the two main inventors. The author was responsible for devising the conceptual approach for the optimization program, defined the design strategy and led the optimization. T. Koskelainen applied the defined design strategy, synthesized the ORM-10103, and developed the first synthesis route. P. Pollesello and J. Levijoki were responsible for the pharmacological conceptualization. The author and J. Levijoki selected the compound ORM-10103 for the study based on the property profile gained during the SAR optimization of the scaffold that is presented in appendix 01. In addition, the author contributed to the paper II by revealing the chemical structure of the study compound. Academic parties were responsible for biological methodologies, and biological testing, as well as data visualization and analyses as follows: J.G. Papp, A. Varró, and L. Virág made a major contribution to the supervising, conceptualization, data curation, project administration, and resources, A. Varró and L. Virág contributed also to the validation. Furthermore L. Virág made a major contribution also to formal analysis, electrophysiological investigation, methodology, software, and visualization. N. Jost and N. Nagy contributed to the formal analysis, electrophysiological and pharmacological investigation, methodology, software, supervision, and visualization. In addition, A. Tóth contributed to the resources, supervision, data curation, formal analysis, and visualization. C. Corici contributed to the pharmacological investigation. Z Kohajda, A. Horváth, K. Acsai, P and P. Biliczki contributed to pharmacological investigation, methodology, visualization and formal analysis. Z Kohajda and K. Acsai contributed also to software. N. Jost, N. Nagy, C. Corici, A. Tóth, A. Varró and L. Virág contributed to the writing.

III The paper III is based on the appendix 02 invention, where the author is the main inventor. The author was responsible for devising the conceptual approach for the optimization program, and defined the design strategy and led the optimization. The author also designed and synthesized ORM-10962, which hence enabled the study. P. Pollesello and J. Levijoki were responsible for the pharmacological conceptualization from Orion. The author and J. Levijoki selected the compound ORM-10962 for the study based on the property profile gained during the SAR optimization of the scaffold that is presented in appendix 02. In addition, the author contributed to the paper III by revealing the chemical structure and supply of the study compound. Academic parties were responsible for biological methodologies and biological testing, as well as data visualization and analyses as follows: Virág L. and Varró A. share senior authorship for the work. They both were responsible for the conceptualization, project administration in academia,

resources, supervision, validation and data curation. In addition, Virág L. contributed to the formal analysis, software and methodologies, as well as electrophysiological investigation. Papp J.G. contributed to the conceptualization, project administration, resources and supervision as well. Geramipour A., Horváth A., Varga R.S., Hornyik T., Corici C., Déri S., Tóth D. and Leprán I. contributed to the pharmacological investigation. Leprán I. also contributed to the validation. Kohajda Z. contributed to the formal analysis, pharmacological investigation, methodology, software, and visualization. Farkas-Morvay N., Acsai K., Horváth B., Prorok J. and Ördög B. contributed to the formal analysis, pharmacological investigation, methodology and visualization. Others contributed to the software, except Farkas-Morvay and Ördög B. Jost N. contributed to the conceptualization, data curation, formal analysis, pharmacological and electrophysiological investigation, methodology, project administration, resources, software, supervision, validation and visualization. Nagy N. also contributed to the formal analysis, cellular electrophysiological investigation, methodology, software, supervision and visualization. Tóth A. and Baczkó I. both contributed to the data curation, formal analysis, methodology, resources, and visualization. In addition, Tóth A. supervised and Baczkó I. contributed to the investigation and validation. Nánási P.P. contributed to the data curation, resources, and validation. Data curation means the lifecycle management of data, from creation to the archiving or deletion when obsolete. Z. Kohajda, N. Jost, N. Nagy, A. Tóth, I. Leprán, and P.P. Nánási, also contributed to the writing.

IV The paper IV is based on the appendix 02 invention, where the author is the main inventor. The author was responsible for devising the conceptual approach for the optimization program, and defined the design strategy and led the optimization. The author also designed and synthesized ORM-10962, which hence enabled the study. P. Pollesello and J. Levijoki were responsible for the pharmacological conceptualization. The author and J. Levijoki selected the compound ORM-10962 for the study based on the property profile gained during the SAR optimization of the scaffold, which is presented in appendix 02. In addition, the author contributed to the paper III by supply of the study compound. Academic parties were responsible for biological methodologies and biological testing, as well as data visualization and analyses as follows: A. Varró and J.G. Papp made a major contribution to the conceptualization, data curation, project administration, supervising, and resources. In addition A. Varró also contributed to the validation. N. Jost contributed to the formal analysis, pharmacological and electrophysiological investigation, methodology, software, supervision, and

visualization. A. Tóth contributed to the data curation, formal analysis, resources, and supervision. K. Oravecz contributed to the biological investigation, methodology, formal analysis, software, and visualization. K. Acsai, and Z. Kohajda contributed to the formal analysis, electrophysiological/pharmacological investigation, methodology, software, and visualization. N. Nagy also contributed to supervision, formal analysis, electrophysiological and pharmacological investigation, methodology, software, and visualization. A. Kormos, A. Gruber., Z. Márton and L. Mirzaei, contributed to the electrophysiological investigation, formal analysis, methodology, and visualization. P.P. Nánási and G. Antoons contributed to the resources. Furthermore P.P. Nánási contributed to the data curation and validation. K. Oravecz, N. Jost, A. Tóth, P.P., Nánási, G. Antoons, A. Varró and N. Nagy also contributed to the writing.

Nomenclature

Abbreviations

AC	Aftercontractions
ADMET	Absorption, Distribution, Metabolism, Excretion and Toxicity
Caco-2	Heterogeneous human epithelial colorectal adenocarcinoma cells
CHF	Congestive heart failure
CHO	Chinese hamster ovary cells
CYP	Cytochrome P450
DAD	Delayed afterdepolarization
DMPK	Drug metabolism and pharmacokinetics
DMTA	Design, make, test, analyse
dP/dt	Left ventricular contractility
2D VS	Two-dimensional virtual screening
EAD	Early afterdepolarization
EDTA	Ethylenediaminetetraacetic acid
EGTA	Egtazic acid
EWG	Electron withdrawing group
F	Forward mode of NCX
Fluo-4-AM	Ca ²⁺ dye
Fluo-3FF	Ca ²⁺ dye
GS	Glucosteril
HAM	Growth medium for mammalian cells
HEK	Human embryonic kidney
HEPES	4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid

hERG	Human ether-a`-go-go-related gene-encoded voltage dependent potassium channel
HF	Heart failure
IC ₅₀	The half maximal inhibitory concentration
I _{NCX}	Na ⁺ /Ca ²⁺ exchanger current
I.U	International Units
IV, iv	Intravenous
LC-MS	Liquid chromatography–mass spectrometry
logP	Logarithm of partition coefficient for compounds
LV	Left ventricle
LVH	Left ventricular hypertrophy
LVP	Left ventricular pressure
m-	Meta-
MAO	Monoamine Oxidase
MEM	Minimal essential medium
MLA	Mouse Lymphoma Assay
mOsm	Milliosmoles
NAD(P)(+)	Nicotinamide adenine dinucleotide phosphate
NAD(P)H	Reduced form of nicotinamide adenine dinucleotide phosphate
NaV 1.5	Voltage-gated sodium channel subunit
NCX	Na ⁺ /Ca ²⁺ exchanger
NMDG	<i>N</i> -Methyl-D-glucamine
o-	Orto-
p-	Para-
PPB	Plasma Protein Binding
R	Reverse mode of NCX
rpm	Rounds per minutes

RT	Room temperature
SAP	Systolic arterial blood pressure
SAR	Structure Activity Relationship
SCD	Sudden cardiac death
Sf9	<i>Spodoptera frugiperda</i> derived insect cell culture
SFR	Start of fast rise
SLC8A1	Solute carrier family 8 member A1
SPR	Structure Property Relationship
SR	Sarcoplasmic reticulum
TEACl	Tetraethylammonium Chloride
TtM	Time to Maximum
UV	Ultraviolet light
VT	Ventricular tachycardia
XIP	Exchange inhibitory peptide

Symbols

n	Number of experimental repetitions
T	Absolute temperature [K]

Contents

Acknowledgements.....	i
List of publications	ii
Author's contribution.....	iii
Nomenclature.....	vii
Abbreviations.....	vii
Symbols	x
1 Introduction.....	1
1.1 Thesis objective	5
2 Rationale for the use of NCX inhibitors in cardiac indications	6
2.1 NCX inhibitors in chronic heart failure indications.....	6
2.2 NCX inhibitors in acute cardiac indications	6
3 Design strategies for the chemical series	8
3.1 Design of novel NCX inhibitors for chronic indications	8
3.1.1 Discovery of chemical series A	8
3.1.2 Optimization strategy and SAR/SPR of the series A.....	14
3.2 Design of novel NCX inhibitors for acute indications.....	21
3.2.1 Discovery of chemical series B.....	21
3.2.2 Optimization strategy and SAR/SPR of the series B	22
3.3 Summary	29
4 Materials and methods	34
4.1 Synthesis routes for series A and B	34
4.1.1 Series A route.....	34
4.1.2 Series B route.....	35
4.2 <i>In vitro</i>	36
4.2.1 NCX1.1 inhibition activity assays	36
4.2.2 Cardiac safety ion channel assays.....	37
4.2.3 ADME assays	39
4.2.4 Other miscellaneous assays.....	41
4.3 <i>Ex vivo</i> and <i>In vivo</i>	42
4.3.1 Ouabain model.....	42
4.3.2 Ischemia-reperfusion induced arrhythmias.....	44
4.3.3 In vivo hemodynamics of ORM-11372	45
5 Results and discussion	47
5.1 <i>In vivo</i> and <i>Ex vivo</i> results for novel NCX inhibitors	47

5.2	Status of NXC inhibitor project in Orion	51
5.3	Legislative perspectives	51
6	Conclusions	53
6.1	Significance of this work.....	54
6.2	Future research recommendations	54
	References	56
	Appendices	59

1 Introduction

The $\text{Na}^+/\text{Ca}^{2+}$ exchanger (NCX) is a transmembrane protein which plays an important role in regulating cardiomyocyte Ca^{2+} homeostasis [1]. NCX1 is the predominant isoform in the heart whereas other isoforms NCX2 and 3 are primarily restricted to the brain and skeletal muscle. NCX is 120 kD and has 10 transmembrane helices with two alpha repeats connected with an intracellular f-loop. The loop contains Ca^{2+} and Na^+ binding sites [2]. NCX1 is encoded by the SLC8A1 gene, which is alternatively spliced leading to different expression in different tissues *e.g.* NCX1.1 is mainly expressed in heart while in kidney NCX1.3 and NCX1.7 are expressed [3].

Figure 1 A topological model of NCX1 structure (corrected from earlier understanding)[2, 4].

NCX1 is a high capacity ionic transporter that exchanges three Na^+ ions for one Ca^{2+} ion. Depending on the electrochemical gradients and membrane potential, NCX functions in either the Ca^{2+} -efflux (forward) or the Ca^{2+} -influx (reverse) mode. As NCX has an electrogenic nature it has also been reported to play a role in reducing the incidence and duration of arrhythmias as well as enhance recovery of developed tension [5].

Figure 2 Illustration of functional mode of NCX (www.AddexBio.com/membrane-exchangers/services)

NCX operates in two modes (**Figure 2**): Forward (exit mode) and Reverse (entry mode). In the forward mode NCX extrudes Ca^{2+} ions from the intracellular matrix which is proposed to have effects on relaxation in conditions where the intracellular Ca^{2+} concentration is high. In the reverse mode NCX intrudes Ca^{2+} ions from the extracellular matrix in conditions where the intracellular Na^{+} concentration is high to enable depolarization of the membrane potential. This could lead to a reperfusion injury after *e.g.* an ischemic event. Selective inhibition of NCX1 has been proposed to create cardioprotection in certain conditions, hence various research groups in academia and industry have been working in the field. Based on current public data, six pharma companies have worked in the field with NCX inhibitors: Kanebo Ltd.[6], Taisho Pharmaceutical [7], Allergan [8][9], Orion Corporation, Astellas Pharma (Yamanouchi Pharmaceutical)[10, 11] and Sanofi [12]. **Table 01** presents the structures of known NCX inhibitors and their publication year with the originator organization.

The main focus of interest of pharma activities have been ischemic heart diseases, but ischemic brain and kidney diseases have been under investigation as well as retinal ganglion cell death associated with glaucoma. NCX inhibitors have shown cell protecting properties in thrombolytic or in surgery for blood vessel formation, or during coronary artery bypasses or organ transplantation. Cardiovascular indications with treatment of disorders caused by an overload of Ca^{2+} ions in myocardial ischemia-reperfusion injuries such as heart function disorders (heart failure, arrhythmias and strokes), myocardial infraction and arrhythmias. The first published and hence most studied NCX inhibitors, KB-R7943 and SEA0400, have differing potency and selectivity. Magee *et al.* have studied these compounds for their cardioprotective properties in isolated rabbit hearts leading to the conclusion that more selective inhibition leads to a marked reduction in myocardial ischemic injury and improved (dP/dt) left ventricular (LV) contractility [1].

Table 1 Illustration of known NCX inhibitor structures

Ref.	Structure	Code	Originator	Year of publishing
[6]		KB-R7943	Kanebo Ltd.	1997
[13]		-	Taisho Pharmaceuticals	1999
[7]		SEA0400	Taisho Pharmaceuticals	1999
I		ORM-10103	Orion Pharma	2003
II		ORM-10962	Orion Pharma	2004

Table 1 (Continue)

Ref.	Structure	Code	Originator	Year of publishing
[10, 11]		YM-244796	Astellas Pharma	2005
[10, 11]		-	Astellas Pharma	2005
[12]		-	Sanofi	2014
III	H ₂ N ⁺ *HCl ⁻	ORM-11372	Orion Pharma	2019

1.1 Thesis objective

The objective of this thesis was to develop potent selective and novel NCX1 inhibitors for chronic treatments and acute on-off use for cardiac indications. Chronic treatments hold potential for arrhythmias and left ventricular hypertrophy (LVH), while acute treatments could be useful new treatments for acute decompensated heart failure or low cardiac output syndrome. The dissertation is based on the optimization work (Figure 3) mainly carried out over the years 1999-2006. The publications of the dissertation (2013-2019) are enabled by the inventions that were made by Otsomaa et al. during the 1999-2006 period. Based on the inventions, Orion filed three patent applications that are appended to the dissertation. Otsomaa was one of the two main inventors with an equal share in the Appendix 01 patent application and the main inventor of the patent applications described in Appendix 02 and 03. Cardiovascular research at Orion was terminated in the year 2007 as a business decision, following the trend of pharma companies globally.

The research tasks for achieving the thesis objectives were:

- (i) To find novel chemistry for NCX inhibition
- (ii) To optimize chemical series of NCX inhibitors with different drug metabolism and pharmacokinetics (DMPK) properties
- (iii) To further characterize pharmacological and other properties of the best compounds of each series

The research work followed the key optimization process (DMTA-cycle, Design, Make, Test, Analyze) as illustrated in **Figure 3** [14]. The design and SAR analysis as well as most of the in vitro testing are described in the dissertation summary. Make part of the each example molecules that were patented are described in the appendices 01-03. Further pharmacological profiling (=testing) of the selected compounds were published in the publications I-IV of the dissertation. Pilot safety evaluation of the best compound for acute indication was also carried out.

Figure 3 Fields of research work described in this thesis.

2 Rationale for the use of NCX inhibitors in cardiac indications

NCX can be modulated for the long term with an inhibitor molecule or relatively short term *i.e.* chronic or acute cardiac indications respectively. Chronic heart failure and heart failure with a preserved ejection fraction (= diastolic heart failure) were considered as chronic cardiac indications. Antiarrhythmic and positive inotropic effects of NCX inhibitors and their ability to prevent ischemia-reperfusion injuries provides the potential to use NCX inhibitors for acute cardiac indications as well. While for the chronic indications the long-term and steady exposure to the inhibitor molecule is the ideal, for the acute indications an inhibitor with relatively fast PD effect followed by fast clearance would be preferred.

2.1 NCX inhibitors in chronic heart failure indications

In cardiomyocytes from failing hearts, the accumulation of insufficient mitochondrial Ca^{2+} secondary to a cytoplasmic Na^+ overload reduces the NAD(P)H/NAD(P)(+) redox potential and increases oxidative stress as the workload increases. These effects are abolished by enhancing the mitochondrial Ca^{2+} with acute treatment with CGP-37157 (CGP), an inhibitor of the mitochondrial $\text{Na}^+/\text{Ca}^{2+}$ exchanger. In a model of heart failure with sudden cardiac death (HF/SCD) induced by aortic constriction combined with daily beta-adrenergic receptor stimulation, chronic CGP treatment attenuates cardiac hypertrophic remodeling, pulmonary edema, and interstitial fibrosis and prevents cardiac dysfunction and SCD [15].

In a model of heart failure with a preserved ejection fraction (= diastolic heart failure), the left ventricle (LV) is hypertrophied, and LV isovolumetric relaxation time as well as decay of intracellular Ca^{2+} transient are prolonged and the diastolic pressure is elevated. NCX inhibitor SEA0400 acutely improves relaxation *in vivo* and in isolated cardiomyocytes. Furthermore, chronic treatment with SEA0400 improves the cardiac remodelling and diastolic functions [16].

2.2 NCX inhibitors in acute cardiac indications

Endogenous inhibitors of cardiac NCX1 effectively suppress both atrial and ventricular arrhythmia [17]. Pre-treatment with a novel NCX inhibitor GYKB-6635 significantly delayed the time to the development of ventricular fibrillation (by about 18%). It was concluded that GYKB-6635 is a potent and highly selective inhibitor of the cardiac NCX and it has also been suggested to contribute to the prevention of delayed after depolarisation (DAD)-based arrhythmias [18]. In an experimental model of CHF induced by rapid

ventricular pacing in rabbits, the acute inhibition of NCX (1) reduces the monophasic action potential duration, (2) decreases dispersion of repolarization, and (3) suppresses early afterdepolarization (EAD) and ventricular tachycardia (VT). The study shows that pharmacological NCX inhibition increases repolarization reserve and protects against VTs in heart failure [19]. Hence NCX inhibitors are considered to have an antiarrhythmic effect.

In myocytes isolated from a canine tachycardic pacing-induced failing heart, the intracellular application of exchange inhibitory peptide (XIP) normalized the intracellular calcium transient ($[Ca^{2+}]_i$) amplitude and concomitantly increased the sarcoplasmic reticulum (SR) Ca^{2+} load. The degree of NCX inhibition at this concentration of XIP was 27% and was selective for NCX. L-type Ca^{2+} currents and plasmalemmal Ca^{2+} pumps were not affected. Hence, the inhibition of the Ca^{2+} efflux mode of the exchanger could potentially be an effective therapeutic strategy for improving contractility (positive inotropic effect) in congestive heart failure [20].

SEA0400 added to cardioplegia limits the accumulation of intracellular Ca^{2+} during ischemic arrest in isolated cardiomyocytes and prevents myocardial injury and improves the recovery of LV functions in isolated hearts [21]. In an ischemia-reperfusion model, NCX inhibitors have been shown to be cardioprotective [22, 23], and reno-protective [24, 25]. Hence, NCX inhibitors have the potential to be used for the prevention of ischemia-reperfusion injuries.

Figure 4 Example structures of target validation compounds found in the literature.

3 Design strategies for the chemical series

When the programme started, as proof of the target protein idea the compound KR-R 7943 was published, but the Taisho Pharmaceutical NCX1 inhibitor series were not. The initial design strategy was based on the known KB-R series and followed by the adoption of the Taisho Pharmaceutical NCX1 inhibitor series. Later, when virtual screening (VS) methods developed and became faster and sufficiently feasible, 2D VS was utilised in the programme. These two approaches resulted in two distinctively different chemical series for further optimization, namely the flavan and phenyl furan series. These series have very different inherent PK properties which enabled us to address both chronic as well as acute cardiac indications. In the following text these series are called the A and B chemical series, respectively. In total altogether 723 different compounds were synthesized for the programme.

Experimental conditions of the primary *in vitro* screening assay were modified during the term of the programme. Therefore, a set of compounds were rescreened with new assay conditions to validate the comparability of the assays. The activity results mainly followed the same tendency and were comparable, but unfortunately some molecules had multitudes of difference in their results, therefore in **Tables 2 - 4** the activity values of some compounds are presented as bands of activity. The SAR was based on the activity values available at the time. The solubility was classified into three screening solubility classes: insoluble compounds with a lower than 10 µg/ml solubility in a pH 7.4 phosphate buffer (50 mM), moderate soluble compounds with a solubility between 10-100 µg/ml in the buffer and soluble compounds which would have a solubility better than 100 µg/ml in the buffer. Unfortunately none of the compounds fell into the soluble compound class.

In Chapter 3.1.1 hit finding and hit optimization for the A series compounds (flavan) are described and in Chapter 3.1.2 the lead optimization is described for the same series. Chapters 3.2.1 and 3.2.2 describe the same phases for the B series compounds (phenyl furan).

3.1 Design of novel NCX inhibitors for chronic indications

3.1.1 *Discovery of chemical series A*

The first design strategy was based on the KR-R 7943 structure, but the same strategy was quickly adapted for the compound series from Taisho Pharmaceutical as the patent applications were published. KBR-7943 is a linear chain of two aromatic rings connected to each other with one methylene group and an oxygen atom, followed by an aliphatic side

chain with a terminal basic functional group. It has many freely rotating bonds, and hence it can be considered a flexible structure. The design strategy was to learn the molecular features (3 dimensional, steric and electronic) that were necessary for binding NCX1 and for triggering its biological response [26]. Any attempts to shorten the distance between aromatic rings or the basic functional group as well as changing the basic functionality to another basic group failed to improve or maintain the activity of the KB-R 7943. Based on the hypothesis made in the beginning, the optimal conformation for binding was not favoured or it had an overly low energy barrier to other nonbinding conformations, and by restricting the structure, the activity could be improved. It was suggested that the KB-R 7943 could be restricted with an aliphatic ring system as described in **Figure 5**. **Figure 5** presents the design strategies for both KB-R 7943 and 2-phenoxyaniline and phenoxypyridine series patented by Taisho Pharmaceutical. The coloured dashed lines represent different options that were followed during the scaffold hopping. ORM-10113 and ORM-10122 modifications (purple dashed lines) gave inactive derivatives and have a minor influence on the number of rotating bonds. Red and blue dashed line modifications reduce the flexibility. **Table 2** presents a sample of molecules synthesized during the scaffold hopping phase that resulted in ORM-10103 and ORM-10285.

Figure 5 Design strategy for scaffold hopping utilising known NCX inhibitors.

Table 2 Lead hopping based on KB-R 7943 and Taisho Pharmaceuticals compounds incl. SEA400. *IC₅₀ for NCX in Sf9 cells. Intracellular [Ca²⁺] recorded with Fluo-4 dye. Extracellular [Ca²⁺] 20 μM Functional modes are undefined.

Code	Ref.	Structure	NCX1 IC ₅₀ *	Inhib. % at 1 μM	Inhib. % at 10 μM	Inhib. % at 50 μM	Solubility class pH 7.4
KB-R 7943	[6]		3.8-10 μM	3-36%	-	87-90%	moderate
SEA 0400	[7]		376 nM	87-91%	-	95%	insoluble
ORM-10103	I (IV)		176 nM	86%	-	81%	insoluble
ORM-10391	-		-	inactive	-	11%	insoluble
ORM-10285	I		13.6 μM	34%	-	75%	insoluble

Table 2 (continued)

Code	Ref.	Structure	NCX1 IC ₅₀ ^a	Inhib. % at 1 μM	Inhib. % at 10 μM	Inhib. % at 50 μM	Solubility class pH 7.4
ORM-10113	-		-	inactive	-	inactive	moderate
ORM-10122	-		-	inactive	-	13%	moderate
ORM-10353	II		-	24%	-	62%	insoluble
ORM-10541	I		1 μM	76%	-	86%	insoluble
ORM-11393	-		>10 μM	-	-	-	moderate

Table 2 (continued)

Code	Ref.	Structure	NCX1 IC ₅₀ ^a	Inhib. % at 1 μ M	Inhib. % at 10 μ M	Inhib. % at 50 μ M	Solubility class pH 7.4
ORM-10310	I		3.3 μ M	62%	-	88%	insoluble
ORM-10402	I		-	36%	-	66%	moderate
ORM-10458	I		246 nM	69%	-	64%	insoluble
ORM-10442	I		-	38%	-	78%	insoluble
ORM-10575	I		149 nM	83%	90%	-	insoluble

Table 2. (continued)

Code	Ref.	Structure	NCX1 IC ₅₀ ^a	Inhib. % at 1 μ M	Inhib. % at 10 μ M	Inhib. % at 50 μ M	Solubility class pH 7.4
ORM-10490	-		-	inactive	-	43%	moderate
ORM-10551	II		-	37%	60%	-	insoluble
ORM-10673	II		2.8 μ M	65%	71%	-	insoluble

The solubility classes in pH 7.4 phosphate buffer: insoluble less than 10 μ g/ml; moderate solubility between 10-100 μ g/ml; soluble has better than 100 μ g/ml solubility. Terminal hydrogen atoms are implicit in the structures (ActivityBase)

Efforts were focused on the optimization of ORM-10103 as it and its modifications had better activity than ORM-10285 and ORM-10543 including their modifications. In addition, aniline is a known structural alert for carcinogenic property [27, 28]. However, there are exceptions among anilines that do not have the undesired genotoxic property. Restricting the aromatic ether moiety (blue dashed line) of the 2-phenoxyaniline and/or phenoxypyridine series led to flatness and did not improve the activity either. The chromane ring modification was the most favoured (the red dashed line in **Figure 5**). The six membered ring was favoured over the five membered ring as shown when comparing indane derivative ORM-10353 with chromane derivative ORM-10103 or 5,6,7,8-tetrahydronaphthalene derivative ORM-10442. Both modifications: the addition of an exocyclic heteroatom to the chromane and the substitution of benzylic methylene of chromane with a heteroatom (examples ORM-10310, ORM-10490, ORM-10551, ORM-10402 and ORM-10458) to improve solubility were shown to reduce activity. The 2,3-dihydrobenz[*b*][1,4]oxathiin ORM-10458 was an exception, but the solubility was not improved. Because of oxidative metabolism and its ability form sulfoxide and/or sulphone derivatives respectively, sulfur is known for its property of being labile [29]. ORM-10391, an isomer of ORM-10103, was synthesized to understand the three dimensional molecular features that were necessary for binding NCX1. Changing the position of the phenyl substituent from position 2 to position 3 (ORM- 10391) made the compound inactive. 5,6,7,8-Tetrahydronaphthalene derivative ORM-10442 and 3-methyl-2-phenylchromane derivative ORM-10763 displayed reduced activity as well. This led us to the conclusion that the oxygen atom of the chromane had a role in the binding and addition to position 3 and is not favoured. Hence, ORM-10103 was selected for focused optimization. Today this phase is called the lead optimization phase.

3.1.2 Optimization strategy and SAR/SPR of the series A

The starting point for the lead optimization of ORM-10103 showed good inhibitory activity of NCX1 and selectivity towards the L-type calcium channel. However, it was poorly soluble. This unfavourable property limited the possibilities to study all the required properties of the compound and scaffold. Therefore one of the major efforts was to improve solubility. **Figure 6** displays each of the substructure features which were optimized in parallel processes. Beneficial structural features of optimization lines were merged during the optimization to identify whether combining some of them would bring an overall synergistic effect to the NCX1 inhibition activity. The B-ring system (chromane) was

optimized as part of the hit optimization described in Chapter 3.1.1. The lead optimization was focused on optimisation of the A-ring (phenyl) and C- ring (5-nitropyridine) systems.

Many electron withdrawing substituents *e.g.* halogens, nitro, trifluoromethyl and 5-nitropyridyl in the A-ring were tolerated without reducing the activity. The position of the EWG had a minor or no effect on the activity.

Figure 6 The optimization strategy followed during the lead optimisation of the ORM-10103 scaffold.

As expected, the modifications did not improve the solubility, but halogen substitutions, especially fluorine atoms, are often used for blocking metabolism [30]. In addition, EWG might have a minor positive effect on the activity compared to a non-substituted derivative (*e.g.* ORM-10743 vs. ORM-10663). Steric reasons could explain the activity difference between *p*- and *m*-substitutions of 5-nitropyridyl as this large group is favoured by the *m*-position. Instead, all of the tested electron donating substituents reduced activity with the exception of *o*-substituted aniline derivative ORM-10656. The *m*- and *p*- substituted A-ring anilines behaved as the other electron donating groups did, *i.e.* reducing the NCX1 inhibition activity. The unexpected behaviour of ORM-10656 might be due to the repulsion caused by electron pairs of oxygen and nitrogen atoms and favouring distorted conformation. Similar additional positive effects on the activity were not observed with *o*-halogen substitution. ORM-10656 also had better inherent solubility at pH 7.4, but due to its genotoxicity alert the aniline substituent was not accepted in the structure. Replacing the A-ring with a heteroaromatic rings was not a tolerated modification as seen *e.g.* in ORM-10720.

However, the C-ring offered the possibility to add solubilizing groups and improve the solubility of the compounds. The best option for a C-ring turned out to be 2-pyridyl allowing solubilizing groups in position 5. The reduction of 5-nitro to the amino group gave

the easiest and quickest route to explore various solubilizing groups. For example, acetamide (ORM-10564), sulphonamide (ORM-10663 and ORM10743) and 2-amino acetamide (ORM-10787) derivatives retained approximately similar activity to ORM-10103, but with an improved pH 7.4 buffer solubility (50 mM). Slightly acidic sulphonamides (*e.g.* ORM-10663 and ORM10743) and weakly basic 2-amino acetamide like compounds (*e.g.* ORM-10787, ORM-10925, ORM-10962, ORM-11037 and ORM-11169) enabled salt formation and pH dependent solubility while they also have improved the pH 7.4 buffer solubility. Thiourea (ORM-10717) and amino sulphonamide (ORM-12647) derivatives showed reduced NCX1 activity. However, reversed amide derivatives *e.g.* ORM-13326 maintained similar activity compared to substituted 5- amino 2-pyridyls.

Many amino acids were well-tolerated substituents of 5- amino 2-pyridyl that improved solubility. ORM-11037 and ORM-11169 are examples of the amino acid derivatives. ORM-11037 has L-proline and ORM-11169 has D-proline as a solubilizing moiety. Activity towards NCX1 was slightly improved compared to ORM-10103 with the L-amino acid, but the amide bond was metabolically unstable. A D-Amino acid derivative tolerated metabolism better, but led to reduced NCX1 activity. Other amino acids behaved similarly. All 2-amino acetamide type compounds had a liability towards the hERG channel. hERG inhibition has shown to trigger drug-induced QT interval prolongation and the appearance of torsade de pointes (TdP), hence indicating proarrhythmic potential of the molecule [30, 31]. It has been suggested that a margin of 30-fold between hERG IC₅₀ and C_{max} would be adequate to ensure an acceptable degree of safety from arrhythmogenesis, with a low risk of obtaining false positives [31].

In **Table 4** the key *in vitro* parameters of some potential compounds are summarized. ORM-10962 is a novel NCX inhibitor which is the most selective towards the L-type Ca²⁺ channel and hERG channel. It is a very potent inhibitor for NCX in both the reverse and forward functional mode. ORM-10962 has reasonable solubility in solutions with a relevant formulation for *in vivo* dosing. The compound did not raise any major concerns concerning human recombinant CYP450 enzymes.

Plasma protein binding of ORM-10962 was studied in two species, rats and guinea pigs. The compound showed extensive binding to the plasma proteins of both species with 98-99% respectively. An *in vivo* PK study with a dose of 5 mg/kg in guinea pigs after intravenous administration showed that the plasma concentration of ORM-10962 was steady from 30 to 120 minutes. The unbound plasma concentration of ORM-10962 was above IC₅₀ (91 ng/ml) 5 minutes after administration, but not after 15 minutes. However, the local concentration of ORM-10962 and its metabolites in the target tissue were not known/studied. *In vitro* microsomal studies suggested that the main metabolism route was

hydrolysis for all derivatives that are connected to the amino pyridyl with acyl bonds. The formed metabolite ORM-10543 was further metabolized to ORM-10564 via acylation. The formation of glucuronides was also detected in *in vivo* studies. Metabolite identification of ORM.-10962 *in vivo* from dog plasma showed that ORM-10543 was the main metabolite. A minor amount of ORM-10564 as well as two different hydroxylation metabolites could be detected. Based on the PK results, ORM-10962 was stable enough for *in vivo* PD studies.

Table 3 The SAR of the lead scaffold. *IC₅₀ for NCX in SF9 cells. Intracellular [Ca²⁺] recorded with Fluo-4 or Fluo-3 dye. Extracellular [Ca²⁺] varied between 10μm to 1 mM: (a) The functional mode is undefined / (b) To elicit reverse (R) mode NCX activity, an Na⁺-deficient medium was used to dilute 50% of the extracellular Na⁺ level [32].

Code	Ref.	Structure	NCX1 IC ₅₀ *	Inhib. % at 1 μM	Inhib. % at 10 μM	Inhib. % at 50 μM	Solubility class pH 7.4
ORM-10103	I		176 nM	86%	-	81%	insoluble
ORM-10543	I		430 nM	73%	-	93%	moderate
ORM-10564	II		140 nM	86%	89%	-	moderate
ORM-10663	II		203 nM	41%	83%	-	moderate
ORM-10717	II		1.4 μM	69%	91%	-	moderate
ORM-10787	II		86 nM	92%	97%	-	moderate

Table 3 (continued)

Code	Ref.	Structure	NCXI IC ₅₀ ^a	Inhib. % at 1 μM	Inhib. % at 10 μM	Inhib. % at 50 μM	Solubility class pH 7.4
ORM-10925	II		278 nM	78%	88%	-	moderate
ORM-10962	II (V, VI)		197 nM R <30 nM	88%	88%	-	moderate
ORM-11037	II		100 nM	86%	90%	-	moderate
ORM-11169	II		1 μM	79%	-	-	moderate
ORM-10743	II		35 nM	84%	94%	-	moderate
ORM-12647	-		R 335 nM	-	-	-	moderate

Table 3 (continued)

Code	Ref.	Structure	NCX1 IC ₅₀ ^a	Inhib. % at 1 μ M	Inhib. % at 10 μ M	Inhib. % at 50 μ M	Solubility class pH 7.4
ORM-10656	-		70 nM	82%	92%	-	moderate
ORM-10876	II		193 nM	86%	-	-	insoluble
ORM-10751	II		18 nM	91%	88%	-	insoluble
ORM-10797	II		60 nM	77%	87%	-	insoluble
ORM-10720	II		-	-	41%	-	moderate
ORM-13326	-		R 45 nM	-	-	-	-

The solubility classes in pH 7.4 phosphate buffer: insoluble less than 10 μ g/ml; moderate solubility between 10-100 μ g/ml; soluble has better than 100 μ g/ml solubility. Terminal hydrogen atoms are implicit in the structures (ActivityBase)

ORM-10962 fulfils the Lipinski rule of 5, the famous and broadly used thumb rule for drug likeness named after its developer Christopher Lipinski [33]. The number of hydrogen bond acceptors of ORM-10962 is 5 (<10) and number of donors is 2 (<5), the molecular weight of free base is 460 (<500), and predicted logP is less than 5 (ADMET Predictor: 3.3, QikProp: 3.8 MOE: 3.9). Therefore, ORM-10962 was considered to be a suitable candidate for further studies *in vivo* and *ex vivo*.

Table 4 Summary table of key *in vitro* parameters

code	IC ₅₀ NCX1 nM (a)	hERG IC ₅₀ μM	NCX IC ₅₀ inward/outward nM (b)	IC ₅₀ CYPs**	L-type Ca ²⁺ μM	Solubility*** mg/ml
SEA0400	57-376	2.8	57R	-	3.4	Insoluble
ORM-10103	176	-	(780/960)*	-	-	Insoluble
ORM-10564	140	11.3	-	>13 μM	6.9	Moderate
ORM-10656	70	0.84	-	-	-	1.49 GS
ORM-10787	86	1.2	34	-	14	>10.72 GS
ORM-10962	197	3.4	<30 R (55/67)*	>10 μM	11.4	>10.35 GS
ORM-11037	100	1.6	59	>13 μM	6.5	39.1 saline
ORM-11169	1000	2.7	-	>11	9.9	>9.88 GS
ORM-12647	-	7.2	335	-	8.7	Moderate
ORM-13395	-	0.4	64	> 8 μM	5	-

IC₅₀ for NCX in Sf9 cells. Intracellular [Ca²⁺] recorded with Fluo-4 or Fluo-3 dye. The extracellular [Ca²⁺] varied between 10 μM to 1 mM: (a) The functional mode is undefined / (b) To elicit reverse (R) mode NCX activity, an Na⁺-deficient medium was used to dilute 50% of the extracellular Na⁺ level [32]. *Studied in the University of Szeged in Hungary using a patch clamp technique (Inhibition curves in Canine cardiomyocytes for ORM-10103 and ORM-10962 can be found from **paper II figure 4 and paper III figure 3**, respectively. Values are means ± SEM in both papers). ** The value is the lowest IC₅₀ value of all the studied CYP's. The studied human recombinant CYPs were CYP1A2, CYP2C9, CYP2C19, CYP3A4. ***RT, 3h time point, GS = 5% Glucosteril, > = solubility at least, while the max solubility was not known/studied. The solubility classes in pH 7.4 phosphate buffer: insoluble less than 10 μg/ml; moderate solubility between 10-100 μg/ml; soluble has better than 100 μg/ml solubility

3.2 Design of novel NCX inhibitors for acute indications

3.2.1 Discovery of chemical series B

The original hit of series B was discovered utilizing ligand based 2D virtual screening (VS) by the computer aided drug discovery team. Features used for the VS were derived mainly from existing Orion internal NCX1 inhibitor structures such as *e.g.* ORM-10103 and ORM-

10787. The VS campaign resulted in a library of 636 purchased compounds for screening at a concentration of 10 μM . The hit ORM-120407 was identified as a singleton potent structure inhibiting NCX1 by 87% at a concentration of 10 μM . The original purchased library contained 18 molecules containing phenyl furan moiety like the hit molecule. However, none of the other 18 molecules showed any activity for NCX1 at a concentration of 10 μM except ORM-120407. Further characterization of the hit structure showed that the IC_{50} for the NCX1 inhibition of the hit was around 200 nM and it inhibited hERG and L-type Ca^{2+} channels with IC_{50} of 2.1 μM and 3.1 μM respectively. These results indicated that the scaffold has the potential for optimization, but preliminary SAR was lacking. To better explore the SAR of ORM-120407, synthesis of new molecules was needed. In addition, a couple of new similar molecules became commercially available 3 years after ORM-120407 was found to inhibit NCX1.1.

Fig 7 The optimization strategy of ORM-120407 scaffold.

3.2.2 Optimization strategy and SAR/SPR of the series B

Figure 7 displays the substructure features that were optimized in parallel processes. Beneficial structural features of optimization lines were merged during the optimization. In order to identify potential overall synergistic effects for NCX1 inhibition, some of the features were combined. **Table 5** summarizes example derivatives of various 5-membered ring systems and 6-membered ring systems that were synthesized during the B-ring (furan) optimization. SAR tolerated 5-membered ring systems better than 6-membered ring systems. Clear evidence of whether the aromaticity was favoured over the aliphatic ring system remains open due to the fact that the synthesized aliphatic ring system derivatives lack a hydrogen bond acceptor in an optimal position. Ring heteroatom *i.e.* oxygen or nitrogen at position one (of the furan) combined with a 1-, 5- substitution was proven to be critical for the binding, *i.e.* carbon at position 1 abolished activity (*e.g.* ORM-11298, ORM-11863 and ORM-12508 versus ORM-11541 or ORM-11628). This finding indicates that a hydrogen bond acceptor is important at that position. This was also shown with ORM-11725, which has only an aliphatic chain between the A and C rings, but a critical hydrogen bond acceptor (ether) remained. The same phenomenon was seen with a plain phenyl ring as a B-ring (ORM-11628) *vs.* 2,6-pyridyl or 2,4-pyrimidyl derivatives (ORM-11680 and

ORM-12508 respectively). The first of the modifications was inactive while the two latter ones brought the activity back. B-ring optimization provided three potential ring systems that were tolerated; furans, oxazoles and thiazoles. The central B-ring oxazole made it possible for 4,2- and 2,4- substitutions which were both proven to have similar activity ca. 30 nM (i.e. ORM-11298 and ORM-11858 respectively). The 4,2-substituted derivatives were easier to synthesize, so therefore the focus was put mainly on 4,2-substituted oxazole derivatives. Thiazole would in principle provide a similar possibility, but within the given timelines, only the synthesis of a 4,2-substituted derivative was successful (ORM-11676). 4,2-Substituted 1,2,3-triazoles and 5,3-substituted 1,2,4-oxadiazoles (ORM-11782 and ORM-11595 respectively) all had reduced activity for NCX1 inhibition.

The lead optimization focused on the optimization of the A-ring (aniline) and C- ring (phenyl) systems. The starting points for the lead optimization were the original hit molecules ORM-120407 and ORM-11298 (4,2-substituted oxazole ring). Since thiazole derivatives tended to have liability for many CYP 450s, it was considered better to focus on furan and oxazole derivatives. During the optimization, 63 furan and 47 4,2 -oxazole derivatives were synthesized. ORM-120407 already showed good inhibitory activity towards NCX1, but the selectivity towards hERG (2.1 μ M) and L-type calcium (3.1 μ M) channels needed to be improved. ORM-120407 also had poor solubility (<10 μ g/ml). ORM-11298 had better inhibitory activity towards NCX1 and good selectivity towards the hERG channel. The oxazole series had also good selectivity towards the L-type calcium channel *e.g.* ORM-11649. Unfortunately both structural starting points had 3-chloro-4-fluoro aniline structural elements that were an alert for genotoxicity if formed through metabolism. Even though the oxazole derivative seemed to be a better starting point due to its better selectivity, many derivatives were chemically unstable. Therefore, furan remained the central B-ring in the original hit molecule.

Table 6 presents examples of the furan derivatives of chemical series B. The hydrogen bond donor property of the aniline moiety was proven to be important for NCX1 activity in the scaffold during the exploration carried out for replacing the unfavourable structural features. The finding was also proven with tertiary amine derivatives such as ORM-11040 and ORM-11407. Therefore, the strategy was to select substituted anilines from literature that were not reported to be genotoxic or pose weak genotoxicity [34, 35]. It was unclear at the optimization phase, if the bond between the aniline and methylene would be cleaved during metabolism and whether any unsubstituted aniline would be formed or how much would be formed. Yet another approach to avoid aniline was to reverse the amine bond, but these derivatives (ORM-11565 and ORM-11630) totally lost the activity needed for NCX1 inhibition. This could be due to increased basicity of molecules. All other attempts to avoid aniline failed as well. For example replacing aniline with a benzylic group, or the use of a

carbon linker (ORM-11382), ethers and esters also produced inactive compounds. The use of 2-aminopyridine instead of aniline was not tolerated either.

Table 5 Exploration of various B-ring systems

Code	Structure	IC ₅₀ NCX1 nM	IC ₅₀ hERG	IC ₅₀ Cyp's	Solubility class
ORM-120407		207-231 53 R	2.1 μM	-	-
ORM-11298		31	>10 μM	2C19	moderate
ORM-11319		inactive	-	-	-
ORM-11411		inactive	-	-	-
ORM-11451		inactive	-	-	-
ORM-11565		inactive	-	-	-
ORM-11595		800	-	>6.9 μM	-
ORM-11628		inactive	-	-	-
ORM-11630		inactive	-	-	-
ORM-11676		10	5.2	many	moderate
ORM-11680		3000	-	-	moderate

Table 5 (continued)

Code	Structure	IC ₅₀ NCX1 nM	IC ₅₀ hERG	IC ₅₀ Cyp's	Solubility class
ORM-11725		800	-	-	moderate
ORM-11755		inactive	-	-	-
ORM-11763		inactive	-	-	-
ORM-11782		175	-	>10 μM	moderate
ORM-11858		35	-	-	Moderate
ORM-11863		2.5	-	>3.5 μM	Insoluble
ORM-12508		181 R	-	>6.3 μM	Moderate

The solubility classes in pH 7.4 phosphate buffer: insoluble less than 10 μg/ml; moderate solubility between 10-100 μg/ml; soluble has better than 100 μg/ml solubility. Terminal hydrogen atoms are implicit in the structures (ActivityBase)

C-Ring optimization included various derivatives with different substitution patterns. Representative sample of synthesized derivatives are presented in **Tables 5** and **6**. A shift of the nitro group of the C-ring from the *o*-position to *m*-position (ORM-11023) improved NCX1 inhibition activity slightly, but the *p*-position (ORM-11066) was not tolerated leading to 10-fold reduced NCX1 inhibition activity. Unlike the nitro group, *p*-methoxy substitution was better tolerated resulting in only a slight reduction in NCX1 inhibition activity. Reduction of the nitro group at the *m*-position (ORM-11065) improved NCX1 inhibition activity 10-fold. Similar phenomena with the *o*-nitro group remained unexplored as the compound was shown to be labile and the compound was not screened. Acylation and urea modification of formed aniline produced inactive compounds. Replacing *o*-, *m*- and *p*-nitro with halogen respectively was a well-tolerated modification. Especially the *ortho* position was favoured and resulted in improved activity. However, as expected, the influence on solubility was minor. In contrast to the *o*-chloro derivative, the ORM-11165 trifluoromethyl modification was not tolerated at all and produced an inactive compound. The trifluoromethyl group was better tolerated in the *m*-position, but still led to reduced NCX1 inhibition. The methyl group as such was better tolerated especially in the *m*-

position. This finding indicates that the trifluoromethyl group was too large a C-ring substituent to retain the binding to the NCX1 protein. Instead, compounds without any substituent in the C-ring *e.g.* ORM-11619 showed good NCX1 inhibition activity. In conclusion, substitution of the C-ring phenyl was not necessary, but some substituents were tolerated, *e.g.* aniline and chlorine. Replacing the phenyl with different heteroaromatic ring systems was mainly tolerated. In **Table 6** ORM-11817, ORM-11875, ORM-11907, ORM-11905 and ORM-12509 represent C-ring hetero aromatic derivatives. Especially the *ortho* pyridyl derivative showed good NCX1 inhibitory activity. The free electron pair in nitrogen can conjugate with the free electron pairs of the oxygen atom of furan which increases the flatness of the structure resulting in poor solubility.

As described earlier in this chapter the original hit molecule ORM-120407 showed an alert for genotoxicity and attempts to replace the aniline moiety with non-aniline structural moiety failed [27, 28]. Hence, the SAR of the A-ring aniline was explored. It was also unknown, if any dealkylation of the aniline would occur *in vivo*. ORM-11024 (aniline), ORM-11817 (3,4-difluoroaniline), ORM-11382 (4-fluoroaniline), ORM-11590 (3-nitroaniline) and ORM-11408 (3-chloroaniline) are illustrative examples of synthesized modifications to explore the SAR of the A-ring aniline. Naked aniline showed a slight decrease in NCX1 inhibition activity. The addition of any polar substituent *e.g.* carboxyl acid were not tolerated, but halogen substitution on the *p*- and *m*- positions were tolerated and resulted in improved NCX1 inhibition. However, neither position was favoured over the other nor showed any synergistic or additive effect. Both chlorine and fluorine were tolerated in both positions and had a similar potency for NCX1 inhibition. The differentiating property was found in the selectivity towards the hERG channel. Inhibition of human recombinant CYP450 enzymes remained a challenge with all the modifications of the B-series molecules. Since the B-series was developed for the short term, acute use with fast metabolism and relatively low CYP450 activities were not considered as no-go properties. ORM-11372 fulfils the Lipinski rule of 5: the number of hydrogen bond acceptors is 1 (<10) and donors is 3 (<5), molecular weight of free base is 282 (<500), and predicted logP is less than 5 (ADMET Predictor: 3.95, MOE: 3.413, QikProp: 3.974). Therefore ORM-11372 was selected for further studies. However, optimization of the B-series remained unfinished due to the termination of the project and cardiovascular research at Orion.

Figure 8 shows the structural features that were found to be critical for the NCX1 inhibition activity. There are still some unexplored opportunities for replacing the aniline as a hydrogen bond donor, *e.g.* indoline or benzimidazole derivatives might solve the issue. There is still room to further explore the heterocyclic C-ring systems for instance by adding solubilizing substituents. For example ORM-11817 has good NCX1 inhibition activity, but

poor solubility. Adding an amino group to the *m*- or *p*-position might lead to interesting derivatives.

Figure 8 2D Pharmacophore of the B-series.

Table 6 A and C-ring system optimization of the B-series.

Code	Structure	IC ₅₀ NCX1 nM	IC ₅₀ hERG	IC ₅₀ Cyp's	Solubility class
ORM-11023		156	-	-	insoluble
ORM-11024		875	-	-	moderate
ORM-11066		2900	-	-	insoluble
ORM-11040		inactive	-	-	insoluble
ORM-11407		714	-	-	-
ORM-11382		2846	-	-	insoluble
ORM-11408		inactive	-	-	-
ORM-11058		inactive	-	-	insoluble
ORM-11271		inactive	-	-	moderate

Table 6 (continued)

Code	Structure	IC ₅₀ NCX1 nM	IC ₅₀ hERG	IC ₅₀ Cyp's	Solubility class
ORM-11065		6 <3 R	2.2 μM	many	moderate
ORM-11904		9.4	-	-	moderate
ORM-11165		4	9 μM	many	insoluble
ORM-11616		80	-	many	moderate
ORM-11190		204	-	many	-
ORM-11217		210	14.6 μM	-	moderate
ORM-11192		inactive	-	-	-
ORM-11243		370	-	-	insoluble
ORM-11244		inactive	-	-	insoluble
ORM-11332		inactive	-	-	moderate
ORM-11372		6 6R	10.6 μM	many	moderate
ORM-11408		3	6.3 μM	many	moderate
ORM-11482		470	-	-	moderate

Table 6 (continued)

Code	Structure	IC ₅₀ NCX1 nM	IC ₅₀ hERG	IC ₅₀ Cyp's	Solubility class
ORM-11590		75	-	many	moderate
ORM-11619		15 11R	9.5 μ M	>4.5 μ M	moderate
ORM-11685		10	3.2 μ M	many	moderate
ORM-11817		3	2.4 μ M	>3.4 μ M	insoluble
ORM-11875		90	-	many	moderate
ORM-11907		38	-	many	moderate
ORM-11905		60	-	-	moderate
ORM-12509		20R	-	2C19	moderate

The solubility classes in pH 7.4 phosphate buffer: insoluble less than 10 μ g/ml; moderate solubility between 10-100 μ g/ml; soluble has better than 100 μ g/ml solubility. Terminal hydrogen atoms are implicit in the structures (ActivityBase).

3.3 Summary

Taking into consideration the shortcomings of the traditional approaches at the time of early phases of the study, the optimization of the chemical series aimed to follow the concept of chemical biology and multiparameter optimization approach that was described by Wess in 2002 [36]. However, the feedback loop from the molecule profiling was sometimes long and some *in vitro* methods were not always available or in standard use in Orion. For example, during the optimization of the A-series (flavans) the *in vitro* metabolic stability method was not in use and metabolic stability was studied *in vivo*. Hence, comparable data for ORM-10962 and ORM-11372 is not fully available. However, both ORM-10962 and ORM-11372 have followed the broad outline of the process shown in **Figure 9**.

The starting points for both series (ORM-10103 and ORM-120407) had relatively good NCX1.1 inhibition activity and there was room to explore novel chemical space. However, other parameters were still suboptimal for the planned indication. **Figures 10** and **11** present the profiles of ORM-10962 and ORM-11372, the end points of the optimization. Ideal values for each parameter are shown by the strong green colour of the sector in the outer circle. Parameters that matched the ideal value by 80% are shown in strong green in the inner circle. The light coloured green sector shows values that were acceptable, but not optimal or could be improved. The closer the center the sector of the pie is, the further the value is from the ideal value. If the value of the parameter should be seen as an alert, the sector is coloured red. The sector is shown as grey if the data point is missing.

Figure 9 Example of a multi parameter optimization of a lead structure [36].

ORM-10103 was a novel structure at the time of invention and also a novel NCX1.1 inhibitor. However, it had poor solubility which prevented further characterization of the compound. It had modest inhibition activity for reverse and forward functional modes of NCX1.1. ORM-10962 has similar NCX 1.1 screening inhibition activity, but improved inhibition activity for reverse and forward functional modes of the NCX1.1 and clearly improved solubility in the buffer (50 mM) at pH 7.4, and a relevant formulation (solution) was developed for *in vivo* dosing. **Table 7** summarizes the parameters studied for ORM-10962 and the profile is shown in **Figure 10**. *In vitro* cardiac ion channel pharmacology studies showed that ORM-10962 is selective towards the L-type Ca^{2+} channel and hERG channel. In addition, passive permeation was predicted to be high. The diversity profile in the receptor panel (purchased from Cerep) revealed that MAO A and B binding should be followed. Potential risks related to the MAO inhibitors would include symptoms such as

tachycardia, increased blood pressure and fever [37]. However, ORM-10962 showed mild to moderate binding to MAO A and B only at high concentrations.

Figure 10 Profile of ORM-10962.

Figure 11 Profile of ORM-11372.

ORM-10962 did not show any major concerns with human recombinant CYP450 enzymes or measured toxicity parameters. However, some data points are missing as the business decision was to focus on acute short-term treatments with the short half-life of the compound. Plasma protein binding, solubility and hERG selectivity might need further optimization if the optimization of the compound series would be continued. Sanofi has continued the optimization of the A series as shown in **Table 1**. They have evidently aimed to improve the solubility by reducing the flatness of the scaffold and replaced the pyridyl ring with an aliphatic ring. This structural change should in theory improve the solubility and it might also have a positive effect on hERG selectivity. However, these parameters have not been measured in Orion from NCX1.1 inhibitors patented by Sanofi. Thus, they can be considered speculative and we do not have experimental evidence.

Table 7 ORM-10962 summary table.

ORM-10962			
IC ₅₀ NCX1	197 nM	IC ₅₀ L-type Ca ²⁺ channel	11.4 μM
IC ₅₀ NCX1 reverse	<30nM (55/67 nM) (Paper III)*	IC ₅₀ NaV 1.5 channel	-
Solubility 5% GS	10.35 mg/ml	IC ₅₀ hERG	3.4 μM
Caco-2 permeability	high	IC ₅₀ 5-HT2B	-
PPB unbound	1-2 %	Uncoupling	>50 μM
Met.stab. IV dosing		MAO A at 10 μM	11%
Dog in vivo	Stable until 210 min	MAO B at 10 μM	52%
Guinea pig in vivo	Stable between 30-120 min		
CYPs	>10 μM	Mini MLA	Negative
IC ₅₀ Alpha 1A	-	Ames	-

* Studied at the University of Szeged in Hungary using a patch clamp technique for canine ventricular cardiomyocytes (Inhibition curve for ORM-10962 can be found from **Paper III, figure 3**. Values are means ± SEM in both papers)

ORM-120407 was an already known and commercially available compound at the time of the experimental phase of this study. It displays similar inhibition activity for NCX1.1 as ORM-10962, but the solubility is poor in a buffer solution at pH 7.4 and the ratio to the hERG inhibition is lower than with ORM-10962. It also shows an alert towards genotoxicity in its structure (aniline moiety). However, ORM-120407 is novel as the NCX1.1 inhibitor structure and had good potential for chemical optimization. The studied parameters of the lead molecule ORM-11372 of the B series are presented in **Table 8** and its profile is shown in **Figure 11**. ORM-11372 is very potent and has low nM NCX1.1 inhibition activity. *In vitro* cardiac ion channel pharmacology studies showed that it has excellent selectivity towards hERG, L-type Ca^{2+} and NaV 1.5 channels. Selectivity towards the alpha 1A receptor was also good. It possesses high passive permeation and has no major concerns regarding cytotoxicity. *In vitro* Mini MLA as well as Ames tests were both negative. In addition, the *in vitro* clearance is fast, and is hence ideal for acute *iv* dosing for on/off use.

Table 8 ORM-11372

ORM-11372			
IC₅₀ NCX1	6 nM (5R/6F)*	IC₅₀ L-type Ca^{2+} channel	6.1 μM
Solubility mQw	5.67 mg/ml	IC₅₀ NaV 1.5 channel	>30 μM
Solubility saline	5.27 mg/ml	IC₅₀ hERG	10.6 μM
Solubility 5% GS	5.44 mg/ml	IC₅₀ 5-HT2B	Negative
Caco-2 permeability	High	MAO A at 10 μM	29%
PPB unbound human	4%	MAO B at 10 μM	77%
Clearance at 1 μM Human/Rat Hepatocytes <i>in vitro</i>	63/161 $\mu\text{l}/\text{min}/\text{unit}$	cytotox	>100 μM
CYPs	1A2,2C19,3A4	Mini MLA	Negative
IC₅₀ Alpha 1A	>10 μM	Ames	Negative

*Inhibition curve for ORM-11372 can be found from **Paper I, figure 3**. Individual values are presented in the figure according to current British Journal of Pharmacology guidelines. For each measured concentration standard error of inhibition value is less than $\pm 3.5\%$ for reverse mode and less than $\pm 6\%$ for forward mode.

ORM-11372 has a liability for human recombinant CYP450 enzymes. However in the planned setting of *iv* use and fast clearance, the liability is considered to be controllable. Liability has to be followed anyway. The free fraction in plasma was 4% which might be enough for the *in vivo* effect, but the solubility might need further optimization especially for *iv* dosing. The major alert (aniline) for genotoxicity remained, but *in vitro* screening

gave negative results and it was considered acceptable to continue further profiling of ORM-11372. Series B optimization was not finalized due to the business decision to stop cardiovascular research in Orion. There was still room left for chemical optimization as well as the possibility to improve solubility and avoid aniline structural moiety as speculated in Chapter 3.2.2, if the optimization of the compound series would be continued.

Figure 12. A few potential derivatives (and their combinations) that might solve remaining undesired properties of series B, but were not synthesized during the study.

4 Materials and methods

4.1 Synthesis routes for series A and B

The synthesis routes and experimental data for compounds that were patented by Otsomaa et al. are described in detail in **Appendices 01-03**. The objective of **Appendix 01** was to protect the chemical space of active scaffolds found during lead hopping and **Appendix 02** aimed to protect chemical space of series A NCX1 inhibitors that were not included in the patent scope of **Appendix 01**. The scope of **Appendix 03** is to protect the chemical space of the prepared compounds of series B. *Scheme 1* shows the general route for the core scaffold of series A compounds. However, the routes for the other compounds included in **Appendix 01** can be found in the original patent application appended. *Scheme 2* shows a general route for the core scaffolds of the series B compounds, respectively. However, routes for the other compounds included in **Appendix 03** can be found in the original patent application appended.

4.1.1 Series A route

Scheme 1 General formula of the synthesis route for series A compounds
(details in appendices 01 and 03).

6-Hydroxyflavanone derivatives were synthesized from 2',5'-dihydroxyacetophenone or the corresponding propiophenone and an appropriate benzaldehyde via condensation resulting in a mixture of the desired 6-hydroxyflavanone and the corresponding chalcone [38, 39]. The chalcone was further cyclized to the desired flavanone by treating it with sodium acetate. 2-Phenylchroman-4,6-diol derivatives were obtained from the corresponding 6-hydroxyflavanones by reduction as shown in **Scheme 1** above. These diol

derivatives further reduced to 6-hydroxy flavanes. Pyridyl-2-oxy derivatives were obtained by reactions with suitable halopyridines and aromatic phenols. Aminopyridinyloxy derivatives were obtained from the corresponding nitro pyridine-2-yloxy by reduction. The amino pyridine derivatives were reacted with suitable amino acids and other carboxylic acid derivatives using 1-(dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride as a coupling agent to result in the amide derivatives as shown in **Scheme 1**. Optionally, the amide derivatives could be obtained by well-known acylation methods. Protecting groups were removed if needed. ORM-10962 was synthesized from a 3-aminopyridine derivative by acylation with 2-chloroacetyl chloride to form a 2-chloroacetamide derivative that further reacted with piperidin-4-ol.

4.1.2 Series B route

Scheme 2 General formula of synthesis routes for the series B compounds (Details in appendix 03).

B-series compounds were made using four alternative routes (**Scheme 2**). The reaction of route 1 was carried out in the presence of a base such as potassium carbonate, in a solvent e.g. water at elevated temperature. The products were isolated from the reaction mixture for example by extraction with a suitable solvent e.g. ethyl acetate and by evaporation of the solvent. Route 2 could be used for compounds wherein ring A is a furan or 4,2 substituted oxazole and ring B is an *o*-/*m*- substituted phenyl. Aniline derivatives were coupled with carboxylic acid derivatives to obtain a carboxamide derivative as shown in **Scheme 2**. The carbonyl group of the intermediate was reduced to obtain the target molecule of the B-series. Condensation between the aniline derivative and the carboxylic acid derivative presented in route 3 was carried out as in route 2. The reaction between

intermediate and ring B was carried out in the presence of *n*-butyllithium, zinc chloride and tetrakis(triphenylphosphine) palladium in a suitable solvent e.g. tetrahydrofuran. Compounds of formula (I) where the ring A is a furan ring could also be prepared according to route 4 with reductive amination, wherein R₁ is H, halogen, cyano or nitro and R₂ is H or halogen. The reaction between the aniline derivative and the carbaldehyde derivative was carried out in the presence a strong acid such as *p*-toluenesulfonic acid in a suitable solvent at elevated temperature. The imine intermediate was reduced using a suitable reducing agent e.g. NaBH₄ in a suitable solvent such as ethanol. The starting materials used in the processes were either commercially available or prepared using synthetic routes found in the literature [40-44]. ORM-11372 was synthesized via reductive amination of 5-(3-nitrophenyl)furan-2-carbaldehyde with the corresponding fluoroaniline followed by hydrogenation of the nitro group under mild condition (route 4).

4.2 *In vitro*

In vitro assays used for profiling of compounds during the optimization cycles are described below. NCX1.1 inhibition activity screening assay conditions can be found from Appendix 03. Modification of the conditions were carried out in the early days of the study. ADME screening and other miscellaneous assays used during the optimization were standardized methods and the conditions are also described below.

4.2.1 *NCX1.1 inhibition activity assays*

General description of the NCX1.1 inhibition assay and the conditions used during series B optimization phase can be found below and also in Appendix 03. The modified assay conditions used for series A optimization (Appendices 01 and 02) are in brackets in the following description. For the measurement of screening the NCX1.1 inhibition activity, NCX1.1 plasmid together with selection resistance gene was transfected into Sf9 cells. Positive clones were selected for the cultures. The cells were loaded with membrane permeable Fluo-4-AM (or Fluo-3FF) which is broken down inside the cell by esterases to give free Fluo-4 (or Fluo-3). The loaded cells were plated onto 96-well plates loaded with the correct amount of each study compound, (standard KB-R7943 or SEA0400) and blanks (without inhibitor). Typically 150 – 200,000 cells were loaded into each well at a volume of 50 µl. The cells were preincubated with intracellular calcium dye Fluo-4 and test compounds for about 1 h at room temperature before the experiments. Using the FLEXstation protocol, fluorescence was measured at 1.5 s intervals, and at 17s from the start a Na⁺ free solution was added to reverse the NCX-mediated Ca²⁺ transport. A normal extracellular solution contained: 172 mM NaCl, 10 mM HEPES, 1 mM CaCl₂, 1.2 mM MgCl₂, 0.33 mM NaH₂PO₄, 5 mM glucose and 5 mM probenecid. The pH was adjusted to

7.4 with NaOH. The Na⁺-free extracellular solution contained: 141 mM NMDG, 10 mM HEPES, 1 mM CaCl₂, 1.2 MgCl₂, 0.33 NaH₂PO₄ and 5 mM glucose. The pH was adjusted to 7.4 with HCl. The osmolality of both solutions was adjusted to 340 - 355 mOsm in order to match the osmolality of Sf9 cell media. Dilution of extracellular Na⁺ induced robust and reproducible intracellular calcium elevations in human NCX expressing Sf9 cells. On each plate, the IC₅₀ determination of test compounds was based on relative fluorescence changes in comparison to a control and a 5 mM nickel acetate-induced NCX inhibition. Briefly, reverse-mode NCX activity was evoked with a 50 % dilution of extracellular Na⁺ using an internal pipettor and FLEXstation fluorescence imaging plate reader (Molecular Devices, Sunnyvale, CA, USA) in 96-well plates while the intracellular calcium level was simultaneously monitored. To elicit reverse mode NCX activity, an Na⁺-deficient medium was used [32].

All the data points were exported and the NCX exchanger rate was calculated using an in-house Orion developed software program. The data was plotted and IC₅₀ calculations were solved graphically using the Hill equation. In older protocols (Victor) fluorescence was measured 13 times at 40 s intervals, lysed with 10% Triton X-100. The data was collected in an Excel sheet, and entered in the ActivityBase DB computer software and the inhibition percent and IC₅₀ were calculated using ActivityBase.

The measurement of the NCX current (**Paper III**) involved measuring the Na⁺/Ca²⁺ exchanger current (I_{NCX}) in Canine ventricular cardiomyocytes, and the method by Hobai *et al.* was applied for this purpose [45]. The electrical properties of canine ventricular cells most resemble those of human ones [46]. Accordingly, the NCX current is defined as an Ni²⁺-sensitive current and is measured in a special K⁺-free solution (composition in mM: NaCl 135, CsCl 10, CaCl₂ 1, MgCl₂ 1, BaCl₂ 0.2, NaH₂PO₄ 0.33, TEACl 10, HEPES 10, glucose 10 and ouabain 20 μM, nisoldipine 1 μM, and lidocaine 50 μM, at pH 7.4) as described in **Paper II** in detail. The pipette solution used for recording the I_{NCX} contained (in mM) CsOH 140, aspartic acid 75, TEACl 20, MgATP 5, HEPES 10, NaCl 20, EGTA 20 and CaCl₂ 10, pH adjusted to 7.2 with CsOH.

4.2.2 Cardiac safety ion channel assays

To carry out the IC₅₀ measurements of the hNav1.5 channel (the primary cardiac sodium channel, encoded by the SCN5A gene) in-house HEK cells were transiently transfected with cDNA of human SCN5A. These cells were then monitored for the next 1-3 days with a manual patch clamp. Sodium currents were evoked with a depolarizing test step to -15 mV for 10ms following a preceding hyperpolarization step to -120 mV. The voltage protocol was repeated every second. The peak current values during the test pulse at -15 mV were used for analysis.

To take the IC₅₀ measurements of the human L-type Ca²⁺ channel (Fluorescence assay), IMR-32 human neuroblastoma cells (from American Type Culture Collection, CCL-127) were maintained in Eagle's minimum essential medium, supplemented with 10% foetal bovine serum (heat inactivated), MEM non-essential amino acids, 100 I.U penicillin and 100 µg/ml streptomycin, 25 mM HEPES at 37°C in a 5% CO₂/ 95% air atmosphere. Twice a week the cells were trypsinized with 0.05 % trypsin-EDTA solution to remove adhered cells and were re-plated. The growth medium was changed every other or on the third day. For the study, the cells were harvested by trypsinization and centrifuged twice (5 min at 1,000 rpm) and resuspended in a probenecid-Ringer. This Ringer consisted of (in mM): 150 NaCl, 3 KCl, 1.2 MgCl₂, 1 CaCl₂, 20 HEPES, 5 glucose and 2.5 probenecid (pH 7.4 adjusted with NaOH, osmolarity 320-324 mOsm). The cell density was determined, and the volume was adjusted so that there would be 200-250,000 cells per well of a 96-well plate. After centrifugation, the pellet was resuspended in either calcium 5 dye (Molecular Devices, USA) and incubated for 45 min or suspended in 0.04% Pluronic F-127 + Fluo-4 (Molecular Probes, USA) and incubated for 30 min, after which the Fluo-4 was removed with centrifugation and resuspension in probenecid Ringer solution. Incubation for both dyes took place in the dark, at room temperature and with constant agitation. The cell suspension (75 µl per well) was pipetted into a black walled, clear bottom 96-well plate to which the compound (75 µl per well at 2,667 x the final concentration) had already been added. The well plate was centrifuged once (5 min, 1,000 rpm) ensuring that the cells were located on the bottom of the wells. Changes in the intracellular calcium concentration were measured with a FLIPRTETRA fluorometric imaging plate reader (Molecular Devices, USA). The excitation wavelength was 470-495 nm and the emission was measured at 515-575 nm. A KCl Ringer (containing (in mM): 200 KCl, 20 CaCl₂, 1.2 MgCl₂, 20 HEPES, 2.5 Probenecid with pH 7.4 KOH adjusted and osmolarity 320-324 mOsm) was applied by FLIPR (50 µl per well) after 10 s. All the experiments were performed at 37 °C. The fluorescence value and maximum minus baseline were calculated for each well with the ScreenWorks 2.0 software. Dose-response curves for the ORM compounds were typically constructed from a mean of 7 separate wells at each of the 5 different concentrations on a plate. The results are expressed as mean ± s.e.m. for n number of plates.

IC₅₀ human ERG channel patch-clamp measurements assay: HEK 293 cells were maintained for 1-2 days in 6 well dishes in an incubator at 37°C, in a 5% CO₂/95% air atmosphere. The growth medium was Dulbeccos modified Eagle's medium, supplemented with 10% foetal bovine serum (heat inactivated), 100 I.U penicillin and 100 µg/ml streptomycin, 10 mM HEPES, 0.5 mg/ml geneticin. All the experiments were performed at the physiological temperature of 36 ± 1°C. Tight-seal whole cell voltage-clamp recordings were made with an Axopatch 200B amplifier (Axon Instruments, USA) [47].

The coverslip with the adhered cells was placed in an extracellular solution which consisted of (in mM): 143 NaCl, 4 KCl, 1.8 CaCl₂, 1.2 MgCl₂, 5 glucose, and 10 HEPES (pH 7.4 adjusted with NaOH). The osmolarity was adjusted to 301 ± 3 mOsm. All drugs were dissolved in the extracellular solution. All doses were applied for about 5 min. The intracellular solution consisted of (in mM): 130 KCl, 7 NaCl, 5 EGTA, 1 MgCl₂, 5 Adenosine triphosphate (disodium salt), and 5 HEPES (pH 7.2 adjusted with KOH) and the osmolarity was carefully adjusted to 290 ± 3 mOsm with an osmometer. The pipettes had resistances between 1 and 2 M Ω . Only recordings with a series resistances below 10 M Ω were used, and the typical values were 2-4 M Ω . In all cells, the cell capacitance was compensated for by the circuitry in the amplifier. No series resistance compensation was used. In all the recordings, the cell voltage was clamped to -75 mV except during the voltage steps. Voltage steps with a duration of 500 ms from a holding potential of -75 mV were applied first to +10 mV and then to -40 mV at a frequency of 0.1 Hz. Tail current values at -40 mV were used for analysis.

4.2.3 ADME assays

Human recombinant cytochrome P450 (CYP) IC₅₀ determinations (fluorescence assay). The human recombinant CYP's studied were CYP1A2, CYP2C9, CYP2C19, CYP3A4. The test substances (stock solutions 10 and 20 mM or stock solutions 5 and 10 mM) were dissolved in methanol. Different stock solutions were used for inhibition tests with different CYP enzymes according to the solubility of each compound and the required test substance concentration for each CYP enzymes. The CYP enzyme inhibition screening was performed using BD Gentest™ Human P450 Inhibition Screening Kits (BD Biosciences Discovery Labware Worldwide, Bedford, MA, USA) or separately ordered reagents from the same vendor. The incubations were done according to the instructions in the BD Gentest™ Human P450 Inhibition Screening Kits for each CYP enzyme studied. The incubation and preparation of incubation mixtures were carried out on black 96-well FluoroNunc plates (cat.no. 237107, Nunc, Thermo Fisher Scientific Inc, Waltham, MA, USA) with Hamilton Microlab STAR automated liquid handling system. The samples were incubated in a Cytomat 2C-LIN incubator at +37 °C without shaking. After incubation, the reactions were stopped according to the suppliers' instructions. The set of incubations consisted of blank, control and actual reactions. The blank and control reactions did not contain test substances, and in the blank reactions the enzyme and the substrate were added after incubation. The samples were analyzed using an EnVision Xcite multilabel plate reader integrated with a Hamilton Microlab STAR. The fluorescence in each sample was determined and the results were converted to a spread sheet format to enable the calculation of the IC₅₀ values. The results were calculated using the XLfit version 4.3.2 add-in module (ID Business Solutions Ltd., Guildford, Surrey, U.K.) in MS Office Excel.

Metabolic stability *in vitro* was tested with a determination of intrinsic metabolic clearance in *in vitro* preparation. Cryopreserved hepatocytes from rats and humans were obtained from Bioreclamation IVT (Baltimore, MD, USA). Thawing the hepatocytes was performed according to the instructions provided by the supplier. Metabolic activities as determined by model metabolic reactions were provided by the supplier. The samples for metabolic stability (for CL_{int} estimation) and for metabolite identification were prepared using the final proportion of DMSO per incubation of 0.5% (v/v). The final incubation concentration of viable cells was 1 million/ml and the final incubation concentration of the test substance was 0.1 and/or 1 μ M with one to three parallel tests. The number of incubation time points was 2 to 6 (e.g. 0, 20, 40, 60, 90, 120 min) and the incubation was terminated with acetonitrile. The results are expressed as μ l/min/unit (unit is 1 million cells). The metabolic stability of the incubations was assessed using LC-MS. The disappearance of the test compound during the time-course of the incubations was followed by measuring the parent peak areas in the positive ion mode.

For the permeability assay, caco-2 cells (clone C2BBel) were obtained from the American Type Culture Collection (Manassas, VA). Cell monolayers were grown to confluence on collagen-coated, microporous, polycarbonate membranes on 12-well Costar Transwell plates. Details of the plates and their certification are shown below. The permeability assay buffer was a Hanks' balanced salt solution containing 10 mM HEPES and 15 mM glucose at a pH of 7.4. The buffer in the receiver chamber also contained 1% bovine serum albumin. The dosing solution concentration was 1 μ M of the test article in the assay buffer. Cell monolayers were dosed on the apical side (A-to-B) or basolateral side (B-to-A) and incubated at 37°C with 5% CO₂ in a humidified incubator. Samples were taken from the donor and receiver chambers after 120 minutes. Each determination was performed in duplicate. The flux of Lucifer yellow was also measured post-experimentally for each monolayer to ensure no damage was inflicted on the cell monolayers during the flux period. All the samples were assayed by LC-MS/MS using electrospray ionization. The apparent permeability (P_{app}) and percent recovery were calculated as follows:

$$P_{app} = (dC_r/dt) \times V_r / (A \times C_A) \quad (1)$$

$$\text{Percent Recovery} = 100 \times ((V_r \times C_r^{\text{final}}) + (V_d \times C_d^{\text{final}})) / (V_d \times C_N) \quad (2)$$

Where,

dC_r/dt is the slope of the cumulative concentration in the receiver compartment versus time in μ M s⁻¹;

V_r is the volume of the receiver compartment in cm³;

V_d is the volume of the donor compartment in cm³;

A is the area of the insert (1.13 cm² for the 12-well Transwell);

C_A is the average of the nominal dosing concentration and the measured 120-minute donor concentration in μM ;

C_N is the nominal concentration of the dosing solution in μM ;

C_r^{final} is the cumulative receiver concentration in μM at the end of the incubation period;

C_d^{final} is the concentration of the donor in μM at the end of the incubation period.

Efflux ratio (ER) is defined as $P_{\text{app}}(\text{B-to-A}) / P_{\text{app}}(\text{A-to-B})$.

The solubility class was determined by nephelometric screening with a pH 7.4 phosphate buffer (50 mM) solution. Study compounds were dissolved in DMSO at a concentration of 50 mM before solubility measurements were taken. The solubility was given as μg per ml followed by classification of the solubility class. Insoluble and moderate refers to the solubility class described on page 8.

Solubility to 5% glucose solution (5% GS, GlucoSteril®) was studied by dissolving the study sample in a mixture of 561 mg of Macrogol 300(PEG) and 127 mg of propylene glycol solution at RT. 5% GS was added until the total volume was 1 ml. The saturated solution was shaken for 3h at RT followed by filtration and determination of the maximum solubility by HPLC (UV detection).

4.2.4 Other miscellaneous assays

For the evaluation of the relevant anti targets a 5-HT-2B receptor/human recombinant assay was conducted. Chinese hamster ovary (CHO-K1) cells stably transfected with human serotonergic 5-HT2B receptors (from Euroscreen, Belgium) were used in this study. Cells were maintained at 37°C in a 5 % CO₂/95 % air atmosphere in an HAM F-12 medium supplemented with 10 % FCS, 25 mM HEPES, 100 I.U/ml penicillin, 100 $\mu\text{g}/\text{ml}$ streptomycin, 500 $\mu\text{g}/\text{ml}$ geneticin and 250 $\mu\text{g}/\text{ml}$ zeocin. The cells were sub cultured 1:20 – 1:30 twice weekly with 0.25 % trypsin and 1 mM EDTA. The growth medium was changed every other or third day.

The day before the experiment the cells were plated in black-walled, clear bottom 96-well plates at a density of 40,000 cells per well with a modified version of the growth medium. The growth medium was removed, and the cells were incubated with calcium 5 dye (Molecular Devices, USA) for 1 hour at 37°C in the dark. Changes in the intracellular calcium concentration were measured with a FLIPRTETRA fluorometric imaging plate reader (Molecular Devices, USA). The excitation wavelength was 470-495 nm and the emission was measured at 515-575 nm. All experiments were performed at 37 °C.

The compound was dissolved in dimethyl sulphoxide (DMSO) to obtain a stock solution of 10 mM from which the final concentrations were prepared daily by dilution with probenecid-Ringer. This Ringer consisted of (in mM): 150 mM NaCl, 3 mM KCl, 1.2 mM MgCl₂, 1 mM CaCl₂, 20 mM HEPES, 5 mM glucose and 2.5 mM Probenecid (pH 7.4 adjusted with NaOH, osmolarity 320-324 mOsm). In the agonist assay compound, 5-HT (10 µM) was applied by FLIPR after 10 s of the initiation of the measurement.

The MDS Pharma service, which today is part of Eurofins (Taiwan) was used for screening the MAO A and B binding and some other *in vitro* methods which were not available in Orion.

4.3 *Ex vivo* and *In vivo*

Ex vivo and *in vivo* ouabain models were based on literature. Methods are also described in the Paper III and hemodynamic model is described in the Paper I. Below, the most relevant methods used for characterization of the two best compounds are presented for each of the series ORM-10962 and ORM-11372.

4.3.1 *Ouabain model*

Spontaneous aftercontractions in guinea-pig papillary muscles can occur when NCX causes triggered arrhythmias via a delayed afterdepolarization (DAD) mechanism. Ouabain by blocking of sodium-potassium ATPase increases the intracellular sodium which is changed for calcium via NCX [48, 49]. Increased calcium leads to an overload of sarcoplasmic reticulum (SR) and the spontaneous release of calcium from the SR resulting net depolarisation effect i.e. delayed after depolarization (DADs). The aftercontractions (ACs) of measured force signal are equivalent for DADs which are seen as spontaneous twitches after a pacing-controlled twitch. In **Figure 12** how spontaneous aftercontractions (AC1) start to rise 9 min after application of ouabain is shown. Thereafter, spontaneous aftercontractions become more prominent which can be seen by their heights and numbers; and second (AC2) and third aftercontractions (**Figure 12B**), and sometimes even more aftercontractions can occur.

An electrically paced papillary muscle was isolated from the right ventricle of guinea-pig. The papillary muscle was mounted onto a horizontal cuvette and perfused with a Tyrode 1 solution (at 37°C) bubbled with carbogen (95% O₂, 5% CO₂). The composition of the Tyrode 1 solution in mM was: NaCl, 135; MgCl₂×6H₂O, 1; KCl, 5; CaCl₂×2H₂O, 2; NaHCO₃, 15; Na₂HPO₄×2H₂O, 1; and glucose, 10; with a pH of 7.3-7.4. The study compounds were applied in a Tyrode 1 solution and equilibrated 15 min before applying ouabain 5 µM. The final dimethyl sulfoxide (DMSO) concentration was 1.1%.

For *in vivo* ouabain model [48, 49] used in **Paper III**, the male animals (weighing 320-440 g) were anaesthetized with an intraperitoneal injection of pentobarbitone (45 mg/kg). The trachea was cannulated, and positive pressure artificial respiration was maintained throughout the experiment by means of a respirator (under room air, with a volume of 1.5 ml/100g at a rate of 35 strokes/min). The left jugular vein was cannulated with a polythene tube and connected to a slow injection apparatus for ouabain infusion and ORM administration. ORM-10962 (solved in DMSO: isotonic saline, 1:9) was administered in bolus 10 minutes before starting the ouabain infusion. The arterial blood pressure was measured in the right *arteria carotis communis*. The ECG (lead II) was continuously recorded and the heart rate was calculated from ECG signals. A ouabain solution (80 µg/ml) was continuously infused at the rate 6 ml/h/kg. In another series of the suspension of ouabain induced arrhythmias, the ouabain (10 µg/kg/min) was infused for 16 min. ORM-10962 was administered *i.v.* after the termination of ouabain infusion and the ECG and blood pressure changes were recorded.

Figure 12 Ouabain-induced aftercontractions in a guinea-pig papillary muscle.

Figure 13 Analysis of ouabain-induced aftercontractions.

Analyses of the aftercontractions were made in the following manner: the ACs heights were extracted with software (IHME v1.0.9) and plotted as a function of the occurrence time (**Figure 13**). Thereafter, the start of a fast rise of the AC heights (SFR), maximum height (MaxHe) and time to maximum (TtM) height were determined. The extra systole seen in the *in vivo* ouabain model is equivalent to the ACs seen in the *ex vivo* ouabain model.

The aim of the ouabain study was to test whether the test molecules could inhibit ouabain induced aftercontractions in guinea-pig papillary muscle tissue.

4.3.2 Ischemia-reperfusion induced arrhythmias

Myocardial ischemia-reperfusion induced arrhythmias in rats were studied in **Paper III**. Male Sprague-Dawley rats were used. A coronary artery occlusion and reperfusion were performed as described previously by Baczkó *et al.* and Leprán *et al.*[50, 51]. During pentobarbitone anaesthesia (60 mg/kg intraperitoneally) the chest was opened between the fourth and fifth ribs and the heart was exposed. A loose loop of atraumatic silk (4/0 Mersilkblack, W582, Ethicon) was placed around the left main coronary artery. After the surgery, artificial respiration was started (at a rate of 60 strokes/min, volume 1.4 ml/min, Harvard Rodent Ventilator 683, South Natick, MA, USA). After 10 minutes normalization the coronary artery was occluded by tightening the loop for 6 minutes, followed by reperfusion for 5 minutes. The arterial blood pressure was measured via the right *arteria carotis communis*. The left *vena femoralis* was prepared and cannulated for drug administration. A solution of ORM- 10962, as described previously, was administered 10 minutes before the coronary artery occlusion. ECG lead II was registered in both protocols.

The heart rate and the time to the development of ventricular arrhythmias were measured on the ECG.

Myocardial ischemia-reperfusion induced arrhythmias in guinea-pigs were studied in **Paper III**. Male guinea pigs weighing 300-400g were used. The animals were anaesthetized with sodium-pentobarbital (300mg/kg, i.p.) and injected with heparin sodium (300 IU, i.p.). The hearts were rapidly excised, mounted via the aorta on a Langendorff apparatus and retrogradely perfused with a warm (37°C), modified Krebs-Henseleit bicarbonate (KHB) buffer at a constant pressure (80 mmHg). The KHB solution contained (in mmol/L): NaHCO_3 25; KCl 4.3; NaCl 118.5; MgSO_4 1.2; KH_2PO_4 1.2; glucose 10; CaCl_2 1.8, having a pH of 7.4 ± 0.05 when gassed with 95% O_2 + 5% CO_2 . Ventricular fibrillations were induced by reperfusion following 20 minutes zero flow ischemia. The ECG was recorded using the WinWCP software (V4.9.1. Whole Cell Electrophysiology Analysis Program, John Dempster, University of Strathclyde, UK).

In the statistical analysis of **Paper III**, the parameters were expressed as means $\pm$ SEM and after an analysis of variance (one-way ANOVA) groups were compared in pairs by means of the modified “t-statistics” method by Wallenstein *et al.*[52]. The survival rate was compared by using the Chi-square test.

4.3.3 *In vivo hemodynamics of ORM-11372*

In **Paper I** a hemodynamic assessment is described for which healthy and myocardial infarct (MI) male rats were used.

The rats were anaesthetized with isoflurane (2.25-2.5 %, Baxter) in carbogen (95 % O_2 and 5 % CO_2) and nitrous oxide (1:1) using a small rodent ventilator (Ugo Basile 7025, ~10 ml/kg, 60 strokes/min). The rats were infused with 0.9 % NaCl in the carotid vein at the stabilization and baseline level. Systemic arterial blood (SAP) and left ventricular pressures (LVP) were measured with pressure transducers (BP: Isotec, Hugo Sachs Elektronik, Germany; LVP: Mikro-Tip transducer SPR-249, Millar Instruments, USA). The signal was amplified (DC-bridge amplifier type 660, Hugo Sachs Elektronik, Germany), digitized (I/O connector block type SCB-68, National Instruments, USA), recorded and analyzed (IHME 1.0.9, Fision Ltd, Finland). The left ventricular inotrophy ($\text{LV} + dP/dt_{\max}$), relaxation ($\text{LV} - dP/dt_{\min}$) and heart rate (HR) were analyzed from the LVP signal. At the end of the hemodynamic experiments, the rats were killed with an overdose of pentobarbital.

For the healthy rats, ORM-11372 and the dobutamine ascending infusion doses were 16, 83, 167, 333, 500, 667, and 833 $\mu\text{g/kg/min}$ (20 min/dose) and 0.05, 0.1, 0.5, 1, 2, 3, and 4 $\mu\text{g/kg/min}$ (15 min/dose), respectively. For the MI and sham rats, ORM-11372 was administered in ascending infusion doses of 1.7, 17, 167 and 417 $\mu\text{g/kg/min}$. All infusions

were given into the jugular vein at the infusion rate of 10 and 5 ml/kg/h (Terumo TE-311, Belgium) for healthy and MI rats, respectively.

At the end of each infusion, blood samples were taken and centrifuged. Plasma was immediately frozen and stored at -20°C for analyses. ORM-11372 and dobutamine were extracted from the plasma samples using a liquid–liquid extraction and analyzed using LC-MS/MS API 4000. The lower limit of quantification for total plasma concentrations were 32.0 and 5.00 ng/ml from the total for ORM-11372 and dobutamine, respectively. Plasma protein binding for ORM-11372 was assessed using the TRANSIL High Sensitivity Binding Assay according to the manufacturer's instructions (Sovicell, GmbH, Germany).

5 Results and discussion

5.1 *In vivo* and *Ex vivo* results for novel NCX inhibitors

ORM-10962 was the most efficient inhibitor of ouabain-induced aftercontractions of the A-series compounds. ORM-10962 was better than SEA0400 (Figure 14) or ORM-10787 in this model despite that SEA0400 and ORM-10787 achieved better IC_{50} values in the primary screening test. However, SEA0400 had high variation in the measured IC_{50} values. The finding can be partially explained by the poor solubility of SEA0400. Additionally, ORM-10564 is classified as poorly soluble which might reduce the effective concentration in the test solution. However, ORM-10787 and ORM-11037 are classified similarly with moderate solubility as is ORM-10962.

Figure 14 The effects of 10 μ M SEA0400 and ORM-10962 on aftercontractions as a function of time.

Figure 15 the effects on the maximum heights of aftercontractions (*ex vivo*).

As a conclusion ORM-10962 effectively inhibits ouabain-induced aftercontractions and was studied in an ouabain-induced arrhythmias-model *in vivo*. ORM-10787 and ORM-10564 can also effectively inhibit ouabain-induced aftercontractions *in vitro* but the poor solubility of ORM-10564 limits its studies *in vivo*.

In **Paper III** *in vivo* studies performed with ORM-10962 and the obtained results are described. The antiarrhythmic effects of ORM-10962 were studied in an anesthetized ouabain model in guinea pigs, and in ischemia-reperfusion models in rats, as well as Langendorff perfused guinea pig hearts. The selective NCX blockade by ORM-10962 proved to be effective against DAD related arrhythmogenesis as the *ex vivo* studies indicated above and *in vivo* prevented ouabain induced cardiac arrhythmias in guinea pigs. In contrast, ORM-10962 had no apparent effect on ischemia-reperfusion induced arrhythmogenesis generated by coronary artery (LAD) occlusion in rats or in Langendorff perfused guinea pig hearts. The findings are explained by the role of that NCX plays in each of the mechanism that can trigger arrhythmias.

Pre-treatment with ORM-10962 significantly delayed the development of ventricular extra systoles or ventricular tachycardia. The ORM-10962 pre-treatment had no significant effects on the measured hemodynamic parameters, but it delayed the known blood pressure increase caused by the ouabain and also delayed the appearance of arrhythmias. However, the treatment of guinea pigs with ORM-10962 after a suspension of ouabain induced arrhythmias, but this was to a much lesser extend compared to the vehicle group and the duration of the arrhythmic period was significantly shorter. Less than half of the animals developed sporadic ventricular extra systoles, while in the vehicle group all of the animals developed some kind of arrhythmias. The frequent arrhythmias in the vehicle group also prevented the determination of the hemodynamic parameters. After the ORM-10962 treatment, the progression of ouabain induced arrhythmias were less expressed: Only one had VT and no VF occurred.

ORM-10962 showed a lack of effect on ischemia-reperfusion induced arrhythmias in rats. A coronary artery occlusion of 6 minutes rarely produced a life-threatening arrhythmia; however, reperfusion after ischemia rapidly induced severe arrhythmias, frequently leading to the death of the animals. During reperfusion after 6-minute of ischemia, serious arrhythmias, such as irreversible ventricular fibrillation (33%), reversible fibrillation (48%) or tachycardia (62%) were triggered. However, ORM-10962 pre-treatment did not result in any change in the development or the severity of these arrhythmias. There was no significant difference in the measured hemodynamic parameters between the groups before the occlusion or after pre-treatment with ORM-10962.

ORM-10962 had no effect on zero flow ischemia-reperfusion induced arrhythmias in guinea pig isolated Langendorff hearts. Reperfusion induced ventricular fibrillations in

both groups. Pre-treatment with ORM-10962 did not show meaningful difference compared the control response.

The results in intact myocytes of **Paper IV** suggest that ORM-10962 inhibits NCX1.1 asymmetrically, favoring a dominant effect on reverse mode NCX activity. A Ca^{2+} influx via reverse mode NCX, results in a reduced cellular Ca^{2+} content and a net negative inotropic effect (three Na^+ ions per Ca^{2+} ion). The positive inotropy caused by forward mode NCX inhibition is enabled only if other mechanisms ensure that Ca^{2+} cycling and reverse mode NCX is slightly attenuated. During heart failure, increased reverse NCX activity is a characteristic feature and which is a rescue mechanism to maintain the cellular Ca^{2+} load. ORM-10962 blocks the reverse mode NCX effectively. However, it is important to note that the expression levels of proteins are often markedly modified during the diseased stage, i.e. similarly during heart failure, ion channel protein levels might be modified. That may further influence the effect of selective NCX inhibition on the Ca^{2+} handling, and thus, emphasises the importance to performing studies in cardiac cells derived from the patients.

In **Paper I** the inotropic effects of novel NCX inhibitor ORM-11372 are assessed in normal and myocardial infarct rats. In order to assure that ORM-11372 is a potent and selective NCX inhibitor, correlations between human and rat NCX were assessed as well as the selectivity against other ion channels. ORM-11372 inhibited human NCX current similarly in reverse and forward mode, at 5 and 6 nM respectively. The inhibition of the NCX current by ORM-11372 in rat cardiomyocytes was same magnitude for reverse ($\text{IC}_{50} = 11$ nM) and forward (10 nM) NCX currents as it was in humans. ORM-11372 is a selective NCX inhibitor. ORM-11372 did not change the L-type calcium inward current at about a 100-fold concentration (1 μM) of IC_{50} value of their NCX inhibition. The small and non-significant decrease of the average value obtained from these calcium current measurements is most likely due to the slight rundown of the current, which is commonly observed during calcium current measurements. At a 1,000-fold concentration (10 μM) ORM-11372 exerted significant and moderate inhibition of the L-type calcium current.

In vivo, ORM-11372 and dobutamine had 25% positive inotropic effects at 36 and 18 nM free plasma concentrations, respectively. ORM-11372 had no significant effects on systolic blood pressure and heart rate at the same dose level as the maximum inotropic response was reached. When the maximum inotropic response to dobutamine was reached, the heart rate had increased significantly at the same dose level (*Figures 16 and 17*).

Based on the results, it can be concluded that ORM-11372 is a very potent NCX inhibitor with very high selectivity compared to the L-type calcium current. ORM-11372 is a positively inotropic compound in normal and myocardial infarct rats.

Figure 16 Hemodynamics in healthy rats.

Figure 17 Hemodynamics in rats with acute decompensated heart failure.

5.2 Status of NXC inhibitor project in Orion

When the NCX1.1 project was initiated in Orion, the main focus of both cardiovascular research and the related business team was on the chronic indication. However, for business reasons, the main focus was shifted to acute indication around 2002-2003. Hence, ORM-10962 was never taken for safety assessment studies. As series A compounds were not good starting points for short acting treatments, the design focus was redirected to the optimization of series B, that had better potential for a short duration of action. However, cardiovascular research was terminated in 2007 as a business decision. The decision followed the trend with pharma companies globally.

However, ORM-11372 was re-evaluated in 2016 and based on the promising hemodynamic results it was decided to take it forward into a safety assessment study. ORM-11372 was studied in a pilot 3-day study for continuous intravenous infusion toxicity in rats.

The histopathological findings from a 3-day continuous infusion in the external jugular vein showed that the test compound ORM-11372 induced myocardial necrosis, especially in the right ventricle, and vacuolation of stromal cells in the spleen. These changes were likely related to the high local concentration and exaggerated pharmacologic (inotropic) effect and to the metabolism, respectively. In the liver, a treatment-related finding was noted in ORM-11372 (10 mg/kg, micro emulsion vehicle) treated rats. 3 out of 5 rats showed diffuse hepatocellular hypertrophy and 1 out of 5 rats displayed an increase in mitotic hepatocytes. Both findings indicated some degree of CYP induction. Histopathological findings were also seen in the spleen. All the rats treated with the test substance ORM-11372 (10 mg/kg, micro emulsion vehicle) showed minimal (1 out of 5 rats) or marked (4 out of 5 rats) vacuolation in red pulp stromal cells, including macrophages. One of the minor metabolic products of ORM-11372, fluoroaniline, is known to induce vacuolation of red pulp cells [53].

These findings led to the final termination of ORM-11372. Both chemical series remained partly unexplored. However, series A optimization was continued by Sanofi as can be seen from their patent application [12]. Furthermore, it remains to be seen whether anyone sees the potential to continue the optimization of the B series after the Orion patent application is published.

5.3 Legislative perspectives

All animal experiments performed in Szeged were carried out in compliance with the Guide for the Care and Use of Laboratory Animals (USA NIH publication No 85–23, revised 1996) and were approved by the Csongrád County Governmental Office for Food Safety and Animal Health, Hungary (approval No.: XIII/1211/2012). The ARRIVE guidelines were adhered to during the study, (NC3Rs Reporting Guidelines Working Group, 2010).

All animal experiments performed in Orion were made according to European Community Guidelines on the use of experimental animals and approved by Finnish National Animal Experiment Board.

6 Conclusions

New chemistry and design approaches were needed to develop more selective NCX1.1 inhibitors than KB-R 7943 and SEA0400. Through the selected design strategy of reducing the number of rotatable bonds, novel and patentable inhibitors were developed. The strategy also produced a training set molecules for lead hopping by VS. VS provided a starting point for even more potent NCX1.1 inhibitors with distinctively different chemical structures. ORM-10962 was the first reported compound that inhibits both forward and reverse NCX currents at well below sub-micromolar concentrations without affecting further essential sarcolemmal transport mechanisms. ORM-11372 is a new structurally distinctively different NCX1.1 inhibitor which is at least ten-fold more potent than ORM-10962 and can inhibit both forward and reverse NCX currents at low nanomolar concentrations without affecting further essential sarcolemmal transport mechanisms.

Selective NCX inhibition exerts a potent antiarrhythmic effect in digitalis induced arrhythmias. ORM-10962 has an antiarrhythmic effect in vivo models where NCX inhibition is believed to play major role. While for arrhythmias that are triggered mainly by other mechanisms or NCX playing minor role, ORM-10962 failed to show antiarrhythmic properties. The development of arrhythmias in ischemia-reperfusion are rather complex phenomena involving several different mechanisms. such as re-entry, spontaneous automaticity, DADs etc. [54] and NCX may not have important role in each mechanism.

The new structurally distinctive NCX1.1 inhibitor ORM-11372 is derived from the VS hit. It inhibits low nanomolar potency for both forward and reverse NCX currents without affecting further essential sarcolemmal transport mechanisms. ORM-11372 is the most potent NCX inhibitor reported to date and it has very high selectivity compared to the L-type calcium current. *In vivo* ORM-11372 had no significant effects on the systolic blood pressure or heart rate at the same dose level as the maximum inotropic response was reached.

The *in vitro* parameters of ORM-10962 and ORM-11372 were optimized to an acceptable level though some of them remained close to the minimum acceptable level. However, the ORM-11372 aniline as a structural moiety remained and it proved to be a show stopper in the pilot toxicity studies. 3-Fluoroaniline, the minor metabolite of ORM-11372, induced vacuolation in red pulp stromal cells, including macrophages. Therefore, the strategy to overcome the genotoxicity of aniline compounds by choosing a less genotoxic or nongenotoxic aniline based on the literature data as a structural moiety failed. Nevertheless, the Ames test of ORM-11372 was negative. A similar approach has turned out to be a common strategy among medicinal chemists currently. It is recommended to avoid aniline as a structural moiety, but in case this seems to be impossible, the genotoxicity of the aniline moiety and final compound should be studied/addressed early on during the optimization.

Genotoxicity should be also studied in several strains as one strain might give negative results, but another strain could give a clearly positive result.

Sanofi, a pharma company, has continued the optimization of the A series as shown in **Table 1**. They have evidently aimed to improve the solubility by reducing the flatness of the scaffold and they have replaced the pyridyl ring with an aliphatic ring. This structural change should in theory improve the solubility and it might also have positive effects on hERG selectivity. However, these parameters have not been measured in Orion for NCX1.1 inhibitors patented by Sanofi. Thus, they can be considered speculative and we do not have experimental evidence.

Series B optimization remained unfinished due to the business decision to stop cardiovascular research in Orion. There was still room left for chemical optimization and the possibility to improve solubility as well as avoid aniline structural moiety as speculated in Chapter 3.2.2, if the optimization of the compound series would be continued. Only the imagination of the medicinal chemist would limit the possibilities. It will remain to be seen whether anyone from industry or the academic or scientific community work further on the B series chemical optimization. And last but not least, further study the role, function and therapeutic value of NCX1.1 with improved tools.

6.1 Significance of this work

The presented work describes the development of two novel potent and selective NCX1.1 inhibitors with different chemical structures which enable scientific communities to study further functions and therapeutic value of NCX inhibition. The work resulted in three patent applications filed by the Orion Corporation.

6.2 Future research recommendations

Even though the presented work reports the development of two structurally different and novel NCX1.1 inhibitors with different PK profiles as tool compounds for further validate the MoA, the ultimate goal of the research programme was not met in the given timeline. However, the B series still has the potential to further optimize the properties and remove the undesired features of the structure. The role of selective NCX inhibition in triggered automaticity suppression should be further evaluated before its therapeutic role can be confirmed. *In vivo* experiments with ORM-10962 were carried out in healthy, un-diseased myocardium, and in an acute ischemia model in rats and in isolated guinea pig hearts. Similarly, an *in vivo* experiment with ORM-11372 was carried out in healthy animals. However, ORM-11372 also showed efficacy in experimentally induced myocardial infarctions in rats. Arrhythmogenesis frequently occurs in diseased hearts (e.g. in heart failure, or chronic myocardial ischemia) in which calcium handling and NCX function may substantially differ from their state in healthy hearts or following acute coronary ligation

in hearts of healthy animals. Furthermore, differences between species exists in Ca^{2+} -handling mechanisms, hence they may respond differently to NCX inhibition. Therefore, further studies are needed to evaluate the potential therapeutic value of NCX inhibition in cardiac arrhythmogenesis and cardiomyopathies.

References

1. Magee, W.P., et al., *Differing cardioprotective efficacy of the Na⁺/Ca²⁺ exchanger inhibitors SEA0400 and KB-R7943*. Am J Physiol Heart Circ Physiol, 2003. **284**(3): p. H903-10.
2. Molinaro, P., et al., *A new cell-penetrating peptide that blocks the autoinhibitory XIP domain of NCX1 and enhances antiporter activity*. Mol Ther, 2015. **23**(3): p. 465-76.
3. Hamming, K.S., et al., *Inhibition of beta-cell sodium-calcium exchange enhances glucose-dependent elevations in cytoplasmic calcium and insulin secretion*. Diabetes, 2010. **59**(7): p. 1686-93.
4. Matsuda, T., K. Takuma, and A. Baba, *Na⁺(+)-Ca²⁺ exchanger: physiology and pharmacology*. Jpn J Pharmacol, 1997. **74**(1): p. 1-20.
5. Ladilov, Y., et al., *Cardioprotective effects of KB-R7943: a novel inhibitor of the reverse mode of Na⁺/Ca²⁺ exchanger*. Am J Physiol, 1999. **276**(6 Pt 2): p. H1868-76.
6. Hori, M., et al. (1997). Preparation of S-[(benzyloxyphenyl)alkyl]isothiurea derivatives as inhibitors of Na⁺-Ca²⁺ exchanger. JP09067336A, Kanebo Ltd, Japan. p. 15 pp.
7. Ota, T., et al. (1999). Preparation of 2-phenoxyaniline derivatives. WO9920598A1, Taisho Pharmaceutical Co., Ltd., Japan. p. 32 pp.
8. Adorante, J.S., G.R. Ehring, and K.L. Kopper (2002). Targeting the reverse mode of the sodium/calcium exchanger for the treatment of optic neuropathy associated with glaucoma and ischemic optic neuropathy. US6479458B1, Allergan, Inc., USA. p. 6 pp.
9. Adorante, J.S., G.R. Ehring, and K.L. Kopper (2005). Blocking the reverse mode of the sodium-calcium exchanger for the treatment of neurodegenerative conditions. WO2005082345A2, Allergan, Inc., USA. p. 23 pp.
10. Kuramochi, T., et al., *Discovery of an N-(2-aminopyridin-4-ylmethyl)nicotinamide derivative: a potent and orally bioavailable NCX inhibitor*. Bioorg Med Chem, 2005. **13**(12): p. 4022-36.
11. Kakefuda, A., et al. (2003). Preparation of phenoxypyridine derivatives or salts thereof as Na⁺/Ca²⁺ exchanger inhibitors. WO2004000813A1, Yamanouchi Pharmaceutical Co., Ltd., Japan. p. 50 pp.
12. Czechitzky, W., et al. (2014). Preparation of substituted chroman-6-yloxy-cycloalkanes and their use as pharmaceuticals. WO2014135674A1, Sanofi, France. p. 175pp.
13. Ota, T., et al. (1999). Phenoxypyridine derivatives as cytoprotective agents and their uses. JP11092454A, Taisho Pharmaceutical Co., Ltd., Japan. p. 12 pp.
14. Andersson, S., et al., *Making medicinal chemistry more effective--application of Lean Sigma to improve processes, speed and quality*. Drug Discov Today, 2009. **14**(11-12): p. 598-604.
15. Liu, T., et al., *Inhibiting mitochondrial Na⁺/Ca²⁺ exchange prevents sudden death in a Guinea pig model of heart failure*. Circ Res, 2014. **115**(1): p. 44-54.
16. Primessnig, U., et al., *Novel pathomechanisms of cardiomyocyte dysfunction in a model of heart failure with preserved ejection fraction*. Eur J Heart Fail, 2016. **18**(8): p. 987-97.
17. Shpak, B., et al., *Effects of purified endogenous inhibitor of the Na⁺/Ca²⁺ exchanger on ouabain-induced arrhythmias in the atria and ventricle strips of guinea pig*. Eur J Pharmacol, 2006. **553**(1-3): p. 196-204.
18. Geramipour, A., et al., *The investigation of the cellular electrophysiological and antiarrhythmic effects of a novel selective sodium-calcium exchanger inhibitor, GYKB-6635, in canine and guinea-pig hearts*. Can J Physiol Pharmacol, 2016. **94**(10): p. 1090-1101.

19. Milberg, P., et al., *Acute inhibition of the Na⁽⁺⁾/Ca⁽²⁺⁾ exchanger reduces proarrhythmia in an experimental model of chronic heart failure*. Heart Rhythm, 2012. **9**(4): p. 570-8.
20. Hobai, I.A., C. Maack, and B. O'Rourke, *Partial inhibition of sodium/calcium exchange restores cellular calcium handling in canine heart failure*. Circ Res, 2004. **95**(3): p. 292-9.
21. Egar, J., et al., *The Na⁺/Ca²⁺ exchange inhibitor SEA0400 limits intracellular Ca²⁺ accumulation and improves recovery of ventricular function when added to cardioplegia*. J Cardiothorac Surg, 2014. **9**: p. 11.
22. Yoshiyama, M., et al., *Cardioprotective effect of SEA0400, a selective inhibitor of the Na⁽⁺⁾/Ca⁽²⁺⁾ exchanger, on myocardial ischemia-reperfusion injury in rats*. J Pharmacol Sci, 2004. **95**(2): p. 196-202.
23. Namekata, I., et al., *Cardioprotection without cardiosuppression by SEA0400, a novel inhibitor of Na⁺-Ca²⁺ exchanger, during ischemia and reperfusion in guinea-pig myocardium*. Life Sci, 2005. **77**(3): p. 312-24.
24. Yatabe, M.S., et al., *Effects of renal Na⁺/Ca²⁺ exchanger 1 inhibitor (SEA0400) treatment on electrolytes, renal function and hemodynamics in rats*. Clin Exp Nephrol, 2015. **19**(4): p. 585-90.
25. Yang, D., et al., *Na⁺/Ca²⁺ exchange inhibitor, KB-R7943, attenuates contrast-induced acute kidney injury*. J Nephrol, 2013. **26**(5): p. 877-85.
26. Wermuth, C.G., et al., *Glossary of terms used in medicinal chemistry (IUPAC recommendations 1998)*. Pure Appl. Chem., 1998. **70**(5): p. 1129-1143.
27. Kugler-Steigmeier, M.E., et al., *Genotoxicity of aniline derivatives in various short-term tests*. Mutat Res, 1989. **211**(2): p. 279-89.
28. Skipper, P.L., et al., *Monocyclic aromatic amines as potential human carcinogens: old is new again*. Carcinogenesis, 2010. **31**(1): p. 50-58.
29. Menichetti, S., et al., *Synthesis and "double-faced" antioxidant activity of polyhydroxylated 4-thiaflavans*. Org. Biomol. Chem., 2005. **3**(16): p. 3066-3072.
30. Wu, Y.-J., et al., *Fluorine Substitution Can Block CYP3A4 Metabolism-Dependent Inhibition: Identification of (S)-N-[1-(4-Fluoro-3-morpholin-4-ylphenyl)ethyl]-3-(4-fluorophenyl)acrylamide as an Orally Bioavailable KCNQ2 Opener Devoid of CYP3A4 Metabolism-Dependent Inhibition*. J. Med. Chem., 2003. **46**(18): p. 3778-3781.
31. Redfern, W.S., et al., *Relationships between preclinical cardiac electrophysiology, clinical QT interval prolongation and torsade de pointes for a broad range of drugs: evidence for a provisional safety margin in drug development*. Cardiovasc Res, 2003. **58**(1): p. 32-45.
32. Long, Y., et al., *Functional comparison of the reverse mode of Na⁺/Ca²⁺ exchangers NCX1.1 and NCX1.5 expressed in CHO cells*. Acta Pharmacol Sin, 2013. **34**(5): p. 691-8.
33. Lipinski, C.A., et al., *Experimental and computational approaches to estimate solubility and permeability in drug discovery and development settings*. Adv. Drug Delivery Rev., 1997. **23**(1-3): p. 3-25.
34. Zimmer, D., et al., *Bacterial mutagenicity and mammalian cell DNA damage by several substituted anilines*. Mutat Res, 1980. **77**(4): p. 317-26.
35. Assmann, N., et al., *Genotoxic activity of important nitrobenzenes and nitroanilines in the Ames test and their structure-activity relationship*. Mutat Res, 1997. **395**(2-3): p. 139-44.
36. Wess, G., *How to escape the bottleneck of medicinal chemistry*. Drug Discov Today, 2002. **7**(10): p. 533-5.
37. Boyer, E.W. and M. Shannon, *The serotonin syndrome*. N Engl J Med, 2005. **352**(11): p. 1112-20.
38. Wang, Y., et al., *A Facile Synthetic Approach to Prenylated Flavanones: First Total Syntheses of (±)-Bonannione A and (±)-Sophoraflavanone A*. J. Nat. Prod., 2001. **64**(2): p. 196-199.

39. Macquarrie, D.J., R. Nazih, and S. Sebt, *KF/natural phosphate as an efficient catalyst for synthesis of 2'-hydroxychalcones and flavanones*. Green Chem., 2002. **4**(1): p. 56-59.
40. Ye, X.Y., et al., *Synthesis and structure-activity relationships of 2-aryl-4-oxazolylmethoxy benzylglycines and 2-aryl-4-thiazolylmethoxy benzylglycines as novel, potent PPARalpha selective activators- PPARalpha and PPARgamma selectivity modulation*. Bioorg Med Chem Lett, 2010. **20**(9): p. 2933-7.
41. Parry, P.R., M.R. Bryce, and B. Tarbit, *5-Formyl-2-furylboronic acid as a versatile bifunctional reagent for the synthesis of pi-extended heteroaryl-furan systems*. Org Biomol Chem, 2003. **1**(9): p. 1447-9.
42. Huguet, F., et al., *Hydroxamic acids as potent inhibitors of Fe(II) and Mn(II) E. coli methionine aminopeptidase: biological activities and X-ray structures of oxazole hydroxamate-EcMetAP-Mn complexes*. ChemMedChem, 2012. **7**(6): p. 1020-30.
43. Jayyosi, Z., et al. (2000). Preparation of diaryl carboxylic acids and derivatives as peroxisome proliferator-activated receptor ligands. WO2000064888A1, Aventis Pharmaceuticals Products Inc., USA. p. 167 pp.
44. Tata, J.R., et al. (2001). Preparation of γ -hydroxy-2-fluoroalkylaminocarbonyl-1-piperazinepentanamides as HIV protease inhibitors. WO2001038332A1, Merck & Co., Inc., USA. p. 536 pp.
45. Hobai, I.A., D. Khananshvil, and A.J. Levi, *The peptide "FRCRCFa", dialysed intracellularly, inhibits the Na/Ca exchange in rabbit ventricular myocytes with high affinity*. Pflugers Arch, 1997. **433**(4): p. 455-63.
46. Szentadrassy, N., et al., *Apico-basal inhomogeneity in distribution of ion channels in canine and human ventricular myocardium*. Cardiovasc Res, 2005. **65**(4): p. 851-60.
47. Hamill, O.P., et al., *Improved patch-clamp techniques for high-resolution current recording from cells and cell-free membrane patches*. Pflugers Arch, 1981. **391**(2): p. 85-100.
48. Thomas, G.P. and R.M. Tripathi, *Effects of alpha-adrenoceptor agonists and antagonists on ouabain-induced arrhythmias and cardiac arrest in guinea-pig*. Br J Pharmacol, 1986. **89**(2): p. 385-8.
49. Watano, T., et al., *Effect of Na⁺/Ca²⁺ exchange inhibitor, KB-R7943 on ouabain-induced arrhythmias in guinea-pigs*. Br J Pharmacol, 1999. **127**(8): p. 1846-50.
50. Baczko, I., I. Lepran, and J.G. Papp, *Influence of anesthetics on the incidence of reperfusion-induced arrhythmias and sudden death in rats*. J Cardiovasc Pharmacol, 1997. **29**(2): p. 196-201.
51. Lepran, I. and L. Szekeres, *Effect of dietary sunflower seed oil on the severity of reperfusion-induced arrhythmias in anesthetized rats*. J Cardiovasc Pharmacol, 1992. **19**(1): p. 40-4.
52. Wallenstein, S., C.L. Zucker, and J.L. Fleiss, *Some statistical methods useful in circulation research*. Circ Res, 1980. **47**(1): p. 1-9.
53. Suttie, A.W., *Histopathology of the spleen*. Toxicol Pathol, 2006. **34**(5): p. 466-503.
54. Carmeliet, E., *Cardiac ionic currents and acute ischemia: from channels to arrhythmias*. Physiol Rev, 1999. **79**(3): p. 917-1017.

Chronic treatments for arrhythmias and LVH; Acute treatments for acute decompensated heart failure or low cardiac output syndrome.

ISBN 978-952-60-8660-6 (printed)
ISBN 978-952-60-8661-3 (pdf)
ISSN 1799-4934 (printed)
ISSN 1799-4942 (pdf)

Aalto University
School of Chemical Engineering
CHEM, CMAT
www.aalto.fi

**BUSINESS +
ECONOMY**

**ART +
DESIGN +
ARCHITECTURE**

**SCIENCE +
TECHNOLOGY**

CROSSOVER

**DOCTORAL
DISSERTATIONS**