T-matrix modelling of plasmonic nanoparticle arrays

Marek Nečada

T-matrix modelling of plasmonic nanoparticle arrays

Marek Nečada

A doctoral dissertation completed for the degree of Doctor of Science (Technology) to be defended, with the permission of the Aalto University School of Science, at a public examination held online (see <https://necada.org/vaitostilaisuus/>) on 27 November 2020 at 12.00 EET.

**Aalto University
School of Science
Department of Applied Physics
Quantum Dynamics**

Supervising professor

Prof. Päivi Törmä, Aalto University, Finland

Thesis advisors

Prof. Päivi Törmä, Aalto University, Finland

Dr. Jani-Petri Martikainen, Aalto University, Finland

Preliminary examiners

Prof. Tero Setälä, University of Eastern Finland

Prof. Jean-Jacques Greffet, Institut d'Optique, France

Opponent

Dr. Vincenzo Giannini, CSIC, Spain

Aalto University publication series

DOCTORAL DISSERTATIONS 194/2020

© 2020 Marek Nečada

ISBN 978-952-64-0151-5 (printed)

ISBN 978-952-64-0152-2 (pdf)

ISSN 1799-4934 (printed)

ISSN 1799-4942 (pdf)

<http://urn.fi/URN:ISBN:978-952-64-0152-2>

Unigrafia Oy

Helsinki 2020

Finland

Printed matter
4041-0619

Author

Marek Nečada

Name of the doctoral dissertation

T-matrix modelling of plasmonic nanoparticle arrays

Publisher School of Science**Unit** Department of Applied Physics**Series** Aalto University publication series DOCTORAL DISSERTATIONS 194/2020**Field of research** Engineering Physics**Manuscript submitted** 19 August 2020**Date of the defence** 27 November 2020**Permission for public defence granted (date)** 19 October 2020**Language** English☐ **Monograph**☒ **Article dissertation**☐ **Essay dissertation****Abstract**

Active nanoparticle arrays are an attractive platform for manipulating light-matter interactions on the nanoscale. The multiscale character of the optical response of those systems gives rise to exciting physical phenomena, but also makes precise numerical modelling challenging, as it encompasses the interplay of mid-to-long-range periodicity, response of the individual nanoparticles with spatial features much smaller than the wavelength, and possibly nonlinear dynamics of the active medium.

This thesis focuses on the development of the multiple-scattering T-matrix method (MSTMM; also known as the superposition T-matrix method) and its applications in modelling the optical properties of plasmonic nanoparticle arrays. MSTMM is a linear model that combines a computationally efficient (due to a dramatic reduction of the degrees of freedom) yet precise description of the optical interactions between the nanoparticles with a faithful model of linear optical response of the individual nanoparticles.

Chapter 1 reviews some of the commonly used approaches to the numerical simulations, compares their theoretical modelling capabilities and practical scalability with regards to nanoparticle arrays, and provides a motivation for employing the multiple-scattering T-matrix approach.

Chapter 2 is dedicated to the theory of MSTMM and its developments aimed to broaden its applicability to a larger class of physical systems. Its first sections provide a brief guide through the basic concepts of vector spherical wavefunctions, T-matrix and translation operators that make the theoretical backbone of MSTMM in finite systems. The method is then expanded in two main directions: (1) Using exponentially convergent lattice summation, MSTMM is extended to infinite periodic arrays. This enables fast computation of transmission and (employing nonlinear eigensolvers) photonic band structure. (2) The computational efficiency of the method is enhanced by taking into consideration the symmetries of the arrays, which considerably improves the array sizes the method can handle in practice. The group theoretical considerations also find their use in lattice mode analysis.

Chapter 3 showcases MSTMM on several examples, including analysis of some real-world lasing experiments with plasmonic nanoarrays, where the method was used for explaining the observed lasing modes.

Chapter 4 discusses possible utilisation of MSTMM in nonclassical context, mainly in the framework of macroscopic quantum electrodynamics.

Keywords nanophotonics, plasmonics, T-matrix, scattering, nanoparticle arrays**ISBN (printed)** 978-952-64-0151-5**ISBN (pdf)** 978-952-64-0152-2**ISSN (printed)** 1799-4934**ISSN (pdf)** 1799-4942**Location of publisher** Helsinki**Location of printing** Helsinki**Year** 2020**Pages** 189**urn** <http://urn.fi/URN:ISBN:978-952-64-0152-2>

Preface

The work summarised in this thesis has been carried out in the Quantum Dynamics (QD) group at the Department of Applied Physics, Aalto University.

I thank Päivi Törmä for giving me the opportunity to start a doctorate in her group, and Jani-Petri Martikainen for helping me getting around especially in the beginning.

I thank Kristian Arjas, Javier Cuerda, Kostas Daskalakis, Sebastiaan van Dijken, Francisco Freire Fernández, Rui Guo, Tommi Hakala, Mikko Kataja, Nicki Källman, Antti Moilanen, Sara Pourjamal, Heikki Rekola, and Aaro Väkeväinen for the fruitful collaboration.

I thank all the aforementioned and also Jildou Baarsma, Arya Dhar, Elmer Doggen, Rebecca Heilmann, Kukka-Emilia Huhtinen, Aleksi Julku, Jami Kinnunen, Pavel Kliuiev, Pramod Kumar, Long Liang, Heikki Muhli, Sebastiano Peotta, Ville Pykönen, Tuomas Vanhala, Anne-Maria Visuri, Jani Taskinen, Marek Tylutki and many others for all the discussions and excitement we had during those years.

I thank those who had to have most patience with me: Alžběta, Naděžda, Věra and Marja.

My special gratitude goes to Alexandra Elbakyan for her dedication to a principle that should be obvious in a reasonable society—the universal availability of scientific knowledge (generated mostly with public funding) to all humanity, without artificial barriers and without letting the knowledge being parasited on by commercial publishers. Thanks for keeping Sci-Hub up and running. I wish that once we will live in a world where the idea of universal availability of scientific knowledge has been embraced by the relevant decision-makers, together with efficient ways of publishing and sharing that are up to date with the technological progress of humanity.

Helsinki, November 05, 2020,

Marek Nečada

Contents

Preface	5
Contents	7
List of Publications	9
Author's Contribution	11
1. Introduction	17
1.1 Field equations and Green's functions	17
1.2 Approaches to modelling	19
1.2.1 Differential methods	19
1.2.2 Integral methods	20
1.2.3 Computational complexity and scaling	21
2. Multiple-scattering T-matrix method	25
2.1 The finite scattering problem	26
2.1.1 The Silver-Müller radiation conditions and the uniqueness theorem	26
2.2 Spherical vector waves	27
2.3 T-matrix	28
2.3.1 Power transport	30
2.3.2 Plane wave scattering; cross sections	31
2.3.3 Computing the T-matrix	31
2.3.4 T-matrix sanity checks	32
2.4 Multiple scattering	33
2.4.1 Translation operator	35
2.4.2 Green's function for the multiple-scattering system	37
2.5 Infinite periodic systems	38
2.6 Resonances and lattice modes	39
2.7 Symmetries	39
2.8 The QPMS suite	40

Contents

2.9	Further extensions	40
2.9.1	Combining with other integral methods	40
2.9.1.1	Planarly layered media	41
2.9.1.2	Retrieving internal fields	41
2.9.1.3	Multilevel fast multipole algorithm	42
3.	T-matrix method in plasmonic nanoarrays	43
3.1	Lasing in nanoparticle arrays	43
3.1.1	Γ point lasing in Ag nanoparticle square array . . .	43
3.1.2	Γ point lasing in Ni nanoparticle rectangular arrays	45
3.1.3	K point lasing in a honeycomb array	50
3.2	Lattice size and quality factor	53
4.	Combining T-matrix method with quantum physics	55
4.1	MSTMM in macroscopic QED	55
4.2	Interactions between quantum emitters near a nanoparticle .	56
	Conclusion	61
	Bibliography	63
	Publications	69

List of Publications

This thesis consists of an overview and of the following publications which are referred to in the text by their Roman numerals.

- I** M. Nečada and P. Törmä. Multiple-scattering T-matrix simulations for nanophotonics: symmetries and periodic lattices. Submitted to *Communications in Computational Physics*, July 2020.
- II** T. K. Hakala, H. T. Rekola, A. I. Väkeväinen, J.-P. Martikainen, M. Nečada, A. J. Moilanen, and P. Törmä. Lasing in dark and bright modes of a finite-sized plasmonic lattice. *Nature Communications*, **8**, 13687, doi: 10.1038/ncomms13687, January 2017.
- III** S. Pourjamal, T. K. Hakala, M. Nečada, F. Freire-Fernández, M. Kataja, H. Rekola, J.-P. Martikainen, P. Törmä, and S. van Dijken. Lasing in Ni Nanodisk Arrays. *ACS Nano*, **13**, 5686, doi: 10.1021/acsnano.9b01006, April 2019.
- IV** R. Guo, M. Nečada, T. K. Hakala, A. I. Väkeväinen, and P. Törmä. Lasing at K Points of a Honeycomb Plasmonic Lattice. *Physical Review Letters*, **122**, 013901, doi: 10.1103/PhysRevLett.122.013901, January 2019.
- V** M. Nečada, J.-P. Martikainen, and P. Törmä. Quantum emitter dipole-dipole interactions in nanoplasmonic systems. *International Journal of Modern Physics B*, **31**, 17400006, doi: 10.1142/S0217979217400069, August 2017.

The author has also contributed to the following related publication which is not included in this thesis:

- A. I. Väkeväinen, A. J. Moilanen, M. Nečada, T. K. Hakala, K. S. Daskalakis, and P. Törmä. Sub-picosecond thermalization dynamics in condensation of strongly coupled lattice plasmons. *Nature Communications*, **11**, 3139, doi: 10.1038/s41467-020-16906-1, June 2020.

Author's Contribution

Publication I: "Multiple-scattering T-matrix simulations for nanophotonics: symmetries and periodic lattices"

M.N. initiated the T-matrix approach usage in the QD group, conceived and developed the ideas of combining MSTMM with group theory and contour-integral based nonlinear eigenvalue solvers, and derived the formula for 1D-periodic SWF Ewald sum with lattice offset. He is also the author of the implementation of the method, i.e. the QPMS suite, with a few contributions (mainly bug reports) from other users of the code. **M.N.** was the main writer of the manuscript. **P.T.** contributed in writing the manuscript and supervised the project regarding the physics problems the method was desired to solve.

Publication II: "Lasing in dark and bright modes of a finite-sized plasmonic lattice"

P.T. initiated and supervised the project. **T.K.H.**, **H.T.R.** and **A.I.V.** designed and performed the experiments and analysed the data. **H.T.R.** and **J.-P. M.** performed the FDTD simulations. **M.N.**, **J.-P.M.** and **A.J.M.** did the numerical modelling of the finite-sized arrays. In particular, **M.N.** did all the T-matrix simulations. **H.T.R.**, **T.K.H.** and **A.I.V.** fabricated the samples. All authors discussed the results. **T.K.H.** and **P.T.** wrote the manuscript together with all authors.

Publication III: "Lasing in Ni Nanodisk Arrays"

S.P. fabricated the samples and carried out the lasing experiments together with **T.K.H.**, **F.F.F.**, **M.K.** and **H.T.R.** **M.N.** processed the experimental data and performed the T-matrix mode calculations. **J.-P.M.** did the FDTD lasing

simulations. T.K.H., S.v.D., **M.N.** and P.T. wrote the manuscript together with all authors.

Publication IV: “Lasing at K Points of a Honeycomb Plasmonic Lattice”

P.T. initiated and supervised the project. R.G. designed and fabricated the samples, performed the experiments together with T.K.H. and A.I.V., and processed the experimental data. **M.N.** performed the theoretical lattice mode analysis, the T-matrix simulations, and the identification of the lasing mode based on the measured data. R.G., **M.N.**, and P.T. wrote the manuscript together with all authors.

Publication V: “Quantum emitter dipole-dipole interactions in nanoplasmonic systems”

M.N. performed the theoretical and numerical analysis and wrote most of the manuscript. J.-P.M. and P.T. provided guidance and contributed in writing the manuscript.

Symbols, abbreviations, and conventions

Symbols

symbol	common meaning	reference
$\mathbf{1}_A$	indicator function of set A ; $\mathbf{1}_A(x) = 1$ if $x \in A$, else $\mathbf{1}_A(x) = 0$	
$\overleftrightarrow{\mathbf{1}}$	identity tensor in three dimensions	
$\mathbf{I}$	identity matrix	
$a/\tilde{a}$	incident (regular) VSWF field expansion coefficient vector, including / excluding the contributions from other scatterers	sec. 2.3,2.4
$B_\rho(\mathbf{R})$	open ball with radius ρ and center $\mathbf{R}$	
$B_{\rho<,\rho>}(\mathbf{R})$	open spherical shell with inner and outer radii $\rho<,\rho>$ and center $\mathbf{R}$	
c	speed of light in vacuum	
$\delta_{\alpha\beta}$	Kronecker delta; $\delta_{\alpha\beta} = 1$ if $\alpha = \beta$, else $\delta_{\alpha\beta} = 0$	
$\mathbf{E}$	electric field	sec. 1.1
$\mathbf{E}_{\text{inc}}/\mathbf{E}_{\text{sc}}$	incident / scattered electric fields	sec. 2.1
$\varepsilon/\overleftrightarrow{\varepsilon}$	(absolute) electric permittivity / permittivity tensor	
ε_{bg}	(absolute) electric permittivity of the background medium	
f	scattered (outgoing) VSWF field expansion coefficient vector	sec. 2.3
$\overleftrightarrow{G}$	electromagnetic Green's function (dyadic)	sec. 1.1
$h_l^{(1)}$	spherical Hankel function of the first kind	[1, §10.47]
$\mathbf{H}$	magnetic field	sec. 1.1
$\mathbf{H}_{\text{inc}}/\mathbf{H}_{\text{sc}}$	incident / scattered magnetic fields	sec. 2.1
$\Im$	imaginary part	
j_l	spherical Bessel function (first kind) of order l	[1, §10.47]

κ	wave number	sec. 1.1
l	multipole degree index	
m	multipole order index	
$\mu/\overleftrightarrow{\mu}$	(absolute) magnetic permeability / permeability tensor	
μ_{bg}	(absolute) magnetic permeability of the background medium	
ω	angular frequency	sec. 1.1
p	scatterer index	
$\mathcal{P}$	index set of scatterers inside a LOSIB (or inside a unit cell of an infinite LOSIB)	
$\mathbf{r}$	real space variable	
$\hat{\mathbf{r}}$	unit vector corresponding to the angular part of the real space variable; $\hat{\mathbf{r}} = \mathbf{r}/r = \mathbf{r}/ \mathbf{r} $	
$\mathbf{r}_p$	position of scatterer p	
$\mathcal{R}$	regular VSWF translation operator	sec.2.4.1
$\Re$	real part	
$\mathbb{R}$	real numbers	
$\mathcal{S}$	outgoing VSWF translation operator	sec.2.4.1
τ	VSWF type index (1 for magnetic, 2 for electric)	sec. 2.2
$T/T_{\tau lm; \tau' l' m'}$	transition matrix (T-matrix) / T-matrix element	sec. 2.3
$\mathbf{u}_{\tau lm}$	outgoing VSWF	eq. (2.2.3)
$\mathbf{v}_{\tau lm}$	regular VSWF	eq. (2.2.2)
$Y_{l,m}$	spherical harmonic	[1, §14.30]
$\mathbb{Z}$	integers	
$\dagger$	Hermitian adjoint	

Abbreviations

abbreviation	meaning	reference
BEM	boundary element method	
BIC	bound state in the continuum	[2]
CD	coupled dipole method	
DGTD	discrete Galerkin time-domain method	
DO	diffractive order	
DoF	degree(s) of freedom	
FDTD	finite-difference time-domain method	
FETD	finite-element time-domain method	
FMM	fast multipole method	
h.c.	Hermitian adjoint term	
irrep	irreducible representation	
LOSIB	linear optical system with isotropic background	def. 3 (p. 25)
MSTMM	multiple scattering T-matrix method	
QED	quantum electrodynamics	
QPMS	The QPMS Photonic Multiple Scattering suite	sec. 2.8
SCUFF-EM	The SCUFF-EM suite, an implementation of BEM (Surface-CURRENT-Field Formulation of ElectroMagnetism)	[3, 4]
SSS	spherically separate scatterer	def. 4 (p. 25)
SSSh	separating spherical shell	def. 4 (p. 25)
VSWF	vector spherical wavefunction	sec. 2.2

Time-frequency transformation

$$f(\omega) = (2\pi)^{-1/2} \int_{-\infty}^{\infty} f(t) e^{i\omega t} dt \quad (0.0.1a)$$

$$f(t) = (2\pi)^{-1/2} \int_{-\infty}^{\infty} f(\omega) e^{-i\omega t} d\omega \quad (0.0.1b)$$

1. Introduction

The problem of electromagnetic response of a system consisting of many relatively small, compact scatterers in various geometries, and its numerical solution, is relevant to several branches of nanophotonics. In practice, the scatterers often form some ordered structure, such as metallic or dielectric nanoparticle arrays [5–10] that offer many degrees of tunability, with applications including structural color, ultra-thin lenses [11], strong coupling between light and quantum emitters [12–14], weak and strong coupling lasing and Bose-Einstein condensation [15–22], magneto-optical effects [23], or sensing [24].

This work focuses on effective modelling techniques of the optical response in aforementioned systems, which are essential for analysing the experimental observations and for engineering new structures with desired properties, and in particular, on the *multiple-scattering T-matrix method* (MSTMM), which will be the main topic of Chapters 2 and 3. The following sections of this introductory chapter overview the field equations, which are the theoretical starting point to any modelling of photonic structures (Section 1.1), and briefly discuss some frequently used computational electromagnetic methods and their applicability to nanoparticle array (Section 1.2), providing a motivation to MSTMM, theory of which is presented in Chapter 2. Chapter 3 demonstrates applications of MSTMM on some real world experiments. Finally, Chapter 4 considers some possibilities of combining MSTMM with quantum mechanics.

1.1 Field equations and Green’s functions

On the classical level (some quantum aspects will be discussed in Chapter 4), electromagnetic field in matter obeys macroscopic Maxwell’s equations, which in the time domain in their differential form read

$$\nabla \cdot \mathbf{D}(\mathbf{r}, t) = \rho_f(\mathbf{r}, t), \quad (1.1.1a)$$

$$\nabla \cdot \mathbf{B}(\mathbf{r}, t) = 0, \quad (1.1.1b)$$

$$\nabla \times \mathbf{E}(\mathbf{r}, t) = -\frac{\partial \mathbf{B}(\mathbf{r}, t)}{\partial t}, \quad (1.1.1c)$$

$$\nabla \times \mathbf{H}(\mathbf{r}, t) = \mathbf{J}_f(\mathbf{r}, t) + \frac{\partial \mathbf{D}(\mathbf{r}, t)}{\partial t}. \quad (1.1.1d)$$

In the optical context, the system is typically assumed to be electrically neutral, without free charges, i.e. $\rho_f = 0$. In order to describe the fields fully, Maxwell's equations have to be supplemented with the connection between magnetic field strength $\mathbf{H}$ and magnetic flux density $\mathbf{B}$, and between electric field $\mathbf{E}$ and electric displacement field $\mathbf{D}$, i.e. the *constitutive relations* characterising the response of the medium (the inclusion of the free current $\mathbf{J}_f$ depends on the particular description of the system). The constitutive relations can take various forms; in practice, certain assumptions have to be imposed in order to make the problem possible to solve with a given method. For instance, one might assume the following form:

$$\mathbf{D}(\mathbf{r}, t) = (2\pi)^{-1/2} \int d\tau \overleftrightarrow{\epsilon}(\mathbf{r}, \tau) \cdot \mathbf{E}(\mathbf{r}, t - \tau), \quad (1.1.2a)$$

$$\mathbf{B}(\mathbf{r}, t) = (2\pi)^{-1/2} \int d\tau \overleftrightarrow{\mu}(\mathbf{r}, \tau) \cdot \mathbf{H}(\mathbf{r}, t - \tau). \quad (1.1.2b)$$

This particular form includes three important assumptions about the system: *space locality*, *linearity*, and *time translation symmetry*. Thanks to the space locality, which tends to be a reasonable assumption unless one gets too close to the atomic length scale, the whole set of equations remains differential (instead of e.g. integro-differential) w.r.t. the variable $\mathbf{r}$. Linearity is quite a strong assumption, but it enables utilising linear transformations—Fourier transform in particular. Time translation symmetry is a natural assumption in a vast majority of cases, and it gives a simple multiplicative form to the constitutive relations in frequency domain. Indeed, applying Fourier transform (0.0.1a) w.r.t. time variable t , Maxwell's equations (1.1.1) transform to their frequency domain versions

$$\nabla \cdot \mathbf{D}(\mathbf{r}, \omega) = \rho_f(\mathbf{r}, \omega), \quad (1.1.3a)$$

$$\nabla \cdot \mathbf{B}(\mathbf{r}, \omega) = 0, \quad (1.1.3b)$$

$$\nabla \times \mathbf{E}(\mathbf{r}, \omega) = i\omega \mathbf{B}(\mathbf{r}, \omega), \quad (1.1.3c)$$

$$\nabla \times \mathbf{H}(\mathbf{r}, \omega) = \mathbf{J}_f(\mathbf{r}, \omega) - i\omega \mathbf{D}(\mathbf{r}, \omega), \quad (1.1.3d)$$

and the constitutive relations (1.1.2) take the form

$$\mathbf{D}(\mathbf{r}, \omega) = \overleftrightarrow{\epsilon}(\mathbf{r}, \omega) \cdot \mathbf{E}(\mathbf{r}, \omega), \quad (1.1.4a)$$

$$\mathbf{B}(\mathbf{r}, \omega) = \overleftrightarrow{\mu}(\mathbf{r}, \omega) \cdot \mathbf{H}(\mathbf{r}, \omega), \quad (1.1.4b)$$

so thanks to the assumptions on the constitutive relations, the whole system reduces to a set of differential equations with the space variable $\mathbf{r}$, which can be solved for each angular frequency ω separately.

If the medium is homogeneous and isotropic, so that $\overleftrightarrow{\varepsilon}(\mathbf{r}, \omega) = \varepsilon_{\text{bg}}(\omega) \overleftrightarrow{\mathbf{1}}$, $\overleftrightarrow{\mu}(\mathbf{r}, \omega) = \mu_{\text{bg}}(\omega) \overleftrightarrow{\mathbf{1}}$, and does not contain free charges ($\rho_{\text{f}} = 0$), the whole system (1.1.3)–(1.1.4) reduces to (inhomogeneous) vector Helmholtz equation

$$\nabla^2 \mathbf{E}(\mathbf{r}, \omega) + (\kappa(\omega))^2 \mathbf{E}(\mathbf{r}, \omega) = +i\omega\mu(\omega) \mathbf{J}_{\text{f}}(\mathbf{r}, \omega) \quad (1.1.5)$$

with transversality condition $\nabla \times \mathbf{E}(\mathbf{r}, \omega) = 0$; here $(\kappa(\omega))^2 = \omega^2 \mu(\omega) \varepsilon(\omega)$ is the (squared) wave number. Note that the sign on the right hand side of (1.1.5) is given by the convention used for Fourier transform (0.0.1). The electric field caused by the currents can be expressed using the corresponding retarded Green's functions

$$\mathbf{E}(\mathbf{r}, \omega) = \int d\mathbf{r}' \overleftrightarrow{G}(\kappa(\omega); \mathbf{r}, \mathbf{r}') \cdot i\omega\mu(\omega) \mathbf{J}_{\text{f}}(\mathbf{r}', \omega). \quad (1.1.6)$$

The retarded Green's function for electric field in isotropic medium has the well-known explicit form

$$\begin{aligned} \overleftrightarrow{G}(\kappa; \mathbf{r}, \mathbf{r}') &= \overleftrightarrow{G}(\kappa; \mathbf{r} - \mathbf{r}') = \left(\overleftrightarrow{\mathbf{1}} + \kappa^{-2} \nabla_{\mathbf{r}} \nabla_{\mathbf{r}} \right) g(\kappa; |\mathbf{r} - \mathbf{r}'|), \\ g(\kappa, r) &= \frac{e^{i\kappa r}}{4\pi r}. \end{aligned}$$

In principle it is possible to write the fields as in (1.1.6) also for inhomogeneous or anisotropic medium, but in such cases the corresponding Green's functions usually lack a closed form [25, chapter 2].

1.2 Approaches to modelling

At a high level, the computational methods of solving Maxwell's equations can be divided into two main classes [26]:

1. The *differential* methods, where Maxwell's equations (1.1.1) are solved directly in their differential form for the fields that permeate all of the solution region.
2. The *integral* methods solve for the sources (polarisations / current densities) after determining the interactions between them.

1.2.1 Differential methods

Differential methods are attractive thanks to their simplicity and generality: The fields are evaluated in a discretised volume, where the interactions between discretisation units are short-ranged, involving only several adjacent cells, avoiding the need to work with large dense matrices. Spatially varying constitutive

relations are typically as easy to deal with as homogeneous media; furthermore, time-domain differential methods, such as FDTD [27–29], FETD [30] or DGTD [31, 32] do not even require linearity of the constitutive relations, which allows for modelling nonlinear phenomena such as lasing [33–35].

However, the scaling properties present an important drawback of the differential methods. The discretisation mesh (or grid) introduces artificial dispersion in the system due to the cumulative phase errors that build up in the field propagation through the mesh. These errors can be mitigated by making the mesh (grid) finer to keep the results accurate. In effect, the number of unknowns (degrees of freedom) N grows faster than linearly with increasing volume of the physical system. The actual growth rate depends on the choice of the differential equation solving scheme; for a second-order accurate scheme, the number of unknowns scales as $(\kappa d)^{3/2}$ per physical dimension, where d is the length scale of the simulation region [36]. There are other problems that might arise, depending on the particular differential method chosen. For example, the rectangular grid used in FDTD would internally break the hexagonal symmetry of a periodic system with $p6m$ symmetry group, potentially leading to a numerical artifact in the form of splitting the doubly degenerate modes at Γ and K points. Using FETD or DGTD with symmetry-conforming mesh instead would avoid this issue.

1.2.2 Integral methods

Integral methods typically require linear constitutive relations (1.1.4) and make use of homogeneous regions, inside which the fields can propagate “freely”, the propagation being described by closed-form Green’s functions. Possible inhomogeneities (scatterers) in the path produce induced currents, which in turn propagate (via the same Green’s functions) to all the other scatterers in the system, inducing new currents in those in a linear manner, etc. Putting all these together, one gets a linear integral equation that describes the electromagnetic response of the whole system. Let us write down a simple volume integral formulation of the problem. The electric field at point $\mathbf{r}$ is given by the fields created by all the induced current densities throughout the simulation region, propagated by the Green’s function, and a given driving field $\mathbf{E}_{\text{inc}}$ originating from external sources,

$$\mathbf{E}(\mathbf{r}) = \mathbf{E}_{\text{inc}}(\mathbf{r}) + \int_{\substack{\text{inhom.} \\ \text{regions}}} d\mathbf{r}' \overleftrightarrow{G}(\mathbf{r}, \mathbf{r}') \cdot \mathbf{J}_{\text{ind}}(\mathbf{r}') \quad (1.2.1)$$

(we don’t explicitly write the frequency dependence here anymore, but it is present implicitly). The induced current, on the other hand, is the “free” current density that is left in by imposing $\mathbf{D}(\mathbf{r}) = \overleftrightarrow{\epsilon}_{\text{bg}} \cdot \mathbf{E}(\mathbf{r})$ in the Maxwell–Ampère equation (1.1.3d),

$$\mathbf{J}_{\text{ind}}(\mathbf{r}) = i\omega (\overleftrightarrow{\epsilon}_{\text{bg}} - \overleftrightarrow{\epsilon}(\mathbf{r})) \cdot \mathbf{E}(\mathbf{r}). \quad (1.2.2)$$

Combining (1.2.1) and (1.2.2) eliminates the scattered field,

$$\begin{aligned} \mathbf{J}_{\text{ind}}(\mathbf{r}) - i\omega (\overleftrightarrow{\epsilon}_{\text{bg}} - \overleftrightarrow{\epsilon}(\mathbf{r})) \int_{\text{inhom. regions}} d\mathbf{r}' \overleftrightarrow{G}(\mathbf{r}, \mathbf{r}') \cdot \mathbf{J}_{\text{ind}}(\mathbf{r}') = \\ = i\omega (\overleftrightarrow{\epsilon}_{\text{bg}} - \overleftrightarrow{\epsilon}(\mathbf{r})) \cdot \mathbf{E}_{\text{inc}}(\mathbf{r}). \end{aligned} \quad (1.2.3)$$

If the obstacles consist of locally homogeneous regions, the problem can be instead reformulated in terms of surface current densities on the region boundaries, leading to reduction of the dimensionality of the unknowns—as in the boundary element method (BEM). In either case, once the induced currents are solved, the total fields can be easily obtained by applying the Green’s functions as in (1.2.1).

After discretisation, the problem gains a matrix form

$$MJ = J_0 \quad (1.2.4)$$

where J is the vector of N unknowns (the currents induced in the discretised volume/surface elements), M is the interaction matrix between the currents, and J_0 is the vector of currents induced directly by the external driving, before scattering further. Unlike in differential methods where the interactions between unknowns are sparse with $O(N)$ non-zero elements, M is a $N \times N$ shaped *dense* matrix which typically has to be kept in memory. On the other hand, if most of the simulation region is filled with homogeneous medium where the field propagates freely and no induced currents are present, the number of unknowns is considerably lower.

The reduction of degrees of freedom (DoF) in integral methods is significant in the case of two-dimensional nanoparticle arrays: the system can be thought as infinite space filled with homogeneous background medium, filled only with a layer of nanoparticles, and possibly a layer of homogeneous substrate. If an index-matching medium is used on both sides of the nanoparticle array, as is often done in the experiments, the substrate interface can be neglected as well.

1.2.3 Computational complexity and scaling

Let us think about how well would these methods scale if—instead of one spherical particle—we had a finite square array of $n \times n$ equal spheres with diameter D , equally spaced with period p . Let $d_{\text{BEM}}, d_{\text{FDTD}}$ denote the approximate BEM mesh and FDTD grid spacings. With BEM, the number of degrees of freedom would scale as $N_{\text{BEM}} \propto O(n^2 D^2 / d_{\text{BEM}}^2)$; if the system is to be solved using LU decomposition, the full interaction matrix would take memory $M_{\text{BEM/LU}} \propto N_{\text{BEM}}^2 \propto O(n^4 D^4 / d_{\text{BEM}}^4)$, and the time complexity of LU decomposition in practice goes like $t_{\text{BEM/LU}} \propto N_{\text{BEM}}^3 \propto O(n^6 D^6 / d_{\text{BEM}}^6)$. For FDTD, one has to consider also the volume between the particles, so the degrees of freedom behave as $N_{\text{FDTD}} \propto O(n^2 D p^2 / d_{\text{FDTD}}^3)$; the memory requirements are directly proportional to the number of degrees of freedom,

and the number T_{FDTD} of time steps must be adjusted to unitless length scale of the system $T_{\text{FDTD}} \propto np/d_{\text{FDTD}}$, so that the driving impulse manages to propagate through the whole system during that time, so the computational time scales as $t_{\text{FDTD}} \propto O(N_{\text{FDTD}} T_{\text{pass}}) = O(n^3 D p^3 / d_{\text{FDTD}}^4)$. At a first glance, this looks (asymptotically) like a clear win for FDTD. However, as mentioned above, the grid spacing has to be adjusted in order to compensate for the cumulative error, $d_{\text{FDTD}} \propto (np)^{-\alpha}$, so for the typical value $\alpha = 0.5$ we already have $t_{\text{FDTD}} \propto O(n^5 D p^5)$. In BEM and other integral methods, the fields are propagated from scatterer to scatterer via exact closed-form Green’s functions, so although the mesh density affects model accuracy at a single-particle level, it typically need not be adjusted when adding other scatterers, so d_{BEM} can be considered constant. Also, the estimated number of FDTD simulation time steps T_{pass} was based on the assumption that resonances are absent and the initial pulse can travel through the whole system “freely” without spending too much time scattering back and forth inside. However, the lattice resonances are where most of the interesting physics happens, so if using FDTD, one has to take into account as well and adjust the number of time steps also by the resonance quality factor Q , so $T_{\text{FDTD}} \propto Qnp/d_{\text{FDTD}}$. Empirical data indicate that the quality factor can be expected to grow linearly with n or even faster (cf. Section 3.2); so taking an optimistic estimate $Q \propto n$ we arrive at $t_{\text{FDTD}} \propto O(n^6 D p^5)$, and FDTD loses the apparent edge in time complexity from several lines earlier. Considering also the constant factors involved, this makes a clear ground for choosing an *integral, frequency domain* method for modelling nanoparticle arrays.

Remark 1. There are also other approaches to solving (1.2.4) than LU factorisation of M . For example, one could solve for J iteratively, without the need to store the interaction matrix M and possibly reducing the computation time. However, iterative methods may have serious convergence problems, especially around resonances [37]. Iterative methods are not considered further in this work.

LU factorisation has also the obvious advantage that once M is factorised, one can easily get solutions for many different right-hand sides (driving fields) of (1.2.4), each one costing only $O(N^2)$ time where N is the number of unknowns.

Remark 2. So the differential methods might not be ideal for large finite systems of scatterers, but what about infinite periodic lattices? In periodic systems, the fields can be solved in terms of quasi-periodic Bloch waves, for which the degrees of freedom can be limited to a single unit cell. In differential methods, this is easily implemented by setting quasi-periodic boundary conditions at the edges of a unit cell—nothing fundamentally changes in the sparse structure of the interactions between unknowns. For integral methods, where the sources (induced currents) are interconnected densely with each other, meaning that in infinite periodic system, each source interacts with an infinite number of sources from all the other unit cells. Due to the imposed Bloch condition, these sources are just quasiperiodic images of the sources in the representative unit cell,

but one has to deal with badly convergent infinite lattice sums of the Green's functions. This makes the integral methods more difficult in infinite periodic systems, but there is a way around in the form of exponentially convergent lattice summation techniques, which are presented in Section 2.5.

On a machine with 8 GB of RAM available for storing the interaction matrix, one could simulate a system of around 80 particles using BEM with LU solver, considering a mesh of ~ 100 surface elements for each particle. This is still far below the number of tens of thousands of particles appearing in the experiments. A possible way around from this lies in reducing the number of unknowns N . An extreme example of this is the coupled dipole (CD) method, in which each scatterer, centered at $\mathbf{r}_i$, is reduced to a single dipolar element characterised by a mere 3×3 polarisability tensor $\overleftrightarrow{\alpha}_i$ coupled with other particles by electric Green's dyadics, so the system (1.2.4) takes the form

$$\mathbf{p}_i - \overleftrightarrow{\alpha}_i \cdot \sum_{j \neq i} \overleftrightarrow{G}(\mathbf{r}_i, \mathbf{r}_j) \cdot \mathbf{p}_j = \overleftrightarrow{\alpha}_i \cdot \mathbf{E}_{\text{inc}}(\mathbf{r}_i). \quad (1.2.5)$$

For ν particles in the system, there would be in total 3ν complex unknowns (dipole polarisation elements) and if each of the interaction matrix element is represented by the IEEE 754 double-precision floating-point format (16 B per one complex number), its size would be $\nu^2 \cdot 144$ B, so with 8 GB RAM available, one could fit in an interaction matrix for 7723 particles, which is an order of magnitude close to what is typically used in the real world experiments.

The CD method is very easy to implement and has indeed been used for modelling optical response of nanoparticle arrays [38]. However, the dipole approximation is obviously too rough for particles with a diameter larger than a small fraction of wavelength, leading to quantitative errors if the particle size is increased. One could argue that if the particles are really tiny, CD would still give fairly accurate results for a routine radar cross section calculation. Nevertheless, CD fails on qualitative level—even for tiny particles—if one wants to study lattice modes of the nanoparticle arrays: there are modes in which the particles' electric dipole moments completely vanish due to symmetry, and regardless of how small the particles are, the excitations have quadrupolar or higher-degree multipolar character. These modes, belonging to what is sometimes called *optical bound states in the continuum (BIC)* [2], tend to have extraordinarily high quality factors and are of particular interest due to the physical phenomena such as lasing that have been observed therein, but the CD method completely fails to capture them. Therefore, higher multipoles need to be taken into consideration. One possible way is to make a step back and represent each scatterer by more surface or volume elements, but this would again cause a steep increase in the number of unknowns and size of the interaction matrix.

A much more effective approach is to represent each scatterer with its *transition matrix*, a.k.a. *T-matrix*, which describes the response to electromagnetic radiation in terms of a multipolar basis especially suitable for scattering calcula-

tions. Instead of Green's functions, the scatterers are coupled to each other by derived objects, the translation operators, whose elements also have closed-form expressions (albeit much more complicated than the original Green's functions). This is the main idea behind the *multiple-scattering T -matrix method*, also called *superposition T -matrix method* in some literature, which is—together with its application in nanophotonics—the main topic of this work, presented in the following chapters. It will become apparent that the T -matrix approach not only has superior computational performance, but also provides a relatively intuitive framework much better suitable for abstract analysis of lattice modes than the other aforementioned methods.

2. Multiple-scattering T-matrix method

The T-matrix method builds on the idea of expanding the electromagnetic field around a scatterer into a basis of vector spherical wavefunctions (VSWFs) in which a single scattering problem is solved. The VSWF basis has the advantage of being suitably localised in space (as much as the Helmholtz equation allows), which allows for the multipolar truncation that makes the T-matrix method so efficient. The multiple-scattering T-matrix method relies on VSWF re-expansion between different origins (according to the locations of scatterers): the driving electromagnetic field incident onto a scatterer is expanded into a vector spherical wavefunction (VSWF) basis in which the single scattering problem is solved, and the scattered field is then re-expanded into VSWFs centered at the other scatterers. Repeating the same procedure with all (pairs of) scatterers yields a set of linear equations, solution of which gives the coefficients of the scattered field in the VSWF bases. Once these coefficients have been found, one can evaluate various quantities related to the scattering (such as cross sections or the scattered fields) quite easily.

In order to facilitate the formal description of the method, let us first introduce some terminology.

Definition 3. Let there be a system satisfying the macroscopic Maxwell's equations (1.1.3) together with the local linear constitutive relations (1.1.4), such that the most of its spatial domain is filled with homogeneous and isotropic medium. We denote such a system as a *linear optical system with isotropic background* (LOSIB). Unless specified otherwise, we will assume that the free source densities $\rho_f(\mathbf{r}, \omega)$, $\mathbf{J}_f(\mathbf{r}, \omega)$ are zero.

Definition 4 (spatial division of LOSIB). The *foreground* F of a LOSIB with background medium specified by $\varepsilon_{bg}(\omega)$, $\mu_{bg}(\omega)$ consists of the closure of all parts of space where the constitutive relations deviate from the background ones, i.e.

$$F = \overline{\left\{ \mathbf{r} \in \mathbb{R}^3 : \overleftrightarrow{\varepsilon}(\mathbf{r}, \omega) \neq \varepsilon_{bg}(\omega) \overleftrightarrow{1} \vee \overleftrightarrow{\mu}(\mathbf{r}, \omega) \neq \mu_{bg}(\omega) \overleftrightarrow{1} \right\}}.$$

The *background* B is the complement of the foreground, i.e. the largest open set where the background constitutive relations do hold,

$$B = \mathbb{R}^3 \setminus F.$$

A *scatterer* S is a part of the foreground that does not touch the rest of the foreground, that is

$$S \subset F, S \cap \overline{F \setminus S} = \emptyset.$$

A scatterer is *compact* if it has a finite diameter.

An open spherical shell $B_{r_<, r_>}(\mathbf{R})$ centered at $\mathbf{R}$, with inner radius $r_<$ and outer radius $r_>$ is a *separating spherical shell* (SSSh) of a compact scatterer S if S is completely contained in the inner ball $\overline{B_{r_<}(\mathbf{R})}$ and the outer ball does not intersect with other parts of the foreground, i.e. if $S \subset \overline{B_{r_<}(\mathbf{R})}$ and $B_{r_>}(\mathbf{R}) \cap (F \setminus S) = \emptyset$.

A *spherically separate scatterer* (SSS) is a compact scatterer that has a nonempty separating spherical shell.

Remark 5. By this definition, a scatterer does not have to be a connected set, and the union of two scatterers is again a scatterer. As a consequence, there can be many ways to divide the foreground into mutually disjoint scatterers. This can be conveniently used if such a division is expected to yield SSSs: it might happen that two disjoint scatterers S_1, S_2 have overlapping circumscribed balls (thus at least either of them is not a SSS) but $S_1 \cup S_2$ is a SSS. If necessary, the whole foreground can be treated as a single scatterer.

2.1 The finite scattering problem

The basic linear scattering problem with a compact scatterer can be formulated as follows. Let there be a given LOSIB and given driving fields $\mathbf{E}_{\text{inc}}, \mathbf{H}_{\text{inc}}$ that satisfy Maxwell's equations in an auxilliary system that is completely filled with the isotropic background medium with material constants $\varepsilon_{\text{bg}}(\omega), \mu_{\text{bg}}(\omega)$; in the auxilliary system, we allow for nonzero free source currents. We search for solutions in the form

$$\mathbf{E} = \mathbf{E}_{\text{inc}} + \mathbf{E}_{\text{sc}}, \quad (2.1.1)$$

$$\mathbf{H} = \mathbf{H}_{\text{inc}} + \mathbf{H}_{\text{sc}}, \quad (2.1.2)$$

fulfilling Maxwell's equations everywhere in the given LOSIB. From the assumed linearity it is obvious that $\mathbf{E}_{\text{sc}}, \mathbf{H}_{\text{sc}}$ themselves must then also satisfy Maxwell's equations in the background. Moreover, in order to give them the interpretation of scattered fields, $\mathbf{E}_{\text{sc}}$ and $\mathbf{H}_{\text{sc}}$ are expected to satisfy the Silver-Müller radiation conditions (2.1.3) below.

2.1.1 The Silver-Müller radiation conditions and the uniqueness theorem

Let D be an open bounded domain in $\mathbb{R}^3$ and let $\mathbf{E}_{\text{sc}}, \mathbf{H}_{\text{sc}} \in C^1(\mathbb{R}^3 \setminus \overline{D})$ be a solution of Maxwell's equations in a lossless, isotropic medium. The

Silver-Müller radiation conditions are [39, Sect. 1.4.1] [40, Sect. 4.2]

$$\lim_{|\mathbf{r}| \rightarrow \infty} \sqrt{\mu} \hat{\mathbf{r}} \times \mathbf{H}_{\text{sc}} + \sqrt{\varepsilon} \mathbf{E}_{\text{sc}} = 0, \quad (2.1.3a)$$

$$\lim_{|\mathbf{r}| \rightarrow \infty} \sqrt{\varepsilon} \hat{\mathbf{r}} \times \mathbf{E}_{\text{sc}} - \sqrt{\mu} \mathbf{H}_{\text{sc}} = 0 \quad (2.1.3b)$$

where the convergence is uniform in all directions $\hat{\mathbf{r}}$. It can be shown that Eqs. (2.1.3a) and (2.1.3b) are equivalent to each other [40, Corollary 4.14], and also to the Sommerfeld radiation conditions (for the scalar Helmholtz equation $\nabla^2 \psi + \kappa^2 \psi = 0$) imposed on all the fields' Cartesian components, that is [40, Corollary 4.14],

$$\hat{\mathbf{r}} \cdot \nabla \psi(\mathbf{r}) - i\kappa \psi(\mathbf{r}) = o(|\mathbf{r}|^{-1}), \quad |\mathbf{r}| \rightarrow \infty$$

uniformly for all directions $\hat{\mathbf{r}}$.

The Silver-Müller radiation conditions provide a suitable definition of *scattered* fields and ensure their uniqueness [40, Theorem 4.18]. The retarded Green's function for homogeneous isotropic medium from Eq. (1.1.7) satisfies the radiation conditions.

2.2 Spherical vector waves

The homogeneous vector Helmholtz equation

$$(\nabla^2 + \kappa^2) \mathbf{\Psi} = 0 \quad (2.2.1)$$

with the transversality condition $\nabla \cdot \mathbf{\Psi} = 0$ can be solved by separation of variables in spherical coordinates to give the solutions – the *regular* and *outgoing* vector spherical wavefunctions (VSWFs) $\mathbf{v}_{\tau lm}(\kappa \mathbf{r})$ and $\mathbf{u}_{\tau lm}(\kappa \mathbf{r})$, respectively, defined as follows:

$$\begin{aligned} \mathbf{v}_{1lm}(\kappa \mathbf{r}) &= j_l(\kappa r) \mathbf{A}_{1,l,m}(\hat{\mathbf{r}}), \\ \mathbf{v}_{2lm}(\kappa \mathbf{r}) &= \frac{1}{\kappa r} \frac{d(\kappa r j_l(\kappa r))}{d(\kappa r)} \mathbf{A}_{2,l,m}(\hat{\mathbf{r}}) + \sqrt{l(l+1)} \frac{j_l(\kappa r)}{\kappa r} \mathbf{A}_{3,l,m}(\hat{\mathbf{r}}), \end{aligned} \quad (2.2.2)$$

$$\begin{aligned} \mathbf{u}_{1lm}(\kappa \mathbf{r}) &= h_l^{(1)}(\kappa r) \mathbf{A}_{1,l,m}(\hat{\mathbf{r}}), \\ \mathbf{u}_{2lm}(\kappa \mathbf{r}) &= \frac{1}{\kappa r} \frac{d(\kappa r h_l^{(1)}(\kappa r))}{d(\kappa r)} \mathbf{A}_{2,l,m}(\hat{\mathbf{r}}) + \sqrt{l(l+1)} \frac{h_l^{(1)}(\kappa r)}{\kappa r} \mathbf{A}_{3,l,m}(\hat{\mathbf{r}}), \end{aligned} \quad (2.2.3)$$

$$\tau = 1, 2; \quad l = 1, 2, 3, \dots; \quad m = -l, -l+1, \dots, +l,$$

where $\mathbf{r} = r\hat{\mathbf{r}} = r(\sin \theta (\hat{\mathbf{x}} \cos \phi + \hat{\mathbf{y}} \sin \phi) + \hat{\mathbf{z}} \cos \theta)$; $j_l(x)$, $h_l^{(1)}(x) = j_l(x) + iy_l(x)$ are the regular spherical Bessel function and spherical Hankel function

of the first kind, respectively, and $\mathbf{A}_{\tau,l,m}$ are the *vector spherical harmonics*, defined by

$$\begin{aligned}\mathbf{A}_{1,l,m}(\hat{\mathbf{r}}) &= \frac{1}{\sqrt{l(l+1)}} \nabla \times (\mathbf{r} Y_{l,m}(\hat{\mathbf{r}})) = \frac{1}{\sqrt{l(l+1)}} \nabla Y_{l,m}(\hat{\mathbf{r}}) \times \mathbf{r}, \\ \mathbf{A}_{2,l,m}(\hat{\mathbf{r}}) &= \frac{1}{\sqrt{l(l+1)}} r \nabla Y_{l,m}(\hat{\mathbf{r}}), \\ \mathbf{A}_{3,l,m}(\hat{\mathbf{r}}) &= \hat{\mathbf{r}} Y_{l,m}(\hat{\mathbf{r}}).\end{aligned}\tag{2.2.4}$$

Vector spherical harmonics are a straightforward extension of the (scalar) spherical harmonics $Y_{l,m}$. Importantly, they constitute a complete orthogonal basis for square integrable vector fields on a sphere [40, p. 170].

Note that the regular waves $\mathbf{v}_{\tau lm}$ (with fields expressed in Cartesian coordinates) have all well-defined limits in the origin, and except for the “electric dipolar” waves $\mathbf{v}_{21m}$, they vanish. While the regular VSWFs therefore solve the homogeneous Helmholtz equation everywhere, the outgoing VSWFs $\mathbf{u}_{\tau lm}$ satisfy them everywhere except the origin where they have a singularity; importantly, the outgoing VSWFs satisfy the Silver-Müller radiation conditions (hence the name). Figure 2.2.1 illustrates the VSWFs of the lowest multipole degrees.

VSWFs are the vector analogues of *scalar* spherical wavefunctions

$$\begin{aligned}v_{l,m}(\kappa \mathbf{r}) &= j_l(\kappa r) Y_{l,m}(\hat{\mathbf{r}}), \\ u_{l,m}(\kappa \mathbf{r}) &= h_l^{(1)}(\kappa r) Y_{l,m}(\hat{\mathbf{r}}),\end{aligned}$$

which solve the scalar Helmholtz equation. The scalar spherical wavefunctions will be useful for expressing the translation operators. The aforementioned transversal VSWFs can be complemented by a set of longitudinal solutions of the vector Helmholtz equation $\mathbf{v}_{3lm}, \mathbf{u}_{3lm}$, but these are not relevant for scattering of electromagnetic waves.

2.3 T-matrix

With the VSWFs properly introduced, we can return to the single particle scattering problem, to introduce the T-matrix. Let there be a LOSIB with a single compact scatterer. We choose the coordinate origin to lie in the center of the scatterer’s circumscribed ball $\overline{B_{R<}(\mathbf{0})}$ we take $(\kappa(\omega))^2 = \omega^2 \mu_{\text{bg}}(\omega) \varepsilon_{\text{bg}}(\omega)$, and assume a time-harmonic driving field $\mathbf{E}_{\text{inc}}(\mathbf{r}), \mathbf{H}_{\text{inc}}(\mathbf{r})$ such that there are no source current densities $\mathbf{J}_f$ inside a larger ball $\overline{B_{R>}(\mathbf{0})}$. This makes possible to expand the driving field inside $\overline{B_{R>}(\mathbf{0})}$ in terms of regular VSWFs (with expansion coefficients $a_{\tau lm}$)

$$\mathbf{E}_{\text{inc}}(\mathbf{r}) = \sum_{\tau=1,2} \sum_{l=1}^{\infty} \sum_{m=-l}^{+l} a_{\tau lm} \mathbf{v}_{\tau lm}(\kappa \mathbf{r}).\tag{2.3.1}$$

The driving field induces response in the scatterer, and as described in Section 2.1, this response produces a unique solution to the scattered fields (with

Figure 2.2.1. Field patterns of a set of regular VSWFs of the two lowest multipole degrees, $l = 1, 2$, rearranged according to the irreducible representations of the point group D_{2h} . The images represent the electric field plotted in the planes along the x, y, z axes. For each VSWF, the electric field is either fully parallel, or fully perpendicular to the given plane (or zero). The parallel fields are represented by their magnitudes and field lines, the perpendicular by the perpendicular component.

the Silver-Müller constraints) $\mathbf{E}_{\text{sc}}(\mathbf{r})$. Outside the ball $\overline{B_{R<}(\mathbf{0})}$, the outgoing VSWFs provide a basis for the scattered field, so it can be expanded as (with expansion coefficients $f_{\tau lm}$)

$$\mathbf{E}_{\text{sc}}(\mathbf{r}) = \sum_{\tau=1,2} \sum_{l=1}^{\infty} \sum_{m=-l}^{+l} (f_{\tau lm} \mathbf{u}_{\tau lm}(\kappa \mathbf{r})). \quad (2.3.2)$$

hence we see that in the spherical shell $B_{R<,R>}(\mathbf{0})$ the total electric field can be expanded as

$$\mathbf{E}(\mathbf{r}) = \mathbf{E}_{\text{inc}}(\mathbf{r}) + \mathbf{E}_{\text{sc}}(\mathbf{r}) \quad (2.3.3)$$

$$= \sum_{\tau=1,2} \sum_{l=1}^{\infty} \sum_{m=-l}^{+l} (a_{\tau lm} \mathbf{v}_{\tau lm}(\kappa \mathbf{r}) + f_{\tau lm} \mathbf{u}_{\tau lm}(\kappa \mathbf{r})). \quad (2.3.4)$$

The VSWF expansion of the corresponding magnetic field follows from the VSWF definitions and the Faraday's law, yielding

$$\begin{aligned} \mathbf{H}(\mathbf{r}) &= \mathbf{H}_{\text{inc}}(\mathbf{r}) + \mathbf{H}_{\text{sc}}(\mathbf{r}) \\ &= \frac{1}{iZ_{\text{bg}}} \sum_{\tau=1,2} \sum_{l=1}^{\infty} \sum_{m=-l}^{+l} (a_{\tau lm} \mathbf{v}_{(3-\tau)lm}(\kappa \mathbf{r}) + f_{\tau lm} \mathbf{u}_{(3-\tau)lm}(\kappa \mathbf{r})), \end{aligned}$$

where $Z_{\text{bg}} = \sqrt{\mu_{\text{bg}}/\varepsilon_{\text{bg}}}$ is the wave impedance of the background.

From the assumed linearity of the system, there exists a linear relation between the expansion coefficients

$$f_{\tau lm} = \sum_{\tau' l' m'} T_{\tau lm; \tau' l' m'} a_{\tau' l' m'} \quad (2.3.5)$$

where $T_{\tau lm; \tau' l' m'} = T_{\tau lm; \tau' l' m'}(\omega)$ are the elements of the *transition matrix*, a.k.a. *T-matrix*. It completely describes the scattering properties of a linear scatterer outside its circumscribed sphere, so with the knowledge of the *T-matrix* we can solve the single-particle scattering problem simply by substituting appropriate expansion coefficients $a_{\tau' l' m'}$ of the driving field into (2.3.5). The outgoing VSWF expansion coefficients $f_{\tau lm}$ are the effective induced electric ($\tau = 2$) and magnetic ($\tau = 1$) multipole polarisation amplitudes of the scatterer, and this is why we sometimes refer to the corresponding VSWFs as to the electric and magnetic VSWFs, respectively.

For later convenience, we introduce a short-hand matrix notation for the expansion coefficients and related quantities, so that we do not need to write the indices explicitly; so for example, Eq. (2.3.5) would be written as $\mathbf{f} = \mathbf{T} \mathbf{a}$, where $\mathbf{a}, \mathbf{f}$ are column vectors with the expansion coefficients. Transposed and complex-conjugated matrices are labeled with the $\dagger$ superscript.

2.3.1 Power transport

Assuming the field expansion (2.3.3) in $B_{R<,R>}(\mathbf{0})$, the net power transported from $B_{R<}(\mathbf{0})$ to $B_{R<,R>}(\mathbf{0})$ by electromagnetic radiation can be shown to

be [25, sect. 7.3]

$$P = \frac{1}{2\kappa^2 Z_{\text{bg}}} \left(\Re(a^\dagger f) + \|f\|^2 \right) = \frac{1}{2\kappa^2 Z_{\text{bg}}} a^\dagger \left(T^\dagger T + \frac{T^\dagger + T}{2} \right) a. \quad (2.3.6)$$

Here P is well-defined only when $\kappa^2 \eta$ is real. We use this result later in Section 2.3.4 to formulate a practical sanity check for numerical computations.

2.3.2 Plane wave scattering; cross sections

In many scattering problems considered in practice, the driving field is at least approximately a plane wave. A transversal ($\hat{\mathbf{k}} \cdot \mathbf{E}_0 = 0$) plane wave propagating in direction $\hat{\mathbf{k}}$ with (complex) amplitude $\mathbf{E}_0$ can be expanded into regular VSWFs [25, 7.7.1] as

$$\mathbf{E}_{\text{PW}}(\mathbf{r}, \omega) = \mathbf{E}_0 e^{i\kappa \hat{\mathbf{k}} \cdot \mathbf{r}} = \sum_{\tau, l, m} a_{\tau lm}(\hat{\mathbf{k}}, \mathbf{E}_0) \mathbf{v}_{\tau lm}(\kappa \mathbf{r}),$$

where the expansion coefficients are obtained from the scalar products of the amplitude and corresponding dual vector spherical harmonics

$$\begin{aligned} a_{1lm}(\hat{\mathbf{k}}, \mathbf{E}_0) &= 4\pi i^l \mathbf{A}'_{1,l,m}(\hat{\mathbf{k}}) \cdot \mathbf{E}_0, \\ a_{2lm}(\hat{\mathbf{k}}, \mathbf{E}_0) &= -4\pi i^{l+1} \mathbf{A}'_{2,l,m}(\hat{\mathbf{k}}) \cdot \mathbf{E}_0. \end{aligned} \quad (2.3.7)$$

With the T -matrix and expansion coefficients of plane waves in hand, we can state the expressions for cross-sections of a single scatterer. Assuming a non-lossy background medium, extinction, scattering and absorption cross sections of a single scatterer irradiated by a plane wave propagating in direction $\hat{\mathbf{k}}$ and (complex) amplitude $\mathbf{E}_0$ are respectively [25, sect. 7.8.2]

$$\sigma_{\text{ext}}(\hat{\mathbf{k}}) = -\frac{1}{\kappa^2 \|\mathbf{E}_0\|^2} \Re(a^\dagger f) = -\frac{1}{2\kappa^2 \|\mathbf{E}_0\|^2} a^\dagger (T + T^\dagger) a, \quad (2.3.8)$$

$$\sigma_{\text{scat}}(\hat{\mathbf{k}}) = \frac{1}{\kappa^2 \|\mathbf{E}_0\|^2} \|f\|^2 = \frac{1}{\kappa^2 \|\mathbf{E}_0\|^2} a^\dagger (T^\dagger T) a, \quad (2.3.9)$$

$$\begin{aligned} \sigma_{\text{abs}}(\hat{\mathbf{k}}) &= \sigma_{\text{ext}}(\hat{\mathbf{k}}) - \sigma_{\text{scat}}(\hat{\mathbf{k}}) = -\frac{1}{\kappa^2 \|\mathbf{E}_0\|^2} \left(\Re(a^\dagger f) + \|f\|^2 \right) \\ &= -\frac{1}{\kappa^2 \|\mathbf{E}_0\|^2} a^\dagger \left(T^\dagger T + \frac{T^\dagger + T}{2} \right) a, \end{aligned} \quad (2.3.10)$$

where $a = a(\mathbf{k}, \mathbf{E}_0)$ is the vector of plane wave expansion coefficients as in (2.3.7).

2.3.3 Computing the T-matrix

In principle, the T-matrix elements for a compact scatterer can be computed by any method that is capable to solve the scattering problem from Section 2.1 with Silver-Müller conditions for the scattered fields $\mathbf{E}_{\text{sc}}, \mathbf{H}_{\text{sc}}$ by the following procedure:

1. Set $\mathbf{E}_{\text{inc}}(\mathbf{r}) = \mathbf{v}_{\tau'l'm'}(\kappa\mathbf{r})$.
2. Choose a radius R such that the whole scatterer fits inside the open ball $B_R(\mathbf{0})$ and evaluate the scattered electric field on the bounding sphere $\mathbf{E}_{\text{sc}}(R\hat{\mathbf{r}})$.
3. Evaluate the inner products

$$-i\kappa \int (\mathbf{v}_{\tau lm} \times (\nabla \times \mathbf{E}_{\text{sc}}) - \mathbf{E}_{\text{sc}} \times (\nabla \times \mathbf{v}_{\tau lm})) \cdot \hat{\mathbf{r}} R^2 d\Omega = T_{\tau lm; \tau' l' m'}$$

to get a column of the T-matrix.

The validity of this procedure follows from the uniqueness theorem for the external boundary problem (the scattered fields in $\mathbb{R}^3 \setminus B_R(\mathbf{0})$ are completely determined by the data $\mathbf{E}_{\text{sc}}(R\hat{\mathbf{r}})$ and the orthogonality and reciprocity relations for VSWFs (for details, see [25, Section 7.3]).

T -matrices of particles with certain simple geometries (most famously spherical) can be obtained analytically [25, 41]; for particles with smooth surfaces one can find them numerically using the *null-field method* [25, 42, 43]. The null-field method works well in the many typical cases, but for less common parameter ranges (such as concave shapes, extreme values of aspect ratios or relative refractive index) they might suffer from serious numerical instabilities [44, Sect. 5.8.4].

Remark 6. In the numerical results presented later in Section 3, the T -matrices are typically evaluated using the null-field method, which is relatively fast for axially symmetric particles and is implemented in the QPMS suite (introduced later in Section 2.8). For more complicated scatterers, we could use the `scuff-tmatrix` tool from the free software SCUFF-EM suite [3, 45], which evaluates the T -matrices using the boundary element method.

Note that the upstream versions of SCUFF-EM contain a bug that renders almost all T -matrix results wrong; I found and fixed the bug in my fork available at <https://github.com/tecnokrates/scuff-em> in revision g78689f5. However, as of August 2020, the bugfix has still not been merged into upstream. My fork of SCUFF-EM corrects also several other bugs, but these are less critical for T-matrix calculations.

2.3.4 T-matrix sanity checks

In practice, it is nontrivial to determine whether a numerically obtained T-matrix is correct or not (e.g. due to multipole truncation, numerical inaccuracies, or possible bugs). However, there are some early warning techniques to catch some of the potential problems.

One of them is to check whether the T-matrix has the symmetries corresponding to the actual physical symmetries of the scatterer, meaning that the T-matrix can be block-diagonalised according to the irreducible representation subspaces of the physical system's symmetry group. For details, see [46].

This approach can be also be reversed – the projector operators from the block-diagonalisation procedure can be used to a posteriori “sweep out” the asymmetries introduced by the method that generated the T-matrix (e.g. due to a nonconforming mesh).

Another practical sanity check for a T-matrix of an arbitrarily shaped scatterer (in nondispersive background) is based on the power transport formula (2.3.6). For a particle without internal gain, the net radiated power can never be positive, regardless of the driving field; as a consequence, hence the “power matrix”

$$\Pi = T^\dagger T + (T^\dagger + T) / 2 \quad (2.3.11)$$

(which is obviously hermitian), must be negative (semi-)definite.

Example 7. Consider a golden cylindrical nanoparticle with 50 nm radius and 50 nm height inside a medium with constant real refractive index of 1.52. We employ two different approaches to compute its T-matrix; in both cases, we use $l_{\max} = 3$ as the multipole cutoff for the final T-matrix:

1. The `scuff-tmatrix` tool from the patched SCUFF-EM suite, which uses BEM. Gold permittivity is interpolated from tabulated values [47].
2. The T-matrix generator for axially symmetric particles from QPMS, which is based on the null-field equations. The intermediate matrices Q, R are calculated up to multipole degree $l_{\max} = 6$. Gold permittivity is obtained from Drude-Lorentz model with parameters from [48].

Fig. 2.3.1 shows the eigenvalues of the power matrix (2.3.11) for the two cases in frequency range 0.6–2.15 eV/ $\hbar$. Both models predict resonances at similar frequencies, as can be seen from the two prominent minima, although their strengths slightly differ (which is not surprising also due to the different material model). At higher frequencies, the negative (semi-)definitivity of the power matrix resulting from BEM T-matrix is violated (it has non-negligible positive eigenvalues). This means that for certain driving fields, the particle would generate energy, which is unphysical. The T-matrix obtained with the null-field method would therefore be more suitable to be used in further computations.

2.4 Multiple scattering

Let us consider a LOSIB with a foreground that can be partitioned into multiple spherically separated scatterers (SSSs). Let $\mathcal{P}$ be an index set labeling the scatterers. We enclose each of the SSSs into closed balls $\overline{B_{R_p}(\mathbf{r}_p)}$ so that they do not intersect, $\overline{B_{R_p}(\mathbf{r}_p)} \cap \overline{B_{R_q}(\mathbf{r}_q)} = \emptyset; p, q \in \mathcal{P}$, so there is a non-empty separating spherical shell (SSSh) $B_{R_p, R_p^>}(\mathbf{r}_p)$ around each one that contains only the background medium. Then the EM field inside each SSSh can be

Figure 2.3.1. Power matrix based sanity check of a T-matrix computed using BEM (left) with the displayed mesh, and using null-field equations (right). In both cases, multipole cutoff was set to $l_{\max} = 3$. Note the non-negligible positive eigenvalues in the matrix obtained with BEM.

expanded in a way similar to (2.3.3), using VSWFs with origins shifted to the centre of the volume:

$$\mathbf{E}(\omega, \mathbf{r}) = \sum_{\tau=1,2} \sum_{l=1}^{\infty} \sum_{m=-l}^{+l} (a_{p,\tau lm} \mathbf{v}_{\tau lm}(\kappa(\mathbf{r} - \mathbf{r}_p)) + f_{p,\tau lm} \mathbf{u}_{\tau lm}(\kappa(\mathbf{r} - \mathbf{r}_p))), \quad (2.4.1)$$

$$\mathbf{r} \in \Theta_{R_p, R_p^>}(\mathbf{r}_p).$$

Unlike the single scatterer case, the incident field coefficients $a_{p,\tau lm}$ here are not only due to some external driving field that the particle does not influence but they also contain the contributions of fields scattered from *all other scatterers*:

$$a_p = \tilde{a}_p + \sum_{q \in \mathcal{P} \setminus \{p\}} \mathcal{S}_{p \leftarrow q} f_q \quad (2.4.2)$$

where $\tilde{a}_p$ represents the part due to the external driving that the scatterers can not influence, and $\mathcal{S}_{p \leftarrow q}$ is a *translation operator* defined below in Sec. 2.4.1, that contains the re-expansion coefficients of the outgoing waves in origin $\mathbf{r}_q$ into regular waves in origin $\mathbf{r}_p$. For each scatterer, we also have its *T*-matrix relation as in (2.3.5),

$$f_q = T_q a_q.$$

Together with (2.4.2), this gives rise to a set of linear equations

$$f_p - T_p \sum_{q \in \mathcal{P} \setminus \{p\}} \mathcal{S}_{p \leftarrow q} f_q = T_p \tilde{a}_p, \quad p \in \mathcal{P} \quad (2.4.3)$$

which defines the multiple-scattering problem. If all the p, q -indexed vectors and matrices (note that without truncation, they are infinite-dimensional) are arranged into blocks of even larger vectors and matrices, this can be written in a short-hand form

$$(I - T\mathcal{S})f = T\tilde{a} \quad (2.4.4)$$

where I is the identity matrix, T is a block-diagonal matrix containing all the individual T -matrices, and $\mathcal{S}$ contains the individual $\mathcal{S}_{p \leftarrow q}$ -matrices as the off-diagonal blocks, whereas the diagonal blocks are set to zeros.

In practice, the multiple-scattering problem is solved in its truncated form, in which all the l -indices related to a given scatterer p are truncated as $l \leq L_p$, leaving only $N_p = 2L_p(L_p + 2)$ different τlm -multi-indices left. The truncation degree can vary for different scatterers (e.g. due to different physical sizes), so the truncated block $[\mathcal{S}_{p \leftarrow q}]_{l_q \leq L_q; l_p \leq L_p}$ has shape $N_p \times N_q$, not necessarily square.

If no other type of truncation is done, there remain $2L_p(L_p + 2)$ different τlm -multi-indices for the p -th scatterer, so that the truncated version of the matrix $(I - T\mathcal{S})$ is a square matrix with $(\sum_{p \in \mathcal{P}} N_p)^2$ elements in total. The truncated problem (2.4.4) can then be solved using standard numerical linear algebra methods (typically, by LU factorisation of the $(I - T\mathcal{S})$ matrix at a given frequency, and then solving with Gauss elimination for as many different incident $\tilde{a}$ vectors as needed).

2.4.1 Translation operator

Let $\mathbf{r}_1, \mathbf{r}_2$ be two different origins; a regular VSWF with origin $\mathbf{r}_1$ can be expanded in terms of regular VSWFs with origin $\mathbf{r}_2$ as follows:

$$\mathbf{v}_{\tau lm}(\kappa(\mathbf{r} - \mathbf{r}_1)) = \sum_{\tau' l' m'} \mathcal{R}_{\tau lm; \tau' l' m'}(\kappa(\mathbf{r}_2 - \mathbf{r}_1)) \mathbf{v}_{\tau' l' m'}(\kappa(\mathbf{r} - \mathbf{r}_2)), \quad (2.4.5)$$

where the regular translation operator $\mathcal{R}$ can be written in terms of scalar spherical wavefunction as in Eq. (2.4.11) below. For singular (outgoing) waves, the form of the expansion differs inside and outside the ball $\overline{B_{\|\mathbf{r}_2 - \mathbf{r}_1\|}(\mathbf{r}_1)}$:

$$\begin{aligned} \mathbf{u}_{\tau lm}(\kappa(\mathbf{r} - \mathbf{r}_1)) &= \\ &= \begin{cases} \sum_{\tau' l' m'} \mathcal{S}_{\tau lm; \tau' l' m'}(\kappa(\mathbf{r}_2 - \mathbf{r}_1)) \mathbf{v}_{\tau' l' m'}(\kappa(\mathbf{r} - \mathbf{r}_2)), & \mathbf{r} \in B_{\|\mathbf{r}_1 - \mathbf{r}_2\|}(\mathbf{r}_2) \\ \sum_{\tau' l' m'} \mathcal{R}_{\tau lm; \tau' l' m'}(\kappa(\mathbf{r}_2 - \mathbf{r}_1)) \mathbf{u}_{\tau' l' m'}(\kappa(\mathbf{r} - \mathbf{r}_2)), & \mathbf{r} \notin \overline{B_{\|\mathbf{r}_1 - \mathbf{r}_2\|}(\mathbf{r}_2)} \end{cases}, \end{aligned} \quad (2.4.6)$$

where the singular translation operator $\mathcal{S}$ has the same form as $\mathcal{R}$ in (2.4.11) except the regular spherical Bessel functions j_l are replaced with spherical Hankel functions $h_l^{(1)}$.

As MSTMM deals most of the time with the *expansion coefficients* of fields $a_{p, \tau lm}, f_{p, \tau lm}$ in different origins $\mathbf{r}_p$ rather than with the VSWFs directly, let us

write down how *they* transform under translation. We assume the field can be expressed in terms of regular waves everywhere, and expand it in two different origins $\mathbf{r}_p, \mathbf{r}_q$,

$$\mathbf{E}(\mathbf{r}, \omega) = \sum_{\tau, l, m} a_{p, \tau l m} \mathbf{v}_{\tau l m}(\kappa(\mathbf{r} - \mathbf{r}_p)) = \sum_{\tau', l', m'} a_{q, \tau' l' m'} \mathbf{v}_{\tau' l' m'}(\kappa(\mathbf{r} - \mathbf{r}_q)).$$

Re-expanding the waves around $\mathbf{r}_p$ in terms of waves around $\mathbf{r}_q$ using (2.4.5),

$$\mathbf{E}(\mathbf{r}, \omega) = \sum_{\tau, l, m} a_{p, \tau l m} \sum_{\tau' l' m'} \mathcal{R}_{\tau l m; \tau' l' m'}(\kappa(\mathbf{r}_q - \mathbf{r}_p)) \mathbf{v}_{\tau' l' m'}(\kappa(\mathbf{r} - \mathbf{r}_q))$$

and comparing to the original expansion around $\mathbf{r}_q$, we obtain

$$a_{q, \tau' l' m'} = \sum_{\tau, l, m} \mathcal{R}_{\tau l m; \tau' l' m'}(\kappa(\mathbf{r}_q - \mathbf{r}_p)) a_{p, \tau l m}. \quad (2.4.7)$$

For the sake of readability, we introduce a shorthand matrix form for (2.4.7)

$$a_q = \mathcal{R}_{q \leftarrow p} a_p \quad (2.4.8)$$

(note the reversed indices) Similarly, if we had only outgoing waves in the original expansion around $\mathbf{r}_p$, we would get

$$a_q = \mathcal{S}_{q \leftarrow p} f_p \quad (2.4.9)$$

for the expansion inside the ball $B_{\|\mathbf{r}_q - \mathbf{r}_p\|}(\mathbf{r}_p)$ and

$$f_q = \mathcal{R}_{q \leftarrow p} f_p \quad (2.4.10)$$

outside.

The regular translation operator elements can be expressed explicitly as

$$\mathcal{R}_{\tau l m; \tau' l' m'}(\mathbf{d}) = \sum_{\lambda=|l-l'|+|\tau-\tau'|}^{l+l'} C_{\tau l m; \tau' l' m'}^{\lambda} Y_{\lambda, m-m'}(\hat{\mathbf{d}}) j_{\lambda}(d), \quad (2.4.11)$$

and analogously the elements of the singular operator $\mathcal{S}$, having spherical Hankel functions ($h_l^{(1)} = j_l + i y_l$) in the radial part instead of the regular Bessel functions,

$$\mathcal{S}_{\tau l m; \tau' l' m'}(\mathbf{d}) = \sum_{\lambda=|l-l'|+|\tau-\tau'|}^{l+l'} C_{\tau l m; \tau' l' m'}^{\lambda} Y_{\lambda, m-m'}(\hat{\mathbf{d}}) h_{\lambda}^{(1)}(d). \quad (2.4.12)$$

Explicit expressions for the rather complicated constant factors $C_{\tau l m; \tau' l' m'}^{\lambda}$ can be found in Publication I.

With a proper normalisation convention for VSWFs, the regular translation operator is unitary, $(\mathcal{R}_{\tau l m; \tau' l' m'}(\mathbf{d}))^{-1} = \mathcal{R}_{\tau l m; \tau' l' m'}(-\mathbf{d}) = \mathcal{R}_{\tau' l' m'; \tau l m}^*(\mathbf{d})$, or in the per-particle matrix notation,

$$\mathcal{R}_{q \leftarrow p}^{-1} = \mathcal{R}_{p \leftarrow q} = \mathcal{R}_{q \leftarrow p}^{\dagger}. \quad (2.4.13)$$

Note that truncation at finite multipole degree breaks the unitarity, $[\mathcal{R}_{q \leftarrow p}]_{l \leq L}^{-1} \neq [\mathcal{R}_{p \leftarrow q}]_{l \leq L} = [\mathcal{R}_{q \leftarrow p}^\dagger]_{l \leq L}$, which has to be taken into consideration when evaluating quantities such as absorption or scattering cross sections. Similarly, the full regular operators can be composed

$$\mathcal{R}_{a \leftarrow c} = \mathcal{R}_{a \leftarrow b} \mathcal{R}_{b \leftarrow c} \quad (2.4.14)$$

but truncation breaks this, $[\mathcal{R}_{a \leftarrow c}]_{l \leq L} \neq [\mathcal{R}_{a \leftarrow b}]_{l \leq L} [\mathcal{R}_{b \leftarrow c}]_{l \leq L}$.

2.4.2 Green's function for the multiple-scattering system

One can use the MSTMM formalism above to calculate the electric Green's function in the full LOSIB with scatterers; as opposed to the Green's functions of the background only. Among other applications, this gives the couplings between point quantum emitters in the macroscopic QED approach, as discussed later in Chapter 4.

To make the distinction for the purposes of this section, let us denote $\overleftrightarrow{G}_0$ the electric Green's function in the homogeneous system with the background medium only, as in (1.1.7); and let $\overleftrightarrow{G}$ stand for the Green's function of the system with scatterers. The background Green's function $\overleftrightarrow{G}_0$ is directly proportional to the electric dipole block of the outgoing VSWF translation operator: on a vector $\mathbf{w}$, the Green's function tensor (dyadic) acts like

$$\overleftrightarrow{G}_0(\kappa; \mathbf{r}_2, \mathbf{r}_1) \cdot \mathbf{w} = \sum_{m'=-1}^1 \underbrace{\sum_{m=-1}^1 F_m(\mathbf{w}) \mathcal{S}_{2,1,m;2,1,m'}(\kappa(\mathbf{r}_2 - \mathbf{r}_1))}_{a_{\mathbf{r}_2;21m'}^{(0)}} v_{21m'}(0) \quad (2.4.15)$$

where the set of linear forms F_m depends on the normalisation and phase conventions for VSWFs; up to a scalar constant γ , it is dual to the values of regular dipole VSWFs in the origin, $F_m \left(\left(\sum_{p \in \mathcal{P}} N_p \right)^2 v_{21m'}(0) \right) = \gamma \delta_{mm'}$. The values $F_m(\mathbf{w})$ here can be interpreted as the VSWF excitation coefficients of an imaginary point scatterer ($\mathbf{w}$ -oriented dipolar source) located at $\mathbf{r}_1$, and the expression $a_{\mathbf{r}_2;21m'}^{(0)}$ on the right hand side of (2.4.15) represents the coefficients of its VSWF re-expansion at $\mathbf{r}_2$ to the lowest (i.e. electric dipole) order.

If we add a set of actual spherically separated scatterers into the system (such that $\mathbf{r}_1, \mathbf{r}_2$ are outside their circumscribed spheres), the field caused by the source at $\mathbf{r}_1$ can be expanded around the p -th scatterer in terms of VSWFs with coefficients

$$\tilde{a}_{p,\tau'l'm'}(\mathbf{r}_1; \mathbf{w}) = \sum_{m=-1}^1 F_m(\mathbf{w}) \mathcal{S}_{2,1,m;\tau',l',m'}(\kappa(\mathbf{r}_p - \mathbf{r}_1)),$$

the collection of which will serve as the right hand side for the multiple scattering problem (2.4.4), which can be solved to yield the scatterers' outgoing

VSWF coefficient vector $f(\mathbf{r}_1; \mathbf{w})$. Propagating the outgoing waves to $\mathbf{r}_2$ and adding their contributions to (2.4.15) gives the values of the full Green's function,

$$\begin{aligned} \overleftrightarrow{G}(\kappa; \mathbf{r}_2, \mathbf{r}_1) \cdot \mathbf{w} = & \overleftrightarrow{G}_0(\kappa; \mathbf{r}_2, \mathbf{r}_1) \cdot \mathbf{w} + \\ & + \sum_{m'=-1}^1 \sum_{p \in \mathcal{P}} \sum_{\tau lm} f_{p, \tau lm}(\mathbf{r}_1; \mathbf{w}) \mathcal{S}_{\tau lm; 2, 1, m'}(\kappa(\mathbf{r}_2 - \mathbf{r}_p)) \mathbf{v}_{21m'}(0). \end{aligned} \quad (2.4.16)$$

2.5 Infinite periodic systems

MSTMM can be extended to handle systems with infinite periodic foreground, as long as it can be partitioned into spherically separated scatterers. Although large finite systems are where MSTMM excels the most, there are several reasons that makes its extension to infinite lattices (where periodic boundary conditions might be applied) desirable as well. Other methods might be already fast enough, but MSTMM will be faster in most cases in which there is enough spacing between the neighboring scatterers. MSTMM works well with any space group symmetry the system might have (as opposed to, for example, FDTD with a cubic mesh applied to a honeycomb lattice), which makes e.g. application of group theory in mode analysis quite easy. Finally, having a method that handles well both infinite and large finite systems gives a possibility to study finite-size effects in periodic scatterer arrays.

MSTMM for periodic systems starts from the usual Bloch wave approach; for a given Bloch quasi-momentum $\mathbf{k}$, both the driving field and scattered field VSWF coefficients are assumed to have a quasiperiodic form $\tilde{a}_{\mathbf{n}, \alpha} = \tilde{a}_{\mathbf{0}, \alpha}(\mathbf{k}) e^{i\mathbf{k} \cdot \mathbf{R}_{\mathbf{n}}}$, $f_{\mathbf{n}, \alpha} = f_{\mathbf{0}, \alpha}(\mathbf{k}) e^{i\mathbf{k} \cdot \mathbf{R}_{\mathbf{n}}}$, where $\mathbf{n}$ is a unit cell index, $\mathbf{R}_{\mathbf{n}}$ is a lattice vector, and the index $\alpha \in \mathcal{P}$ distinguishes between different scatterers in a single unit cell.

The multiple scattering problem 2.4.4 can then be rewritten in terms of the VSWF expansion coefficients of fields around scatterers in a representative unit cell,

$$f_{\mathbf{0}, \alpha}(\mathbf{k}) - T_{\alpha} \sum_{\beta \in \mathcal{P}} W_{\alpha\beta}(\mathbf{k}) f_{\mathbf{0}, \beta}(\mathbf{k}) = T_{\alpha} \tilde{a}_{\mathbf{0}, \alpha}(\mathbf{k}) \quad (2.5.1)$$

where $W_{\alpha\beta}(\mathbf{k})$ is a lattice sum of the VSWF translation operator modulated according to the Bloch vector $\mathbf{k}$

$$W_{\alpha\beta}(\mathbf{k}) \equiv \sum_{\mathbf{m} \in \mathbb{Z}^d} (1 - \delta_{\alpha\beta} \delta_{\mathbf{m}\mathbf{0}}) \mathcal{S}_{\mathbf{0}, \alpha \leftarrow \mathbf{m}, \beta} e^{i\mathbf{k} \cdot \mathbf{R}_{\mathbf{m}}}. \quad (2.5.2)$$

Computation of $W_{\alpha\beta}(\mathbf{k})$ represents the main challenge of MSTMM for infinite periodic systems due to bad convergence properties of the lattice sum. However, there fortunately exist advanced analytical techniques which enable transforming the lattice sums into an exponentially convergent form, as explained in Section 3.3 of Publication I.

2.6 Resonances and lattice modes

If we set the right hand side of the equation 2.5.1 to zero,

$$(I - TW) f_0(\mathbf{k}) = 0, \quad (2.6.1)$$

we can solve it to get the solutions that propagate in the system without requiring external driving, that is, the *lattice modes*. Non-trivial solutions exist if the matrix on the left-hand side $M(\omega, \mathbf{k}) = I - T(\omega) W(\omega, \mathbf{k})$ is singular – this condition gives the *dispersion relation* for the periodic structure. In plasmonic lattices, the scatterers are absorbing, so the frequencies ω will be in general complex, which makes finding the dispersion relations nontrivial. However, if the matrix $M(\omega, \mathbf{k})$ is an analytic function of ω , it is possible to use the Beyn’s algorithm or another nonlinear eigensolver based on contour integration [49, 50] that makes the solution quite efficient, see Publication I, Section 3.2.

Analogous approach can be used for finding the resonances of finite arrays when the right hand side is set to zero in Eq. (2.4.4). We apply this to study the quality factor of array resonances later in Section (3.2).

2.7 Symmetries

In Publication I, we show that if the LOSIB in question has nontrivial symmetries, they can be exploited to significantly reduce the computational costs of solving the multiple-scattering T-matrix problem. The idea is to find a unitary transformation and symmetry-adapted basis for the excitation coefficient vectors f that takes the interaction matrix $(I - TS)$ from the multiple-scattering problem (2.4.4) into a block-diagonal form. Each of the blocks corresponds to an irreducible representation of the LOSIB symmetry group. The linear algebraic operations then proceed on each block separately.

If the original truncated system has N degrees of freedom and is factorised into s almost equally large blocks, a single LU factorisation with time complexity $O(N^3)$ and memory complexity $O(N^2)$ can be replaced with s LU factorisations, each requiring $O(N^3/s^3)$ time and $O(N^2/s^2)$, reducing the total time and memory demands both by the factor of s^{-2} (if the s LU factorisations are done in sequence). This significantly increases the number of scatterers that MSTMM can handle in practice.

Symmetries also play an important role in lattice mode analysis of infinite periodic systems. Among other things, they give a priori knowledge about the modes and their degeneracies in high-symmetry parts of the Brillouin zone, as used in Publications III and IV. The technical details of exploiting the symmetries in MSTMM for both finite and periodic infinite systems are explained in Publication I.

2.8 The QPMS suite

The *QPMS Photonic Multiple Scattering suite* is a software package that implements MSTMM including the enhancements introduced in sections 2.5 and 2.7. It contains a core C library, Python bindings and several utilities for routine computations, such as scattering cross sections under plane wave irradiation or lattice modes of two-dimensional periodic arrays. QPMS includes Doxygen documentation together with description of the API. It has been written with customisability and extendibility in mind, so that including e.g. alternative methods of *T*-matrix calculations of a single particle's matrix are as easy as possible.

QPMS is a free software published under the GNU General Public License version 3. The source code can be downloaded from the public git repository <https://repo.or.cz/qpms.git>. An online version of the Doxygen documentation is available at <https://qpms.necada.org>.

2.9 Further extensions

The “pure” MSTMM as described in the previous sections of this chapter (and currently implemented in QPMS) has certain limitations: It requires an unbounded LOSIB with a foreground that can be divided into mutually disjoint SSSs. Moreover, by itself it can only evaluate fields *outside* the scatterers' circumscribed spheres. While some of these limitations are too fundamental to overcome (mainly that the system has to be a LOSIB), others can be lifted using a realistic amount of effort. Moreover, some problems can be solved more efficiently using another approach than LU factorisation in the linear system (2.4.4). Some of the possible extensions are discussed in this section.

2.9.1 Combining with other integral methods

If the foreground contains SSSs but cannot be completely divided into SSSs in a meaningful way, MSTMM can be combined with another integral method, such as BEM or some approach based on the volume integral equations, resulting into linear system of the form (1.2.4) in which the unknowns are both multipolar excitations for the SSSs and whichever degrees of freedom the other method works with (such as surface or volume current elements) for the rest of the system. The couplings inside the two subsystems are obtained in their respective usual manners. As long as the other method is based on the usual Green's dyadics, the couplings between the two subsystems are also straightforward, and exponentially convergent lattice sums can then still be used for periodic systems. In the worst case scenario, this basic hybrid approach brings at least the typical DoF reduction for the SSS subsystem.

2.9.1.1 Planarly layered media

Systems consisting of compact scatterers and parallel planar layers make a special but significant class: for example, they include a typical experimental situation where a nanoparticle array lies on a planar substrate with different refractive index than the environment above the array. The interactions between the nanoparticles are then no longer given simply just by the translation operators (2.4.9), and the interactions with the interfaces have to be included. Fortunately, it is feasible to do this without increasing the number of unknowns (i.e. the interfaces themselves do not need their own degrees of freedom): the transmissions and reflections from the interfaces just lead to additional interaction terms between the compact scatterers [51]. The new interaction terms contain Sommerfeld integrals, which makes their evaluation more difficult than the translation operators in homogeneous medium (whose elements do have closed-form expressions, although complicated). Efficient methods of evaluating Sommerfeld integrals can be found in the literature [52].

2.9.1.2 Retrieving internal fields

In fact, MSTMM uses a hybrid approach from the very beginning: the T-matrix of a single SSS has to be retrieved by another method (as described in Section 2.3.3), and MSTMM itself handles only the mutual interactions between the SSSs and the fields outside their circumscribed spheres (in practice—due to truncation—only the fields *far enough* from the circumscribed spheres).

However, after solving the multiple-scattering problem with MSTMM, the method used to compute the T-matrix can be plugged in again in order to retrieve the internal fields, and the near fields possibly lost in MSTMM due to the multipole truncation.

Let $\mathbf{E}_{\text{sc}(p,\tau lm)}$ be the electric fields that solve the single-scatterer problem involving only the p -th scatterer placed in the origin with the incident fields in the form of regular VSWFs, $\mathbf{E}_{\text{inc}(p,\tau lm)} = \mathbf{v}_{\tau lm}$, and let $a_{p,\tau lm}$ be the incoming VSWF coefficients for the p -th scatterer obtained by MSTMM for the whole multiple-scattering problem with incident field $\mathbf{E}_{\text{inc}}$. The total electric field is then

$$\begin{aligned} \mathbf{E}(\mathbf{r}) &= \mathbf{E}_{\text{inc}}(\mathbf{r}) + \mathbf{E}_{\text{sc}}(\mathbf{r}) \\ &= \mathbf{E}_{\text{inc}}(\mathbf{r}) + \sum_{\tau lm} a_{p,\tau lm} \mathbf{E}_{\text{sc}(p,\tau lm)}(\mathbf{r} - \mathbf{r}_p) \\ &\quad + \sum_{p' \in \mathcal{P} \setminus \{p\}} \sum_{\tau' l' m'} f_{p',\tau' l' m'} \mathbf{u}_{\tau' l' m'}(\mathbf{r} - \mathbf{r}_{p'}). \end{aligned} \quad (2.9.1)$$

This expansion is valid everywhere outside the circumscribed spheres of the *other* scatterers $p' \in \mathcal{P} \setminus \{p\}$, and in particular, it is valid inside the p -th scatterer and in its vicinity. This enables numerical evaluation of the fields anywhere: far from the scatterers, the field is obtained using the “pure” MSTMM formula (2.3.2) as usual; and inside and nearby the scatterers, the field is evaluated using the hybrid formula (2.9.1). In practice, the multipole truncation will

cause a small jump in the field values between the regions where the different methods are used. If the regions cannot be chosen in a way that makes these jumps negligible, it is an indication that the multipole truncation has been excessive from the very beginning.

2.9.1.3 Multilevel fast multipole algorithm

MSTMM, as presented here, can be interpreted as a type of two-level fast multipole method (FMM) [26, 53–55]. FMM in general uses aggregation of degrees of freedom into larger blocks (according to position) in such way that the interaction between them can be, up to a desired accuracy, described by less numbers than the number of interactions between the DoFs in the original system. Here we have used aggregation at the level of single scatterer, and the DoFs have been aggregated into the electric and magnetic multipole excitations of the scatterer, the interactions between the aggregates (i.e. scatterers) are given by the truncated translation operators, which is computationally much more concise than e.g. interactions between all the surface elements in BEM, as discussed in Chapter 1.

In principle, the aggregation process can be repeated on multiple levels (e.g. the aggregated volumes from one level are joined to even larger aggregates on the higher level). As a result, one can get an algorithm that effectively enables to reduce the speed of matrix multiplication in systems like (1.2.4) from $O(N^2)$ to $O(N \log N)$, where N is the original number of DoFs. This gives a significant boost to solving the scattering problem on large systems with iterative methods (note that the matrix does not have to be saved in the memory). The algorithm is known as the *multilevel fast multipole algorithm* (MLFMA) [26].

One could imagine that in MSTMM the multilevel aggregation would correspond to assembling an aggregated T -matrix of several adjacent SSSs and treating them as one – the resulting T -matrix would have a cutoff at higher multipole degree than those of the elementary scatterers; however, this would work only if such aggregation yields scatterers that are again spherically separated (which is usually not the case); hence in practice, the aggregation has to be done in a more sophisticated way [26, 56]. There exists a free software implementation of such method [57] (but it does not have many of the other features that QPMS does).

3. T-matrix method in plasmonic nanoarrays

This chapter shows several applications of MSTMM and its implementation in QPMS, mostly building on the results contained in Publications II, III and IV. However, as all of them predate the implementation of some important features in the QPMS suite, such as the lattice sums and Beyn’s algorithm, some of the approaches used in the original articles might look deprecated in today’s view. Therefore, rather than restating the old results, I try to provide a current perspective, especially in the case of Publication II.

3.1 Lasing in nanoparticle arrays

Due to the assumption of linear constitutive relations that lies in the core of the T-matrix method, it cannot by itself explain the phenomena with nonlinear time dynamics such as lasing or Bose-Einstein condensation. However—sometimes combined with other methods or using suitable heuristics—it can give an unprecedented insight into experimentally observed phenomena, which, regardless of their internal nonlinearity, develop on top of lattice resonances that appear in the linear regime.

Lasing action requires high effective gain, which in periodic photonic structures normally peaks at the band edges where the group velocity is zero [58]. Such band edges are mostly guaranteed to exist in high-symmetry points of the Brillouin zone. The most obvious candidate is the Γ point corresponding to the zero Bloch vector, $\mathbf{k} = 0$, which is a high-symmetry point for most space groups with nontrivial point symmetries.

3.1.1 Γ point lasing in Ag nanoparticle square array

Let us consider the experiment performed in Publication II. The investigated system consists of a regular $100\,\mu\text{m} \times 100\,\mu\text{m}$ square array of cylindrical silver nanoparticles with the nominal height $h = 30\,\text{nm}$, radius $r = 30\,\text{nm}$ and periodicity $375\,\text{nm}$ along x and y directions, fabricated on a borosilicate substrate. The array is doped with a medium containing a fluorescent dye, Rhodamine

6G, and on top there is a cover glass. For modelling purposes, we assume that this gives a roughly homogeneous background medium with a constant refractive index of 1.52. The frequencies corresponding to the first diffraction order crossing at the Γ and the assumed band edge right below fall well within the emission spectrum of the dye, so when excited (by a light pump at higher frequency), the dye can emit photons into these lattice modes, serving as gain for lasing action.

When pumped, the array emits light at up to two frequencies at a time, depending on the pump fluence, see Figure 3.1.1, with two remarkably different spatial emission patterns:

- Case 1.* The emission with the higher pump threshold makes a single beam in the direction normal to the array plane. In real space images, the whole array appears bright, with some variation.
- Case 2.* The lower threshold emission shows two peaks along the k_x axis in the spectrometer image. Experiments with different array sizes (keeping the other parameters constant) show that the distance between the peaks is inversely proportionate to the length of the array side. Real space images show maximal intensities at two opposite edges of the array, while the middle of the array remains dark.

An explanation for the observations from Case 1 is that the lasing action happens in a *bright mode*, characteristic by in-phase electric dipolar excitations of the nanoparticles in the xy plane. Quite intuitively, such polarisation pattern creates the simple beam radiating in the normal (z) direction. In Case 2, on the other hand, the lasing happens in a *dark mode* in which the nanoparticles exhibit such multipole polarisation patterns that—in the ideal case of an infinite periodic lattice—would not radiate outside the array because far from the array, the field contributions from all the sources would cancel out due to symmetries. However, in the real-world case of a finite lattice, the light would “scatter from the edges”: both the real space and k -space measurements are consistent with the hypothesis that dipoles are induced near the edges, with π phase difference between the opposite sides.

The hypothesis can be tested using a multiple-scattering T-matrix simulation by choosing the multipole polarisation pattern expected for the dark or bright mode in question, and setting this as the right-hand side of the scattering equation (2.4.4) for *all* nanoparticles. Such a simulation is depicted in Figure 3.1.2. The driving of each particle with the electric xy quadrupole wave $\mathbf{v}_{22-2} - \mathbf{v}_{22+2}$ (irreducible representation A'_2 ; see also Figure 2.2.1) gives results that are in excellent qualitative agreement with the experimental observation, supporting the hypothesis about the electric xy quadrupole character of the output. However, this is not conclusive, as the constant driving of each particle with the magnetic z dipole wave $\mathbf{v}_{110}$ (irrep B'_2) would produce the same qualitative results for the x and y linear polarisations. To distinguish these patterns from each other, it

would be necessary to perform the measurements also with output polarisation filter along one of the diagonal directions.

Remark 8. In reality, we cannot expect that the lasing action would happen in such homogeneous manner throughout the array. Hence instead of setting the right hand side of (2.4.4) as constant for all particles, we might rather choose to drive only some particles, for example in certain region, and observe whether the results qualitatively change. This is also a good indicator of the level of spatial coherence that the array can offer.

3.1.2 Γ point lasing in Ni nanoparticle rectangular arrays

Arrays with nanoparticles made of ferromagnetic materials, such as nickel, are interesting due to their magneto-optic properties [23]. Publication III studies lasing in rectangular (and square) arrays of nickel nanoparticles. The array periodicity in the x direction in all samples is $p_x = 380$ nm, while in the y direction, the periodicities are $p_y = 370, 375, 380, 385$, and 390 nm; nominal radius of the particles is 30 nm and height 60 nm. Otherwise the samples have similar properties as in the system considered in Section 3.1.1.

Figure 3.1.3(a-e) shows the angle and frequency resolved emission data from the arrays when pumped with nominal fluence of 4.6 mJ cm^{-2} . For the $p_y = 370, 375$ nm cases, there are clear lasing emission peaks around the Γ point diffracted order crossings corresponding to both x and y periodicities. For the other cases, the lasing action seems to happen only around the diffractive order crossing corresponding to $p_x = 380$ nm, which might be attributed to insufficient overlap with the gain medium emission spectrum. Intriguing lasing dynamics are apparent from Figure 3.1.4, which shows experimental array emission spectra at $k_y = 0$ for series of pump fluences: around the diffracted order crossing corresponding to the $p_x = 380$ nm periodicity, there appear several lasing peaks (up to three clearly distinguishable ones), the occurrence of which depends on pump fluence. The origin of these peaks can be elucidated using MSTMM.

Figure 3.1.3(f-j) shows lattice band diagrams obtained using the Beyn's algorithm. The wavelengths on the horizontal axes come from the real parts of the eigenfrequencies found, whereas the imaginary part of the frequencies determines the color of the data points. The lasing patterns follow some of the high quality lattice modes from the simulation; interestingly, this covers also the cases where the lasing output reaches over a broader interval in k_y . (This phenomenon seems to be related to the nanoparticle material; with silver or golden nanoparticles, at the Γ point one typically sees a sharp drop in the magnitude of the simulated eigenfrequency imaginary part, and also experimental output is clearly directed in the normal direction, except for the size-dependent beam deviation due to the dark modes mentioned in Section 3.1.1.)

Figure 3.1.4 shows the emission spectra at the normal direction for several

Figure 3.1.1. Observed lasing in Ag nanoparticle square array. (a–b) Measured emission intensities with (a) $P = 1.3P_{\text{th}}$ and (b) $P = 2.0P_{\text{th}}$ where P_{th} is the threshold pump fluence for the higher energy lasing mode. (c) Emission spectra and (d) the mode energies of the lower (red) and higher (blue) energy modes as a function of pump fluence. (e–f) Examples of real space images of the lower (e) and upper (f) energy lasing modes. All the measurements are performed with a linear polarisation filter in the y direction in the output. *Reproduced from Publication II (CC BY 4.0).*

Figure 3.1.2. Simulated real and k -space patterns for constant driving with different multipole polarisations, according to the irreps of D_{4h} . The electric xy quadrupole pattern $\mathbf{v}_{12-2} - \mathbf{v}_{12+2}$ is in the A'_2 irrep (third row) and electric xy (circularly polarised) dipole pattern $\mathbf{v}_{11-1}$ in the E' irrep (ninth row). The first four columns depict the induced x and y dipoles; the last three columns show the radiated far field patterns with x , y , and x - y diagonal linear polarisation filters on the output.

values of pump fluence, with horizontal and vertical linear polarisation filter on the output, and also without filter. The emission peaks can be attributed to the simulated lattice modes based on their frequencies and also their polarisations. At the Γ point, the B'_2/B'_1 irreducible representations of D_{2h} (which merge into the two-dimensional irrep E' in the square-symmetric case D_{4h}) correspond to predominantly electric x/y dipole excitations, respectively, and will appear as x/y linearly polarised emission in the far field. The broad higher frequency peaks appearing near the diffracted order crossings corresponding to $p_y = 370, 375$ nm can be unambiguously attributed to the corresponding B'_2 mode due to its linear polarisation in the x direction, the frequencies also fitting well to the B'_2 frequencies obtained in the lattice mode simulations.

The emission around 580 nm, which is seen in all cases, is harder to analyse without more detailed polarisation measurements, as most of the irreducible representations are expected to emit far-field radiation with both x and y field components. Based on the lattice mode simulations, it is likely that the z -dipole dominated A''_2 mode is lasing, but due to the multiple peaks visible in some cases it is possible that other modes are lasing as well. For $p_y = 380$ nm, the smaller additional peak at the diffraction order crossing most probably belongs to the E' irrep.

Remark 9. In Figure 3(f-j) of Publication III the approach was slightly different: as the eigensolver was not implemented in QPMS back then, we used only the “dips” in the lowest singular values of the left-hand side of the lattice mode equation (2.6.1) as the estimates for the lattice modes, scanning over the real frequencies only. Such an approach has some obvious weaknesses: the width or depth of such dip does not exactly correspond to the quality of the modes, and even more importantly, the presence of a dip in the singular values for real ω *does not guarantee the existence of a mode*, that is, an actual solution to Eq. (2.6.1), in the complex frequency plane.

Moreover, spherical particles were used in the original simulations in Publication III, as the scuff-tmatrix results were known to be unreliable due to the bug mentioned in Subsection 2.3.3, and the T-matrix generator for cylindrical scatterers was not implemented in QPMS yet. However, one always has to be cautious about the particle shape and size, as the lattice modes are often sensitive to these parameters, and the real nanoparticles in the experimental sample do never perfectly match the nominal parameters.

And finally, in Publication III we a priori excluded the modes which have zero in-plane components at the $z = 0$ plane (i.e. the modes which are odd with respect to the $z = 0$ mirror symmetry). This was based on the incorrect assumption that such modes do not significantly radiate in the normal direction (or with small deviations from it). Far field simulations, performed later, disproved this assumption.

Figure 3.1.3. (a–e) Angle- and wavelength-resolved emission data for Ni nanoparticle array samples with p_y ranging from 370 to 390 nm at a pump fluence of 4.6 mJ cm^{-2} . (f–j) Dispersions of respective ideal infinite arrays computed using MSTMM and Beyn’s algorithm; the color scale indicates the ratio between imaginary and real frequencies of the lattice modes found as a measure of the mode quality.

Figure 3.1.4. Pump fluence dependence of emission spectra for Ni nanoparticle array samples at the Γ point. Polarisation of the lasing output is analysed with an x filter (left column), y filter (centre column), or no filter (right column). Reproduced from the supplementary material of [Publication III](#).

3.1.3 K point lasing in a honeycomb array

Publication IV is a study of lasing in a mode outside of the center of the Brillouin zone, where we extensively use the symmetry machinery, which itself is explained in detail in Section 4 of Publication I.

The investigated system is an array of golden cylindrical-shaped nanoparticles (nominal height $h = 50$ nm, base radius $r = 50$ nm) arranged into a honeycomb structure with two particles per unit cell and $p6m$ space group symmetry. The nominal distance between neighbouring particles is varied between 569–583 nm. This time, the sample is put into a liquid with a refractive index matching that of the surface (nominally $n = 1.52$), so the assumption about homogeneous background and mirror symmetry in the lattice plane is expected to hold much better than in the cases presented above.

We study lasing around the third diffracted order crossing at the K points, see 3.1.5(a–b). Organic dye IR-792 is used for the gain medium to match the frequency of the diffracted order crossing. In contrast to the previously presented cases (first diffracted order crossing at the Γ point) where the light left the array (up to small deviation due to the array size effects) in the normal direction, now there are 12 natural beam directions corresponding to the 12 diffracted orders meeting at the K points that have enough energy to leave the array plane, and the 6 of those beams closer to the normal direction are further investigated to identify the lasing mode.

The geometrical character of the possible modes is given by the irreducible representations of the point group D_{3h} , which corresponds to the little co-group of $p6m$ at the K point. The dipole excitations in the three irreducible representations A'_1, A'_2, E' that are even with respect to the $z = 0$ mirror plane, are depicted in Figure 3.1.5(c). In contrast to the Γ point lasing cases studied previously, all of these modes do have dipole contributions from at least either particle in the unit cell and are fundamentally bright, and the innermost beams are expected to have linear polarisation patterns.

We take two different paths to identify the lasing mode, both based on polarisation-resolved measurements of the emitted light and comparing the measurements to the theory. The first, “far-field” approach is based on measuring the polarisation of the six beams (Figure 3.1.6) which is compared to the theoretically predicted polarisations of the respective modes, see Figure 3.1.5(d). The second, “near-field” approach makes use of real space microscopic images of the lasing arrays—choosing a sufficient amount of different polarisation filter directions on the output, each irreducible representation will have a characteristic predicted set of patterns. The characteristic patterns can be deduced by superposition of Airy disks generated by the dipole configurations of the respective irreps; comparing the real space with the predicted patterns, one can classify the observed mode (Figure 3.1.7; see also Supplemental Material of Publication IV). In our case, both methods show that the observed lasing mode belongs to the A_1 irreducible representation.

Figure 3.1.5. (a) Schematic of the lasing measurements with hexagonal arrays. (inset: scanning electron microscope image of the gold nanoparticles, scale bar 500 nm, with the A and B unit cell sites marked) enables lasing emission in distinct off-normal angles, six of which are collected by an objective and further analysed. In the Fourier image, the six angles correspond to lasing at the six K points of the first Brillouin zone, with distinct polarisation direction of the electric field $\mathbf{E}$. (b) Band structure of the empty lattice model, as given by the by diffracted orders of a periodic structure without an actual coupling to the nanoparticles. The lasing is observed near the third diffracted order crossing at the K points (highlighted). (c) Real space electric dipole polarisations of the nanoparticles (circles) corresponding to two singlet modes and a doublet mode at a specific time. The dipole polarizations depicted by orange and magenta arrows evolve in time rotating clockwise and counterclockwise, respectively, for the K mode, and in the opposite directions for the K' mode. (d) Fourier transform of the dipole polarizations in the corresponding eigenmodes. *Reproduced from Publication IV.*

Remark 10. To make the analysis complete, one would need to rule out also the other three irreducible representations A_1'', A_2'', E'' that are odd with respect to the horizontal mirror symmetry, which was not properly done in [Publication IV](#). In fact, the measured k -space polarisation pattern could be attributed also to the A_2'' irrep as well. However, according to my preliminary survey, these irreps should be brightest on the edges of their real space images, which is not the case here (see Figure S7 from the Supplemental Material), supporting the conclusion from [Publication IV](#).

Interestingly, the lasing is observed in the two distinct K points (with opposite sign of the Bloch vector $\mathbf{k}$) simultaneously, with apparent correlation in phase, although these modes in the opposite K points are independent in the infinite lattice case. This might possibly be yet another finite size effect.

Figure 3.1.6. Lasing mode polarization. (a) Angle resolved emission of the sample without any polarizer in detection. All six K points are clearly visible. (b) Polar emission intensities at each K point in the presence of a linear polarizer. The angles refer to the polarizer angles and the radii refer to the measured intensities. The red dashed lines are the calculated intensity distributions for linearly polarized light (along the Γ - K directions) passing through the polarizer at the corresponding angle. *Reproduced from Publication IV.*

Figure 3.1.7. Upper row: examples of real-space images for different polarization filters (no filter, horizontal, vertical) used for the analysis of the lasing mode. In each case, the expected positions of the nanoparticles (small yellow-cyan circles) and the dipole polarizations (arrows) of the singlet mode A'_1 for the ideal (zero) relative phase between the two K points are depicted, projected to the corresponding filter direction. Lower row: theoretical prediction for the intensities for the ideal relative phase between the two distinct K -points. The scale bar length is $1\ \mu\text{m}$. *Reproduced from Publication IV.*

3.2 Lattice size and quality factor

One of the finite size effects in nanoparticle arrays are the radiative losses, which are unavoidable even if there were a corresponding dark mode in the infinite periodic case. The relative radiative losses are expected to decrease with increasing size of the array. One can study this phenomenon using MSTMM and Beyn’s algorithm.

Let us consider a square array of $n \times n$ nanoparticles. We choose the same basic lattice properties as in the system from Section 3.1.1, i.e. constant background refractive index of 1.52 and lattice constant of 375 nm. However, now we instead use spherical nanoparticles with radius $r = 40$ nm and constant real refractive index of 3. The choice of real refractive index eliminates absorption, and the choice of spherical scatterers is expected to eliminate possible artificial loss or gain caused by the multipole truncation (as the T-matrix of a sphere is diagonal [41, 46]).

Let us vary the array size parameter n and observe how it affects resonant frequencies of the array. Using Beyn’s algorithm, we search for the resonances (defined as the solutions of Eq. (2.4.4) with zero right-hand side) around the frequency corresponding to the first diffraction order crossing in the Γ point.

The results are displayed on top of Fig. 3.2.1. Note that the integration contours inside which the resonances are searched differ for various n , as they have to be adjusted according to the regions in which the frequencies are actually located.

As we eliminated absorption in our system, the imaginary parts of the resonant frequencies are due to the radiative losses. As expected, the larger the array, the closer to the real axis resonances can be found. Also the total number of solutions grows with the number of degrees of freedom. One can see “band-like” resonance clusters forming on top, below which lower quality resonances are scattered in a more random fashion – presumably, a lot more of them would be found if we extended or moved the integration contours lower.

If we define the quality factor of a resonance as the ratio of the real and imaginary parts of its frequency, $Q = |\Re\omega/\Im\omega|$, we can extract its dependence on the array size from the data. This is done in the bottom plot of 3.2.1, which shows a log-log plot of the quantiles of Q . The lower quantiles are strongly affected by the choice of the integration contour, but they are not important as we are mostly interested in the high quality resonances that are expected to approach the periodic system lattice modes as the array size increases. The bulk of the higher-quality resonances can be seen in the upper quantiles. There is no obvious simple function to fit to the data, but the quality factor of the upper quantiles does exhibit slightly faster than linear growth w.r.t. n , which justifies the estimate used in Section 1.2.3.

Figure 3.2.1. Resonances of nonabsorbing finite square arrays of spherical nanoparticles, as obtained with Beyn’s algorithm. Top: real and imaginary parts of the frequencies found; the integration contours used for the different sizes are indicated (ellipses in the background). Bottom: log-log plot of quality factor quantiles of the resonances found, as a function of array size; the dotted line represents exact linear proportionality.

4. Combining T-matrix method with quantum physics

Quantum effects can appear on different levels of a nanoplasmonic system. For example, if a scatterer is sufficiently small ($\lesssim 10$ nm in diameter), its scattering properties can no longer be reliably modelled by the classical approaches based on its shape and macroscopic constitutive relations. Or a small distance between two metallic nanoparticles might cause quantum tunneling of electrons (this is sometimes called the *conductive coupling regime*) or the non-local effect, drastically affecting the overall electromagnetic response of the adjacent nanoparticles. Such cases require some type of treatment that takes into account the microscopic electronic structure of the scatterer [59–62]. Here we look at quantum physics in plasmonic systems from a different viewpoint, taking the *macroscopic* description of plasmonic nanostructures as the initial point for quantisation. While these models usually *do not* give insight into the effects that come from the microscopic electronic structure, such as tunneling and the non-local effect, they *do* provide a way to describe couplings between quantum emitters e.g. placed near a plasmonic nanoarray, and they *do* explain other quantum phenomena, such as Casimir-Polder or van der Waals forces [63].

4.1 MSTMM in macroscopic QED

There have been several approaches to quantise macroscopic electromagnetism in dissipative inhomogeneous media [64–74] and the question of correctness of the different approaches is a controversial topic [75] that is beyond the scope of the present work.

One class of the quantisation approaches is based on models of microscopic harmonic oscillator continuum for the medium, coupled to the electromagnetic field. The Hamiltonian of the system is then expressed, via a Fano-diagonalisation procedure [76–78], in terms of polaritonic operators $\hat{\mathbf{f}}(\mathbf{r}, \omega)$ representing the elementary excitations of the system

$$H_{\text{macro}} = \int d^3\mathbf{r} \int_0^\infty d\omega \hbar\omega \hat{\mathbf{f}}^\dagger(\mathbf{r}, \omega) \hat{\mathbf{f}}(\mathbf{r}, \omega). \quad (4.1.1)$$

Importantly, the quantum electric field operator can be expressed using the

Green's function of the classical system [79, eq. (3)]

$$\hat{\mathbf{E}}(\mathbf{r}) = i\sqrt{\frac{\hbar}{\pi\epsilon_0}} \int_0^\infty d\omega \frac{\omega^2}{c^2} \int d^3\mathbf{r}' \sqrt{\Im\epsilon(\mathbf{r}', \omega)} \overleftrightarrow{G}(\omega; \mathbf{r}, \mathbf{r}') \hat{\mathbf{f}}(\mathbf{r}', \omega) + \text{h.c.} \quad (4.1.2)$$

As demonstrated in Section 2.4.2, MSTMM provides a suitable way to calculate the classical Green's function $\overleftrightarrow{G}(\omega; \mathbf{r}, \mathbf{r}')$ in a system consisting of compact scatterers. Model 4.1.1 can be taken as a basis for introduction of quantum emitters (QEs) into the system: for example a set of n two-level QEs can be plugged in via fermionic operators $\hat{S}_i^{\pm, z}$ that interact via the dipolar interaction with the electric field to yield [79, eq. (4)]

$$H = \int d^3\mathbf{r} \int_0^\infty d\omega \hbar\omega \hat{\mathbf{f}}^\dagger(\mathbf{r}, \omega) \hat{\mathbf{f}}(\mathbf{r}, \omega) + \sum_{i=1}^n \hbar\epsilon_i \left(\hat{S}_i^z + \frac{1}{2} \right) - \sum_{i=1}^n \left(\hat{S}_i^- + \hat{S}_i^+ \right) \boldsymbol{\mu}_i \cdot \hat{\mathbf{E}}(\mathbf{r}_i) \quad (4.1.3)$$

with

$$\hat{S}_i^+ = |g_i\rangle \langle e_i|, \hat{S}_i^- = |e_i\rangle \langle g_i|, \hat{S}_i^z = \frac{1}{2} (|e_i\rangle \langle e_i| - |g_i\rangle \langle g_i|), \quad (4.1.4)$$

where $|g_i\rangle, |e_i\rangle$ are the ground and excited states of the i -th quantum emitter.

The validity of the (widely used) Fano diagonalisation approach leading to the expressions above has been recently questioned, especially in the case when the background is empty space, i.e. $\epsilon_{\text{bg}} = \epsilon_0$ [75]. As a note for future research, it might be possible to use the building blocks of MSTMM (as introduced in Chapter 2) in the very process of canonical quantisation. The typical approach has been to use plane waves in the initial quantisation of free electromagnetic fields; however, another bases, such as regular VSWFs, could be used as well. This, as a modification of the approach used in [64], could lead to a quantum MSTMM description without the problems pointed out in [75].

4.2 Interactions between quantum emitters near a nanoparticle

In Publication V, we used two different approaches to model a system of dipolar quantum emitters (QEs), such as fluorescent molecules, in the vicinity of a plasmonic nanoparticle.

The first approach is a simplistic “relaxed Dicke” model with Hamiltonian of the form

$$\begin{aligned} H = & H_{\text{res}} + H_{\text{QE}} + H_{\text{res-QE}} + H_{\text{QE-QE}} \\ & \sum_{\lambda} \hbar\omega_{\lambda} \hat{b}_{\lambda}^{\dagger} \hat{b}_{\lambda} + \sum_i \hbar\epsilon_i \left(\hat{S}_i^z + \frac{1}{2} \right) + \sum_{i,\lambda} V_{i\lambda} \left(\hat{b}_{\lambda}^{\dagger} \hat{S}_i^- + \hat{S}_i^+ \hat{b}_{\lambda} \right) \\ & + \sum_{i < j} g_{ij} \left(\hat{S}_i^+ \hat{S}_j^- + \hat{S}_j^+ \hat{S}_i^- \right), \end{aligned} \quad (4.2.1)$$

where the nanoparticle is represented by a collection of resonant bosonic modes (indexed by λ) with frequency $\hbar\omega_{\lambda}$ and creation and annihilation operators

Figure 4.2.1. Example configurations (a) of quantum emitters in the equatorial plane of a spherical nanoparticle (with dipoles oriented perpendicular to the plane), the corresponding energy spectra obtained from the modified Dicke model for varying nanoparticle resonance (b), and light spectra from the multiple scattering model (blue lines, arbitrary units) together with the expectation values of the observable $\hat{P}$ for the energy eigenstates (red dots, arbitrary units) for the nanoparticle dipole resonance set at 2.8 eV (c), 3.0 eV (d) and 3.2 eV (e). The single QEs have transition energy $\hbar\epsilon = 3.0$ eV and dipole moment (c,d,e) $\mu = 0.19$ eV $\cdot$ nm = 9.1 D; the Drude damping is set to $\hbar\gamma_D = 1$ meV. Rows (f,g) are analogous to rows (b,d) but for a lower dipole moment $\mu = 0.04$ eV $\cdot$ nm = 1.9 D. *Reproduced from Publication V.*

$\hat{b}_\lambda^\dagger, \hat{b}_\lambda$, and the QEs are represented by two-level systems with energy level difference $\hbar\epsilon$, with fermionic operators $\hat{S}_i^{\pm,z}$ as in (4.1.4). The couplings g_{ij} , $V_{i\lambda}$ depend on system geometry, QEs' dipole moments $\boldsymbol{\mu}$ and nanoparticle resonant modes' field strengths. The model is based on extensive use of quasistatic approximation, which in turn relies on the neglect of coupling to external radiative modes and other loss channels. We solve for the energies of H via exact diagonalisation. The expression (4.2.1) assumes the rotating wave approximation, which provides an a priori block-diagonalisation of H with respect to the “total excitation number” N .

If we limit ourselves to the single-excitation states, $N = 1$, the interactions typically (depending on the actual geometry) lead to splitting of the originally constant QE energy level $\hbar\epsilon$ into a broader “band” of QE collective states and—if only one bosonic mode is considered with its energy close to the two-level system energy step $\omega \simeq \epsilon$ —to creation of two “polariton” states; this is best seen when the bosonic mode energy ω is varied, keeping the other parameters constant, see Fig. 4.2.1(f). We are interested in the radiative properties of these states – however, Hamiltonian (4.2.1) does not include coupling to the external radiation. Therefore, we couple the system to a continuum of external transverse EM field modes and apply the Fermi's golden rule to calculate the decay rate of the eigenstates of H . We show that the decay rate is proportional to the expectation value of an operator $\hat{P}$ related to the total dipole moment of the QEs and the nanoparticle, which allows to quantify the “brightness” of the calculated eigenstates of H .

Numerical experiments with varying different parameters show that when the direct QE dipole-dipole couplings g_{ij} are reasonably small, the two outstanding “polariton” states show much larger expectation value of $\hat{P}$ (therefore being “bright”) compared to the other (“dark”) states. However, the situation changes when the dipole-dipole couplings g_{ij} are large, either due to the large dipole moment of the QEs or small distance between them: some of the originally “dark” states attain a considerable expectation value of $\hat{P}$.

We benchmark the “relaxed Dicke” model results to the macroscopic QED model (4.1.3); following [79, Eq. (5)], we calculate the light spectrum $S(\mathbf{R}, \omega)$ measured at a detector position $\mathbf{R}$ if the QEs are incoherently pumped; see also [80, Supplemental Material eq. (29)].

The relevant quantities $\langle \hat{P} \rangle$, $S(\mathbf{R}, \omega)$ of the respective methods are compared on several configurations in Figure 4.2.1(c–e). The system consists of several QEs in a vicinity of a nanosphere with radius $r = 7$ nm, relative permittivity approximated by the Drude model $\epsilon(\omega) = \epsilon_\infty - \omega_p^2 / (\omega^2 + i\omega\gamma_p)$ with parameters $\epsilon_\infty = 4.6$, $\hbar\gamma_p = 0.001$ eV (this parameter is chosen arbitrarily low in order to create peaks comparable to the modified Dicke model spectrum), and we vary its plasma frequency ω_p . There is a clear qualitative correspondence between the peaks of the spectra $S(\mathbf{R}, \omega)$ in the macroscopic QED model and the $\omega, \langle \hat{P} \rangle$ pairs of the relaxed Dicke model, although the values do not fit exactly, which is expected due to the naive character of the relaxed Dicke model

and the amount of approximations made. Clearly, the correspondence would worsen with increasing γ_p .

The main goal of Publication V was to find out whether (and under which conditions) the dipole-dipole interactions between the QEs can have experimentally observable effects; i.e. whether the result into “bright” collective modes (other than the two “polaritons”) that can be probed by external radiation. Using the relaxed Dicke model, we randomly generated a large number of configurations with different parameters and followed a set of statistics related to the eigenstates. We concluded that in order to get any observable effects, one would need QEs with such large concentration and/or transition dipole moments that can be considered to be near the threshold of physical feasibility. Recently, there have been important advances in studies of phenomena arising from the dipole-dipole interactions between the emitters such as suppression of scattering [81].

Conclusion

This thesis has demonstrated that MSTMM is the method of choice for simulating optical response and mode properties of systems of compact nanoscatterers, plasmonic nanoparticle arrays in particular.

Publication I presents three notable improvements of MSTMM that largely broaden the set of problems it can solve: 1) the exponentially convergent lattice summation (newly also for 1d lattices with offset), enabling simulations with infinite periodic arrays; 2) employing group theory to transform the scattering problem into the symmetry-adapted basis; and 3) using contour-integration based nonlinear eigensolvers. The possibility to use the same method for finite and infinite systems alike is important, as it enables to study the finite size effects without contaminating the data by numerical discrepancies possibly emerging just due to different methods used. The symmetry-adapted basis transformation brings a significant decrease in the computational demands and gives additional insight in the mode analysis. Finally, nonlinear eigensolvers allow for efficient finding of the lattice modes in the complex frequency space. These improvements have been implemented as part of the QPMS suite, a free software implementation of MSTMM written from scratch as a part of the work presented here.

In Chapter 3 and Publications II, III and IV, MSTMM was proven to be a powerful method not only for simulating the scattering response of plasmonic nanoparticle arrays, but (with the improvements described in Chapter 2 of this thesis and Publication I) also for explaining the mode structure and radiation patterns observed in real-world lasing experiments. It has substantially contributed in understanding the “dark” modes, also known as the optical bound states in the continuum, and their outcoupling in the realistic setup of finite-sized arrays.

As mentioned in Chapter 4, MSTMM can be applied also in macroscopic QED context where it is used to calculate the underlying Green’s functions. A possible topic for future research would be to employ the MSTMM toolbox in the very process of canonical quantisation of macroscopic electromagnetism.

There are still also some interesting open questions related to classical MSTMM for infinite periodic systems, especially about the exact physical inter-

pretation of the two-branchedness in the long-range part of the two-dimensional lattice sum, and its possible impact on the lattice modes acquired with the method.

The basic idea of MSTMM might be simple, but its implementation (even in the basic finite version) certainly is not easy; this is partly due to the complicated prefactors and the jungle of possible normalisation and phase conventions used in spherical harmonics and other building blocks of VSWFs and translation operators, and partly because even benchmarking to other tools tends to be nontrivial (especially if they contain bugs themselves). The development of the QPMS suite has taken years so far. Although not perfect, now it is mature enough and available to be used or—even better—further developed by the whole photonics community. Out of the not so many existing MSTMM implementations, many features are (to my best knowledge) unique to QPMS, such as the handling of symmetries or the nonlinear eigensolver for mode calculations. However, others are still waiting for their inclusion, be it the support of layered media [52], evaluation of fields inside the scatterers, a solver based on the multi-level fast multipole algorithm [26, 56], or any other from the TODO list available at the project home page <https://qpms.necada.org/>. Everyone is welcome to contribute!

Bibliography

- [1] NIST Digital Library of Mathematical Functions. URL: <http://dlmf.nist.gov/>.
- [2] Chia Wei Hsu, Bo Zhen, A. Douglas Stone, John D. Joannopoulos, and Marin Soljačić. Bound states in the continuum. *Nature Reviews Materials*, 1(9):1–13, July 2016. doi:10.1038/natrevmats.2016.48.
- [3] Homer Reid. SCUFF-EM, 2018. URL: <http://homerreid.dyndns.org/scuff-EM/>.
- [4] Homer Reid. Electromagnetism in the Spherical-Wave Basis, August 2016. URL: <https://homerreid.github.io/scuff-em-documentation/tex/scuffSpherical.pdf>.
- [5] V. G. Kravets, A. V. Kabashin, W. L. Barnes, and A. N. Grigorenko. Plasmonic Surface Lattice Resonances: A Review of Properties and Applications. *Chem. Rev.*, 118(12):5912–5951, June 2018. doi:10.1021/acs.chemrev.8b00243.
- [6] Weijia Wang, Mohammad Ramezani, Aaro I. Väkeväinen, Päivi Törmä, Jaime Gómez Rivas, and Teri W. Odom. The rich photonic world of plasmonic nanoparticle arrays. *Materials Today*, 21(3):303–314, April 2018. doi:10.1016/j.mattod.2017.09.002.
- [7] V. Giannini, G. Vecchi, and J. Gómez Rivas. Lighting Up Multipolar Surface Plasmon Polaritons by Collective Resonances in Arrays of Nanoantennas. *Phys. Rev. Lett.*, 105(26):266801, December 2010. doi:10.1103/PhysRevLett.105.266801.
- [8] G. Vecchi, V. Giannini, and J. Gómez Rivas. Shaping the Fluorescent Emission by Lattice Resonances in Plasmonic Crystals of Nanoantennas. *Phys. Rev. Lett.*, 102(14):146807, April 2009. doi:10.1103/PhysRevLett.102.146807.
- [9] F. J. García de Abajo. Colloquium: Light scattering by particle and hole arrays. *Rev. Mod. Phys.*, 79(4):1267–1290, October 2007. doi:10.1103/RevModPhys.79.1267.
- [10] Shengli Zou, Nicolas Janel, and George C. Schatz. Silver nanoparticle array structures that produce remarkably narrow plasmon lineshapes. *The Journal of Chemical Physics*, 120(23):10871–10875, June 2004. doi:10.1063/1.1760740.
- [11] Mohammadreza Khorasaninejad and Federico Capasso. Metalenses: Versatile multifunctional photonic components. *Science*, 358(6367), December 2017. doi:10.1126/science.aam8100.
- [12] A. I. Väkeväinen, R. J. Moerland, H. T. Rekola, A.-P. Eskelinen, J.-P. Martikainen, D.-H. Kim, and P. Törmä. Plasmonic Surface Lattice Resonances at the Strong Coupling Regime. *Nano Lett.*, 14(4):1721–1727, April 2014. doi:10.1021/nl4035219.

- [13] Mohammad Ramezani, Matthijs Berghuis, and Jaime Gómez Rivas. Strong light-matter coupling and exciton-polariton condensation in lattices of plasmonic nanoparticles [Invited]. *J. Opt. Soc. Am. B, JOSAB*, 36(7):E88–E103, July 2019. doi:10.1364/JOSAB.36.000E88.
- [14] P. Törmä and W. L. Barnes. Strong coupling between surface plasmon polaritons and emitters: A review. *Rep. Prog. Phys.*, 78(1):013901, 2015. doi:10.1088/0034-4885/78/1/013901.
- [15] Wei Zhou, Montacer Dridi, Jae Yong Suh, Chul Hoon Kim, Dick T. Co, Michael R. Wasielewski, George C. Schatz, and Teri W. Odom. Lasing action in strongly coupled plasmonic nanocavity arrays. *Nat Nano*, 8(7):506–511, July 2013. doi:10.1038/nnano.2013.99.
- [16] T. K. Hakala, H. T. Rekola, A. I. Väkeväinen, J.-P. Martikainen, M. Nečada, A. J. Moilanen, and P. Törmä. Lasing in dark and bright modes of a finite-sized plasmonic lattice. *Nature Communications*, 8:13687, January 2017. doi:10.1038/ncomms13687.
- [17] R. Guo, M. Nečada, T. K. Hakala, A. I. Väkeväinen, and P. Törmä. Lasing at Γ Points of a Honeycomb Plasmonic Lattice. *Phys. Rev. Lett.*, 122(1):013901, January 2019. doi:10.1103/PhysRevLett.122.013901.
- [18] Tommi K. Hakala, Antti J. Moilanen, Aaro I. Väkeväinen, Rui Guo, Jani-Petri Martikainen, Konstantinos S. Daskalakis, Heikki T. Rekola, Aleksu Julku, and Päivi Törmä. Bose–Einstein condensation in a plasmonic lattice. *Nature Phys*, 14(7):739–744, July 2018. doi:10.1038/s41567-018-0109-9.
- [19] Ankun Yang, Thang B. Hoang, Montacer Dridi, Claire Deeb, Maiken H. Mikkelsen, George C. Schatz, and Teri W. Odom. Real-time tunable lasing from plasmonic nanocavity arrays. *Nat Commun*, 6:6939, April 2015. doi:10.1038/ncomms7939.
- [20] Mohammad Ramezani, Alexei Halpin, Antonio I. Fernández-Domínguez, Johannes Feist, Said Rahimzadeh-Kalaleh Rodriguez, Francisco J. Garcia-Vidal, and Jaime Gómez Rivas. Plasmon-exciton-polariton lasing. *Optica, OPTICA*, 4(1):31–37, January 2017. arXiv:1606.06866, doi:10.1364/OPTICA.4.000031.
- [21] Aaro I. Väkeväinen, Antti J. Moilanen, Marek Nečada, Tommi K. Hakala, Konstantinos S. Daskalakis, and Päivi Törmä. Sub-picosecond thermalization dynamics in condensation of strongly coupled lattice plasmons. *Nature Communications*, 11(1):1–12, June 2020. arXiv:1905.07609, doi:10.1038/s41467-020-16906-1.
- [22] Danqing Wang, Weijia Wang, Michael P. Knudson, George C. Schatz, and Teri W. Odom. Structural Engineering in Plasmon Nanolasers. *Chem. Rev.*, 118(6):2865–2881, March 2018. doi:10.1021/acs.chemrev.7b00424.
- [23] M. Kataja, T. K. Hakala, A. Julku, M. J. Huttunen, S. van Dijken, and P. Törmä. Surface lattice resonances and magneto-optical response in magnetic nanoparticle arrays. *Nat Commun*, 6, May 2015. doi:10.1038/ncomms8072.
- [24] Christian Kuttner. Plasmonics in Sensing: From Colorimetry to SERS Analytics. *Plasmonics*, November 2018. doi:10.5772/intechopen.79055.
- [25] Gerhard Kristensson. *Scattering of Electromagnetic Waves by Obstacles*. Scitech Publishing, Edison, NJ, July 2016.
- [26] Weng Cho Chew, Jian-Ming Jin, Eric Michielssen, and Jiming Song. *Fast and Efficient Algorithms in Computational Electromagnetics*. Artech House Antennas and Propagation Library. Artech House Publishers, 2000.
- [27] Kane Yee. Numerical solution of initial boundary value problems involving maxwell’s equations in isotropic media. *IEEE Transactions on Antennas and Propagation*, 14(3):302–307, May 1966. doi:10.1109/TAP.1966.1138693.

- [28] John B. Schneider. Understanding the FDTD Method, 2020. URL: <https://www.eecs.wsu.edu/~schneidj/ufdtd/>.
- [29] Dennis M. Sullivan. *Electromagnetic Simulation Using the FDTD Method*. Wiley-IEEE Press, Hoboken, New Jersey, 2 edition edition, June 2013.
- [30] S. Pernet, X. Ferrieres, and G. Cohen. High spatial order finite element method to solve Maxwell's equations in time domain. *IEEE Transactions on Antennas and Propagation*, 53(9):2889–2899, September 2005. doi:10.1109/TAP.2005.856046.
- [31] Bernardo Cockburn, George E. Karniadakis, and Chi-Wang Shu. *Discontinuous Galerkin Methods: Theory, Computation and Applications*. Springer Science & Business Media, December 2012.
- [32] Stéphane Descombes, Clément Durochat, Stéphane Lanteri, Ludovic Moya, Claire Scheid, and Jonathan Viquerat. Recent advances on a DGTD method for time-domain electromagnetics. *Photonics and Nanostructures - Fundamentals and Applications*, 11(4):291–302, November 2013. doi:10.1016/j.photonics.2013.06.005.
- [33] Shih-Hui Chang, Hui Cao, and A. Taflove. Four-level two-electron FDTD model of lasing action in a semiconductor. In *IEEE Antennas and Propagation Society International Symposium. Digest. Held in Conjunction with: USNC/CNC/URSI North American Radio Sci. Meeting (Cat. No.03CH37450)*, volume 4, pages 382–385 vol.4, June 2003. doi:10.1109/APS.2003.1220201.
- [34] Montacer Dridi and George C. Schatz. Lasing action in periodic arrays of nanoparticles. *Journal of the Optical Society of America B*, 32(5):818, May 2015. doi:10.1364/JOSAB.32.000818.
- [35] J. Cuerda, F. Rütting, F. J. García-Vidal, and J. Bravo-Abad. Theory of lasing action in plasmonic crystals. *Phys. Rev. B*, 91(4):041118, January 2015. doi:10.1103/PhysRevB.91.041118.
- [36] R. Lee and A.C. Cangellaris. A study of discretization error in the finite element approximation of wave solutions. *IEEE Transactions on Antennas and Propagation*, 40(5):542–549, May 1992. doi:10.1109/8.142629.
- [37] Karl F. Warnick. *Numerical Analysis for Electromagnetic Integral Equations*. Artech House Electromagnetic Analysis. Artech House, 2008.
- [38] LinLin Zhao, K. Lance Kelly, and George C. Schatz. The Extinction Spectra of Silver Nanoparticle Arrays: Influence of Array Structure on Plasmon Resonance Wavelength and Width†. *J. Phys. Chem. B*, 107(30):7343–7350, July 2003. doi:10.1021/jp034235j.
- [39] P. A. Martin. *Multiple Scattering: Interaction of Time-Harmonic Waves with N Obstacles*. Encyclopedia of Mathematics and Its Applications. Cambridge University Press, first edition, 2006.
- [40] David Colton and Rainer Kress. *Integral Equation Methods in Scattering Theory*. Classics in Applied Mathematics. SIAM-Society for Industrial and Applied Mathematics, 2013.
- [41] Gustav Mie. Beiträge zur Optik trüber Medien, speziell kolloidaler Metallösungen. *Ann. Phys.*, 330(3):377–445, January 1908. doi:10.1002/andp.19083300302.
- [42] P. C. Waterman. New Formulation of Acoustic Scattering. *The Journal of the Acoustical Society of America*, 45(6):1417–1429, June 1969. doi:10.1121/1.1911619.
- [43] P. C. Waterman. Symmetry, Unitarity, and Geometry in Electromagnetic Scattering. *Phys. Rev. D*, 3(4):825–839, February 1971. doi:10.1103/PhysRevD.3.825.

- [44] Michael I. Mishchenko, Larry D. Travis, and Andrew A. Lacis. *Scattering, Absorption, and Emission of Light by Small Particles*. Cambridge University Press, first edition, 2002.
- [45] M. T. Homer Reid and Steven G. Johnson. Efficient Computation of Power, Force, and Torque in BEM Scattering Calculations. *IEEE Transactions on Antennas and Propagation*, 63(8):3588–3598, August 2015. [arXiv:1307.2966](#), [doi:10.1109/TAP.2015.2438393](#).
- [46] F. Michael Schulz, Knut Stamnes, and J. J. Stamnes. Point-group symmetries in electromagnetic scattering. *Journal of the Optical Society of America A*, 16(4):853, April 1999. [doi:10.1364/JOSAA.16.000853](#).
- [47] P. B. Johnson and R. W. Christy. Optical Constants of the Noble Metals. *Phys. Rev. B*, 6(12):4370–4379, December 1972. [doi:10.1103/PhysRevB.6.4370](#).
- [48] Aleksandar D. Rakić, Aleksandra B. Djurišić, Jovan M. Elazar, and Marian L. Majewski. Optical properties of metallic films for vertical-cavity optoelectronic devices. *Appl. Opt.*, *AO*, 37(22):5271–5283, August 1998. [doi:10.1364/AO.37.005271](#).
- [49] Wolf-Jürgen Beyn. An integral method for solving nonlinear eigenvalue problems. *Linear Algebra and its Applications*, 436(10):3839–3863, May 2012. [doi:10.1016/j.laa.2011.03.030](#).
- [50] Brendan Gavin, Agnieszka Międlar, and Eric Polizzi. FEAST eigensolver for nonlinear eigenvalue problems. *Journal of Computational Science*, 27:107–117, July 2018. [doi:10.1016/j.jocs.2018.05.006](#).
- [51] Krzysztof M. Czajkowski, Maria Bancerek, and Tomasz J. Antosiewicz. Multipole analysis of substrate-supported dielectric nanoresonator arrays with T-matrix method. *arXiv:2006.09137 [physics]*, June 2020. URL: <http://arxiv.org/abs/2006.09137>, [arXiv:2006.09137](#).
- [52] Amos Egel, Siegfried W. Kettlitz, and Uli Lemmer. Efficient evaluation of Sommerfeld integrals for the optical simulation of many scattering particles in planarly layered media. *J. Opt. Soc. Am. A*, *JOSAA*, 33(4):698–706, April 2016. [doi:10.1364/JOSAA.33.000698](#).
- [53] M. Epton and B. Dembart. Multipole Translation Theory for the Three-Dimensional Laplace and Helmholtz Equations. *SIAM J. Sci. Comput.*, 16(4):865–897, July 1995. [doi:10.1137/0916051](#).
- [54] V. Rokhlin. Diagonal Forms of Translation Operators for the Helmholtz Equation in Three Dimensions. *Applied and Computational Harmonic Analysis*, 1(1):82–93, December 1993. [doi:10.1006/acha.1993.1006](#).
- [55] R. Coifman, V. Rokhlin, and S. Wandzura. The fast multipole method for the wave equation: A pedestrian prescription. *IEEE Antennas and Propagation Magazine*, 35(3):7–12, June 1993. [doi:10.1109/74.250128](#).
- [56] Johannes Markkanen and Alex J. Yuffa. Fast superposition T-matrix solution for clusters with arbitrarily-shaped constituent particles. *Journal of Quantitative Spectroscopy and Radiative Transfer*, 189:181–188, March 2017. [doi:10.1016/j.jqsrt.2016.11.004](#).
- [57] Johannes Markkanen. FaSTMM, 2018. available at https://bitbucket.org/planetarysystemresearch/fastmm_v1.0/. URL: https://bitbucket.org/planetarysystemresearch/fastmm_v1.0/src/master/.
- [58] Jonathan P. Dowling, Michael Scalora, Mark J. Bloemer, and Charles M. Bowden. The photonic band edge laser: A new approach to gain enhancement. *Journal of Applied Physics*, 75(4):1896–1899, February 1994. [doi:10.1063/1.356336](#).

- [59] Tuomas Rossi. *Computational Modeling of Quantum Aspects in Plasmonic Nanostructures*. Aalto University, 2017. URL: <https://aaltodoc.aalto.fi:443/handle/123456789/28074>.
- [60] Rubén Esteban, Asier Zugarramurdi, Pu Zhang, Peter Nordlander, Francisco J. García-Vidal, Andrei G. Borisov, and Javier Aizpurua. A classical treatment of optical tunneling in plasmonic gaps: Extending the quantum corrected model to practical situations. *Faraday Discuss.*, 178(0):151–183, May 2015. doi:10.1039/C4FD00196F.
- [61] Wenqi Zhu, Ruben Esteban, Andrei G. Borisov, Jeremy J. Baumberg, Peter Nordlander, Henri J. Lezec, Javier Aizpurua, and Kenneth B. Crozier. Quantum mechanical effects in plasmonic structures with subnanometre gaps. *Nature Communications*, 7(1):11495, June 2016. doi:10.1038/ncomms11495.
- [62] Da Xu, Xiao Xiong, Lin Wu, Xi-Feng Ren, Ching Eng Png, Guang-Can Guo, Qihuang Gong, and Yun-Feng Xiao. Quantum plasmonics: New opportunity in fundamental and applied photonics. *Adv. Opt. Photon., AOP*, 10(4):703–756, December 2018. doi:10.1364/AOP.10.000703.
- [63] Stefan Scheel and Stefan Yoshi Buhmann. Macroscopic Quantum Electrodynamics - Concepts and Applications. *Acta Phys. Slovaca*, 58(5):675–809, August 2008. URL: <http://www.physics.sk/aps/pub.php?y=2008&pub=aps-08-05>.
- [64] V. Dorier, J. Lampart, S. Guérin, and H. R. Jauslin. Canonical quantization for quantum plasmonics with finite nanostructures. *Phys. Rev. A*, 100(4):042111, October 2019. doi:10.1103/PhysRevA.100.042111.
- [65] Aurélien Drezet. Equivalence between the Hamiltonian and Langevin noise descriptions of plasmon polaritons in a dispersive and lossy inhomogeneous medium. *Phys. Rev. A*, 96(3):033849, September 2017. doi:10.1103/PhysRevA.96.033849.
- [66] S. A. R. Horsley and T. G. Philbin. Canonical quantization of electromagnetism in spatially dispersive media. *New J. Phys.*, 16(1):013030, January 2014. doi:10.1088/1367-2630/16/1/013030.
- [67] T. G. Philbin. Canonical quantization of macroscopic electromagnetism. *New J. Phys.*, 12(12):123008, December 2010. doi:10.1088/1367-2630/12/12/123008.
- [68] Christian Raabe, Stefan Scheel, and Dirk-Gunnar Welsch. Unified approach to QED in arbitrary linear media. *Phys. Rev. A*, 75(5):053813, May 2007. doi:10.1103/PhysRevA.75.053813.
- [69] Navin A. R. Bhat and J. E. Sipe. Hamiltonian treatment of the electromagnetic field in dispersive and absorptive structured media. *Phys. Rev. A*, 73(6):063808, June 2006. doi:10.1103/PhysRevA.73.063808.
- [70] L. G. Suttorp and A. J. van Wonderen. Fano diagonalization of a polariton model for an inhomogeneous absorptive dielectric. *EPL*, 67(5):766, September 2004. doi:10.1209/epl/i2004-10131-8.
- [71] L. G. Suttorp and Martijn Wubs. Field quantization in inhomogeneous absorptive dielectrics. *Phys. Rev. A*, 70(1):013816, July 2004. doi:10.1103/PhysRevA.70.013816.
- [72] Omar Di Stefano, Salvatore Savasta, and Raffaello Girlanda. Mode expansion and photon operators in dispersive and absorbing dielectrics. *Journal of Modern Optics*, 48(1):67–84, January 2001. doi:10.1080/09500340108235155.
- [73] Ho Trung Dung, Ludwig Knöll, and Dirk-Gunnar Welsch. Spontaneous decay in the presence of dispersing and absorbing bodies: General theory and application to a spherical cavity. *Phys. Rev. A*, 62(5):053804, October 2000. doi:10.1103/PhysRevA.62.053804.

- [74] T. Gruner and D.-G. Welsch. Green-function approach to the radiation-field quantization for homogeneous and inhomogeneous Kramers-Kronig dielectrics. *Phys. Rev. A*, 53(3):1818–1829, March 1996. doi:10.1103/PhysRevA.53.1818.
- [75] Vincent Dorier, Stéphane Guérin, and Hans-Rudolf Jauslin. Critical review of quantum plasmonic models for finite-size media. *Nanophotonics*, -1(ahead-of-print), June 2020. doi:10.1515/nanoph-2020-0061.
- [76] Kurt Friedrichs. Über die Spektralzerlegung eines Integraloperators. *Math. Ann.*, 115(1):249–272, December 1938. doi:10.1007/BF01448941.
- [77] K. O. Friedrichs. On the perturbation of continuous spectra. *Communications on Pure and Applied Mathematics*, 1(4):361–406, December 1948. doi:10.1002/cpa.3160010404.
- [78] U. Fano. Effects of Configuration Interaction on Intensities and Phase Shifts. *Phys. Rev.*, 124(6):1866–1878, December 1961. doi:10.1103/PhysRev.124.1866.
- [79] A. Delga, J. Feist, J. Bravo-Abad, and F. J. García-Vidal. Theory of strong coupling between quantum emitters and localized surface plasmons. *J. Opt.*, 16(11):114018, November 2014. doi:10.1088/2040-8978/16/11/114018.
- [80] A. Delga, J. Feist, J. Bravo-Abad, and F. J. García-Vidal. Quantum Emitters Near a Metal Nanoparticle: Strong Coupling and Quenching. *Phys. Rev. Lett.*, 112(25):253601, June 2014. doi:10.1103/PhysRevLett.112.253601.
- [81] N. J. Schilder, C. Sauvan, Y. R. P. Sortais, A. Browaeys, and J.-J. Greffet. Near-Resonant Light Scattering by a Subwavelength Ensemble of Identical Atoms. *Phys. Rev. Lett.*, 124(7):073403, February 2020. doi:10.1103/PhysRevLett.124.073403.

ISBN 978-952-64-0151-5 (printed)
ISBN 978-952-64-0152-2 (pdf)
ISSN 1799-4934 (printed)
ISSN 1799-4942 (pdf)

Aalto University
School of Science
Department of Applied Physics
www.aalto.fi

**BUSINESS +
ECONOMY**

**ART +
DESIGN +
ARCHITECTURE**

**SCIENCE +
TECHNOLOGY**

CROSSOVER

**DOCTORAL
DISSERTATIONS**